

# A Uniformization Theorem Of Complete Noncompact Kähler Surfaces With Positive Bisectional Curvature

Bing-Long Chen\*, Siu-Hung Tang\*\*, Xi-Ping Zhu\*

\* Department of Mathematics, Zhongshan University,  
Guangzhou 510275, P. R. China, and

The Institute of Mathematical Sciences, The Chinese University of Hong  
Kong, Hong Kong

\*\* The Institute of Mathematical Sciences, The Chinese University of Hong  
Kong, Hong Kong

## Abstract

In this paper, by combining techniques from Ricci flow and algebraic geometry, we prove the following generalization of the classical uniformization theorem of Riemann surfaces. Given a complete noncompact complex two dimensional Kähler manifold  $M$  of positive and bounded holomorphic bisectional curvature, suppose its geodesic balls have Euclidean volume growth and its scalar curvature decays to zero at infinity in the average sense, then  $M$  is biholomorphic to  $\mathbf{C}^2$ . During the proof, we also discover an interesting gap phenomenon which says that a Kähler manifold as above automatically has quadratic curvature decay at infinity in the average sense.

## §1. Introduction

One of the most beautiful results in complex analysis of one variable is the classical uniformization theorem of Riemann surfaces which states that a simply connected Riemann surface is biholomorphic to either the Riemann sphere, the complex line or the open unit disc. Unfortunately, a direct analog of this beautiful result to higher dimensions does not exist. For example, there is a vast variety of biholomorphically distinct complex structures on  $\mathbf{R}^{2n}$  for  $n > 1$ , a fact which was already known to Poincaré (see [3], [9] for a modern

treatment). Thus, in order to characterize the standard complex structures for higher dimensional complex manifolds, one must impose more restrictions on the manifolds.

From the point of view of differential geometry, one consequence of the uniformization theorem is that a positively curved compact or noncompact Riemann surface must be biholomorphic to the Riemann sphere or the complex line respectively. It is thus natural to ask whether there is similar characterization for higher dimensional complete Kähler manifold with positive "curvature". That such a characterization exists in the case of compact Kähler manifold is the famous Frankel conjecture which says that a compact Kähler manifold of positive holomorphic bisectional curvature is biholomorphic to a complex projective space. This conjecture was solved by Andreotti–Frankel [10] and Mabuchi [18] in complex dimensions two and three respectively and the general case was then solved by Mori [24], and Siu–Yau [34] independently. In this paper, we are thus interested in complete noncompact Kähler manifolds with positive holomorphic bisectional curvature. The following conjecture provides the main impetus.

**Conjecture** (Green–Wu [13], Siu[33], Yau [37]) A complete noncompact Kähler manifold of positive holomorphic bisectional curvature is biholomorphic to a complex Euclidean space.

In contrary to the compact case, very little is known about this conjecture. The first result in this direction is the following isometrically embedding theorem.

**Theorem** (Mok–Siu–Yau[20], Mok[21]) Let  $M$  be a complete noncompact Kähler manifold of nonnegative holomorphic bisectional curvature of complex dimension  $n \geq 2$ . Suppose there exist positive constants  $C_1, C_2$  such that

for a fixed base point  $x_0$  and some  $\varepsilon > 0$ ,

$$(i) \quad \text{Vol}(B(x_0, r)) \geq C_1 r^{2n} \quad 0 \leq r < +\infty ,$$

$$(ii) \quad R(x) \leq \frac{C_2}{1 + d^{2+\varepsilon}(x_0, x)} \quad \text{on } M ,$$

where  $\text{Vol}(B(x_0, r))$  denotes the volume of the geodesic ball  $B(x_0, r)$  centered at  $x_0$  with radius  $r$ ,  $R(x)$  denotes the scalar curvature and  $d(x_0, x)$  denotes the geodesic distance between  $x_0$  and  $x$ . Then  $M$  is isometrically biholomorphic to  $\mathbf{C}^n$  with the flat metric.

Their method is to consider the Poincaré–Lelong equation  $\sqrt{-1}\partial\bar{\partial}u = \text{Ric}$ . Under the condition (ii) that the curvature has faster than quadratic decay, they proved the existence of a bounded solution  $u$  to the Poincaré–Lelong equation. By virtue of Yau’s Liouville theorem on complete manifolds with nonnegative Ricci curvature, this bounded plurisubharmonic function  $u$  must be constant and hence the Ricci curvature must be identically zero. This implies that the Kähler metric is flat because of the nonnegativity of the holomorphic bisectional curvature. However, this argument breaks down if the faster than quadratic decay condition (ii) is weakened to a quadratic decay condition. In this case, although we can still solve the Poincaré–Lelong equation with logarithmic growth, the boundedness of the solution can no longer be guaranteed.

In [21], Mok also developed a general scheme for compactifying complete Kähler manifolds of positive holomorphic bisectional curvature. This allowed him to obtain the following improvement of the above theorem.

**Theorem** (Mok[21]) Let  $M$  be a complete noncompact Kähler manifold of complex dimension  $n$  with positive holomorphic bisectional curvature. Suppose there exist positive constants  $C_1, C_2$  such that for a fixed base point

$x_0$ ,

$$(i) \quad \text{Vol}(B(x_0, r)) \geq C_1 r^{2n} \quad 0 \leq r < +\infty ,$$

$$(ii)' \quad 0 < R(x) \leq \frac{C_2}{1 + d^2(x_0, x)} \quad \text{on } M ,$$

then  $M$  is biholomorphic to an affine algebraic variety. Moreover, if in addition the complex dimension  $n = 2$  and

$$(iii) \quad \text{the Riemannian sectional curvature of } M \text{ is positive,}$$

then  $M$  is biholomorphic to  $\mathbf{C}^2$ .

To the best of our knowledge, the above result of Mok and its slight improvements by To [35], and Chen-Zhu [7] are the best results in complex dimension two of the above stated conjecture. Here, we would also like to recall the remark pointed out in [7] that there is a gap in the proof of Shi [31] (see [7] for more explanation) which would otherwise constitute a better result than that of Mok [21].

In this paper, we consider only the case of complex dimension two. Our principal result is the following

**Main Theorem** Let  $M$  be a complete noncompact complex two-dimensional Kähler manifold of positive and bounded holomorphic bisectional curvature. Suppose there exists a positive constant  $C_1$  such that for a fixed base point  $x_0$ , we have

$$(i) \quad \text{Vol}(B(x_0, r)) \geq C_1 r^4 \quad 0 \leq r < +\infty ,$$

$$(ii)'' \quad \lim_{r \rightarrow +\infty} \frac{1}{\text{Vol}(B(x_0, r))} \int_{B(x_0, r)} R(x) dx = 0 ,$$

then  $M$  is biholomorphic to  $\mathbf{C}^2$ .

We remark that the condition (ii)'' means that the scalar curvature tends to zero at infinity in average sense. In view of the classical Bonnet–Myers theorem, this condition is almost necessary to make sure that the manifold is

noncompact under our positive bisectional curvature condition. It is also clear that the pointwise decay condition

$$\lim_{d(x,x_0) \rightarrow +\infty} R(x) = 0$$

is stronger than (ii)''.

The proof of the Main Theorem will be divided into three parts. In the first part, we will show that  $M$  is a Stein manifold homeomorphic to  $\mathbf{R}^4$ . For this, we evolve the Kähler metric on  $M$  by the Ricci flow first studied by Hamilton. Note that the underlying complex structure of  $M$  is unchanged under the Ricci flow, thus we can replace the Kähler metric in our main theorem by any one of the evolving metric. The advantage is that, in our case, the properties of the evolving metric are improving during the flow. Moreover, we know that the Euclidean volume growth condition (i) as well as the positive holomorphic bisectional curvature condition are perserved by the evolving metric. More importantly, by a blow up and blow down argument as in [6], we can prove that the curvature of the evolving metric decays linearly in time. This implies that the injectivity radius of the evolving metric is getting bigger and bigger and any geodesic ball with radius less than half of the injectivity radius is almost pseudoconvex. By a perturbation argument as in [7], we are then able to modify these geodesic balls to a sequence of exhausting pseudoconvex domains of  $M$  such that any two of them form a Runge pair. From this, it follow readily that  $M$  is a Stein manifold homeomorphic to  $\mathbf{R}^4$ .

In the second part of the proof, we consider the algebra  $P(M)$  of holomorphic functions of polynomial growth on  $M$  and we will prove that its quotient field has transcendental degree two over  $\mathbf{C}$ . For this, we first need to construct two algebraically independent holomorphic functions in the algebra  $P(M)$ . Using the  $L^2$  estimates of Andreotti–Vesentini [1] and Hörmander [17], it suffices to construct a strictly plurisubharmonic function of logarithmic growth on  $M$ . Now, if the scalar curvature decays in space at least quadratically, it

was known from [20], [21] that such a strictly plurisubharmonic function of logarithmic growth can be obtained by solving the Poincaré–Lelong equation, as we mentioned before. However, our decay assumption (ii)'' is too weak to apply their result directly. To resolve this difficulty, we make use of the Ricci flow to verify a new gap phenomenon which was already predicted by Yau in [38]. More explicitly, by using the time decay estimate of evolving metric in the previous part, we prove that the curvature of the initial metric must decay quadratically in space in certain average sense. Fortunately, this turns out to be enough to insure the existence of a strictly plurisubharmonic function of logarithmic growth. Next, by using the time decay estimate and the injectivity radius estimate of the evolving metric, we prove that the dimension of the space of holomorphic functions in  $P(M)$  of degree at most  $p$  is bounded by a constant times  $p^2$ . Combining this with the existence of two algebraically independent holomorphic functions in  $P(M)$  as above, we can prove that the quotient field  $R(M)$  of  $P(M)$  has transcendental degree two over  $\mathbf{C}$  by a classical argument of Poincaré–Siegel. In other words,  $R(M)$  is a finite extension field of some  $\mathbf{C}(f_1, f_2)$ , where  $f_1, f_2 \in P(M)$  are algebraically independent over  $\mathbf{C}$ . Then, from the primitive element theorem, we have  $R(M) = \mathbf{C}(f_1, f_2, g/h)$  for some  $g, h \in P(M)$ . Hence the mapping  $F : M \rightarrow \mathbf{C}^4$  given by  $F = (f_1, f_2, g, h)$  defines, in an appropriate sense, a birational equivalence between  $M$  and some irreducible affine algebraic subvariety  $Z$  of  $\mathbf{C}^4$ .

In the last part of proof, we will basically follow the approach of Mok in [21] and [23] to establish a biholomorphic map from  $M$  onto a quasi-affine algebraic variety by desingularizing the map  $F$ . Our essential contribution in this part is to establish uniform estimates on the multiplicity and the number of irreducible components of the zero divisor of a holomorphic function in  $P(M)$ . Again, the time decay estimate of the Ricci flow plays a crucial role in the arguments. Based on these estimates, we can show that the mapping  $F :$

$M \rightarrow Z$  is almost surjective in the sense that it can miss only a finite number of subvarieties in  $Z$ , and can be desingularized by adjoining a finite number of holomorphic functions of polynomial growth. This completes the proof that  $M$  is a quasi-affine algebraic variety. Finally, by combining with the fact that  $M$  is homeomorphic to  $\mathbf{R}^4$ , we conclude that  $M$  is indeed biholomorphic to  $\mathbf{C}^2$  by a theorem of Ramanujam [27] on algebraic surfaces.

This paper contains eight sections. From Sections 2 to 4, we study the Ricci flow and obtain several geometric estimates for the evolving metric. In Section 5, we show that the two dimensional Kähler manifold is homeomorphic to  $\mathbf{R}^4$  and is a Stein manifold. Based on the estimates on the Ricci flow, a space decay estimate on the curvature and the existence of a strictly plurisubharmonic function of logarithmic growth are obtained in Section 6. In Section 7, we establish uniform estimates on the multiplicity and the number of irreducible components of the zero divisor of a holomorphic function of polynomial growth. Finally, in Section 8 we construct a biholomorphic map from the Kähler manifold onto a quasi-affine algebraic variety and complete the proof of the Main Theorem.

We are grateful to Professor L. F. Tam for many helpful discussions and Professor S. T. Yau for his interest and encouragement. This work was partially supported by The IMS of The Chinese University of Hong Kong and the third author was also partially supported by the Foundation for Outstanding Young Scholars of China.

## §2. Preserving the volume growth

Let  $(M, g_{\alpha\bar{\beta}})$  be a complete, noncompact Kähler surface (i.e., a Kähler manifold of complex dimension two) satisfying all the assumptions in the Main Theorem. We evolve the metric  $g_{\alpha\bar{\beta}}$  according to the following Ricci flow

equation

$$\begin{cases} \frac{\partial g_{\alpha\bar{\beta}}(x, t)}{\partial t} = -R_{\alpha\bar{\beta}}(x, t) & x \in M \quad t > 0, \\ g_{\alpha\bar{\beta}}(x, 0) = g_{\alpha\bar{\beta}}(x) & x \in M, \end{cases} \quad (2.1)$$

where  $R_{\alpha\bar{\beta}}(x, t)$  denotes the Ricci curvature tensor of the metric  $g_{\alpha\bar{\beta}}(x, t)$ .

Since the curvature of the initial metric is bounded, it is known from [29] that there exists some  $T_{\max} > 0$  such that (2.1) has a maximal solution on  $M \times [0, T_{\max})$  with either  $T_{\max} = +\infty$  or the curvature becomes unbounded as  $t \rightarrow T_{\max}$  when  $T_{\max} < +\infty$ . By using the maximum principle, one knows ( see Mok [22], Hamilton [14], or Shi [32] ) that the positivity of holomorphic bisectional curvature and the Kählerity of  $g_{\alpha\bar{\beta}}$  are preserved under the evolution of (2.1). In particular, the Ricci curvature remains positive.

Our first result for the solution of the Ricci flow (2.1) is the following proposition.

**Proposition 2.1** Suppose  $(M, g_{\alpha\bar{\beta}})$  is assumed as above. Then the maximal volume growth condition (i) is preserved under the evolution of (2.1), i.e.,

$$\text{Vol}_t(B_t(x, r)) \geq C_1 r^4 \quad \text{for all } r > 0, \quad x \in M, \quad (2.2)$$

with the same constant  $C_1$  as in condition (i). Here,  $B_t(x, r)$  is the geodesic ball of radius  $r$  with center at  $x$  with respect to the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$ , and the volume  $\text{Vol}_t$  is also taken with respect to the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$ .

**Proof.** Define a function  $F(x, t)$  on  $M \times [0, T_{\max})$  as follows,

$$F(x, t) = \log \frac{\det(g_{\alpha\bar{\beta}}(x, t))}{\det(g_{\alpha\bar{\beta}}(x, 0))}.$$

By (2.1), we have

$$\begin{aligned} \frac{\partial F(x, t)}{\partial t} &= g^{\alpha\bar{\beta}}(x, t) \cdot \frac{\partial}{\partial t} g_{\alpha\bar{\beta}}(x, t) \\ &= -R(x, t) \leq 0, \end{aligned} \quad (2.3)$$



which implies that  $F(\cdot, t)$  is nonincreasing in time. Since  $R_{\alpha\bar{\beta}}(x, t) \geq 0$ , we know from (2.1) that the metric is shrinking in time. In particular,

$$g_{\alpha\bar{\beta}}(x, t) \leq g_{\alpha\bar{\beta}}(x, 0) \quad \text{on} \quad M \times [0, T_{\max}) . \quad (2.4)$$

This implies that

$$\begin{aligned} e^{F(x,t)} R(x, t) &= g^{\alpha\bar{\beta}}(x, t) R_{\alpha\bar{\beta}}(x, t) \cdot \frac{\det(g_{\gamma\bar{\delta}}(x, t))}{\det(g_{\gamma\bar{\delta}}(x, 0))} \\ &\leq g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(x, t) \\ &= g^{\alpha\bar{\beta}}(x, 0) (R_{\alpha\bar{\beta}}(x, t) - R_{\alpha\bar{\beta}}(x, 0)) + R(x, 0) \\ &= -\Delta_0 F(x, t) + R(x, 0) , \end{aligned} \quad (2.5)$$

where  $\Delta_0$  denotes the Laplace operator with respect to the initial metric  $g_{\alpha\bar{\beta}}(x, 0)$  and  $R(x, t)$  denotes the scalar curvature of the metric  $g_{\alpha\bar{\beta}}(x, t)$ .

Combining (2.3) and (2.5) gives

$$e^{F(x,t)} \frac{\partial F(x, t)}{\partial t} \geq \Delta_0 F(x, t) - R(x, 0) \quad \text{on} \quad M \times [0, T_{\max}) . \quad (2.6)$$

Next, we introduce a cutoff function which will be used several times in this paper. Now, as the Ricci curvature of the initial metric is positive, we know from Schoen and Yau (Theorem 1.4.2 in [28]) or Shi [32] that there exists a positive constant  $C_3$  depending only on the dimension such that for any fixed point  $x_0 \in M$  and any number  $0 < r < +\infty$ , there exists a smooth function  $\varphi(x)$  on  $M$  satisfying

$$\begin{cases} e^{-C_3\left(1+\frac{d_0(x,x_0)}{r}\right)} \leq \varphi(x) \leq e^{-\left(1+\frac{d_0(x,x_0)}{r}\right)} , \\ |\nabla\varphi|_0(x) \leq \frac{C_3}{r}\varphi(x) , \\ |\Delta_0\varphi|(x) \leq \frac{C_3}{r^2}\varphi(x) , \end{cases} \quad (2.7)$$

for all  $x \in M$ , where  $d_0(x, x_0)$  is the distance between  $x$  and  $x_0$  with respect to the initial metric  $g_{\alpha\bar{\beta}}(x, 0)$  and  $|\cdot|_0$  stands for the corresponding  $C^0$  norm of the initial metric  $g_{\alpha\bar{\beta}}(x, 0)$ .

Combining (2.6) and (2.7), we obtain

$$\begin{aligned} \frac{\partial}{\partial t} \int_M \varphi(x) e^{F(x,t)} dV_0 &\geq \int_M (\Delta_0 F(x, t) - R(x, 0)) \varphi(x) dV_0 \\ &\geq \frac{C_3}{r^2} \int_M F(x, t) \varphi(x) dV_0 - \int_M R(x, 0) \varphi(x) dV_0, \end{aligned}$$

where  $dV_0$  denotes the volume element of the initial metric  $g_{\alpha\bar{\beta}}(x, 0)$ .

Recall that  $F(\cdot, t)$  is nonincreasing in time and  $F(\cdot, 0) \equiv 0$ . We integrate the above inequality from 0 to  $t$  to get

$$\int_M \varphi(x) (1 - e^{F(x,t)}) dV_0 \leq \frac{C_3 t}{r^2} \int_M (-F(x, t)) \varphi(x) dV_0 + t \int_M R(x, 0) \varphi(x) dV_0. \quad (2.8)$$

Since the metric is shrinking under the Ricci flow, we have

$$B_t(x_0, r) \supset B_0(x_0, r) \quad \text{for } t \geq 0, \quad 0 < r < +\infty,$$

and

$$\begin{aligned} \text{Vol}_t(B_t(x_0, r)) &\geq \text{Vol}_t(B_0(x_0, r)) \\ &= \int_{B_0(x_0, r)} e^{F(x,t)} dV_0 \\ &= \text{Vol}_0(B_0(x_0, r)) + \int_{B_0(x_0, r)} (e^{F(x,t)} - 1) dV_0. \end{aligned} \quad (2.9)$$

Then by (2.7) and (2.8), the last term in (2.9) satisfies

$$\begin{aligned} \int_{B_0(x_0, r)} (e^{F(x,t)} - 1) dV_0 &\geq e^{2C_3} \int_M (e^{F(x,t)} - 1) \varphi(x) dV_0 \\ &\geq \frac{C_3 e^{2C_3} t}{r^2} \int_M F(x, t) \varphi(x) dV_0 \\ &\quad - e^{2C_3} t \int_M R(x, 0) \varphi(x) dV_0. \end{aligned} \quad (2.10)$$

To estimate the two terms of the right hand side of (2.10), we consider any fixed  $T_0 < T_{\max}$ . Since the curvature is uniformly bounded on  $M \times [0, T_0]$ ,

it is clear from the equation (2.3) that  $F(x, t)$  is also uniformly bounded on  $M \times [0, T_0]$ .

Set

$$A = \sup \{ |F(x, t)| \mid x \in M, t \in [0, T_0] \}$$

and

$$M(r) = \sup_{a \geq r} \frac{1}{\text{Vol}_0(B_0(x_0, a))} \int_{B_0(x_0, a)} R(x, 0) dV_0 .$$

Then condition (ii)" says that  $M(r) \rightarrow 0$  as  $r \rightarrow +\infty$ . By using the standard volume comparison theorem and (2.7), we have

$$\begin{aligned} \int_M R(x, 0) \varphi(x) dV_0 &\leq \int_M R(x, 0) e^{-\left(1 + \frac{d_0(x, x_0)}{r}\right)} dV_0 \\ &= \int_{B_0(x_0, r)} R(x, 0) e^{-\left(1 + \frac{d_0(x, x_0)}{r}\right)} dV_0 \\ &\quad + \sum_{k=0}^{\infty} \int_{B_0(x_0, 2^{k+1}r) \setminus B_0(x_0, 2^k r)} R(x, 0) e^{-\left(1 + \frac{d_0(x, x_0)}{r}\right)} dV_0 \\ &\leq \int_{B_0(x_0, r)} R(x, 0) dV_0 + \sum_{k=0}^{\infty} e^{-2^k} (2^{k+1})^4 . \\ &\quad \frac{\text{Vol}_0(B_0(x_0, r))}{\text{Vol}_0(B_0(x_0, 2^{k+1}r))} \int_{B_0(x_0, 2^{k+1}r)} R(x, 0) dV_0 \\ &\leq C_4 \cdot M(r) \cdot \text{Vol}_0(B_0(x_0, r)) , \end{aligned} \tag{2.11}$$

and similarly

$$\begin{aligned} \int_M \varphi(x) dV_0 &\leq \int_{B_0(x_0, r)} e^{-\left(1 + \frac{d_0(x, x_0)}{r}\right)} dV_0 \\ &\quad + \sum_{k=0}^{\infty} \int_{B_0(x_0, 2^{k+1}r) \setminus B_0(x_0, 2^k r)} e^{-\left(1 + \frac{d_0(x, x_0)}{r}\right)} dV_0 \\ &\leq \text{Vol}_0(B_0(x_0, r)) + \sum_{k=0}^{\infty} e^{-2^k} (2^{k+1})^4 \cdot \text{Vol}_0(B_0(x_0, r)) \\ &\leq C_4 \text{Vol}_0(B_0(x_0, r)) , \end{aligned} \tag{2.12}$$

where  $C_4$  is some positive constant independent of  $r$ .

Substituting (2.10), (2.11) and (2.12) into (2.9) and dividing by  $r^4$ , we obtain

$$\begin{aligned} \frac{\text{Vol}_t(B_t(x_0, r))}{r^4} &\geq \frac{\text{Vol}_0(B_0(x_0, r))}{r^4} - \frac{C_3 e^{2C_3} A T_0}{r^2} \left( C_4 \frac{\text{Vol}_0(B_0(x_0, r))}{r^4} \right) \\ &\quad - e^{2C_3} T_0 \left( C_4 M(r) \cdot \frac{\text{Vol}_0(B_0(x_0, r))}{r^4} \right) \\ &\geq C_1 - \frac{C_3 e^{2C_3} A T_0 \cdot C_4 C_1}{r^2} - e^{2C_3} T_0 C_4 C_1 \cdot M(r) \end{aligned}$$

by condition (i). Then letting  $r \rightarrow +\infty$ , we deduce that

$$\lim_{r \rightarrow +\infty} \frac{\text{Vol}_t(B_t(x_0, r))}{r^4} \geq C_1 .$$

Hence, by using the standard volume comparison theorem we have

$$\text{Vol}_t(B_t(x, r)) \geq C_1 r^4 \quad \text{for all } x \in M, 0 \leq r < +\infty, t \in [0, T_0] .$$

Finally, since  $T_0 < T_{\max}$  is arbitrary, this completes the proof of the proposition.  $\square$

### §3. Singularity Models

In this and the next section we will continue our study of the Ricci flow (2.1). We will use rescaling arguments to analyse the behavior of the solution of (2.1) near the maximal time  $T_{\max}$ .

First of all, let us recall some basic terminologies. According to Hamilton ( see for example, definition 16.3 in [16]), a solution to the Ricci flow, where either the manifold is compact or at each time  $t$  the evolving metric is complete and has bounded curvature, is called a singularity model if it is not flat and is of one of the following three types. Here, we have used  $Rm$  to denote the Riemannian curvature tensor.

**Type I:** The solution exists for  $-\infty < t < \Omega$  for some  $0 < \Omega < +\infty$  and

$$|Rm| \leq \frac{\Omega}{\Omega - t}$$

everywhere with equality somewhere at  $t = 0$ ;

**Type II:** The solution exists for  $-\infty < t < +\infty$  and

$$|Rm| \leq 1$$

everywhere with equality somewhere at  $t = 0$ ;

**Type III:** The solution exists for  $-A < t < +\infty$  for some  $0 < A < +\infty$  and

$$|Rm| \leq \frac{A}{A + t}$$

everywhere with equality somewhere at  $t = 0$ .

The singularity models of Type I and II are called ancient solutions in the sense that the existence time interval of the solution contains  $(-\infty, 0]$ .

Next, we recall the local injectivity radius estimate of Cheeger, Gromov and Taylor [5]. Let  $N$  be a complete Riemannian manifold of dimension  $m$  with  $\lambda \leq$  sectional curvature of  $N \leq \Lambda$  and let  $r$  be a positive constant satisfying  $r \leq \frac{\pi}{4\sqrt{\Lambda}}$  if  $\Lambda > 0$ , then the injectivity radius of  $N$  at a point  $x$  is bounded from below as follows,

$$\text{inj}_N(x) \geq r \cdot \frac{\text{Vol}(B(x, r))}{\text{Vol}(B(x, r)) + V_\lambda^m(2r)},$$

where  $V_\lambda^m(2r)$  denotes the volume of a ball with radius  $2r$  in the  $m$  dimensional model space  $V_\lambda^m$  with constant sectional curvature  $\lambda$ . In particular, it implies that for a complete Riemannian manifold  $N$  of dimension 4 with sectional curvature bounded between  $-1$  and  $1$ , the injectivity radius at a point  $x$  can be estimated as

$$\text{inj}_N(x) \geq \frac{1}{2} \cdot \frac{\text{Vol}(B(x, \frac{1}{2}))}{\text{Vol}(B(x, \frac{1}{2})) + V} \quad (3.1)$$

for some absolute positive constant  $V$ . Furthermore, if in addition  $N$  satisfies the maximal volume growth condition

$$\text{Vol}(B(x, r)) \geq C_1 r^4, \quad 0 \leq r < +\infty,$$

then (3.1) gives

$$\text{inj}_N(x) \geq \beta > 0 \tag{3.2}$$

for some positive constant  $\beta$  depending only on  $C_1$  and  $V$ .

Now, return to our setting. Let  $(M, g_{\alpha\bar{\beta}})$  be a complete, noncompact Kähler surface satisfying the same assumptions as in the Main Theorem and let  $g_{\alpha\bar{\beta}}(x, t)$  be the solution of the Ricci flow (2.1) on  $M \times [0, T_{\max})$ . Denote

$$R_{\max}(t) = \sup_{x \in M} R(x, t).$$

We have shown in Proposition 2.1 that the solution  $g_{\alpha\bar{\beta}}(\cdot, t)$  satisfies the same maximal volume growth condition (i) as the initial metric. Since condition (i) is invariant under rescaling of metrics, by a simple rescaling argument, we get the following injectivity radius estimate for the solution  $g_{\alpha\bar{\beta}}(\cdot, t)$ ,

$$\text{inj}(M, g_{\alpha\bar{\beta}}(\cdot, t)) \geq \frac{\beta}{\sqrt{R_{\max}(t)}} \quad \text{for } t \in [0, T_{\max}). \tag{3.3}$$

Then, by applying a result of Hamilton (see Theorems 16.4 and 16.5 in [16]), we know that there exists a sequence of dilations of the solution which converges to one of the singularity models of Type I, II or III. We will analyse this limit in the next section.

We conclude this section with the following lemma which will be very useful in our analysis of the Type I and Type II limits.

**Lemma 3.1** Suppose  $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(\cdot, t))$  is a complete ancient solution to the Ricci flow on a noncompact Kähler surface with nonnegative and bounded holomorphic bisectional curvature for all time. Then the curvature operator of the metric  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  is nonnegative definite everywhere on  $\widetilde{M} \times (-\infty, 0]$ .

**Proof** Choose a local orthonormal coframe  $\{\omega_1, \omega_2, \omega_3, \omega_4\}$  on an open set  $U \subset \widetilde{M}$  so that  $\omega_1 + \sqrt{-1}\omega_2$  and  $\omega_3 + \sqrt{-1}\omega_4$  are  $(1, 0)$  forms over  $U$ . Then the self-dual forms

$$\varphi_1 = \omega_1 \wedge \omega_2 + \omega_3 \wedge \omega_4, \quad \varphi_2 = \omega_2 \wedge \omega_3 + \omega_1 \wedge \omega_4, \quad \varphi_3 = \omega_3 \wedge \omega_1 + \omega_2 \wedge \omega_4$$

and the anti-self-dual forms

$$\psi_1 = \omega_1 \wedge \omega_2 - \omega_3 \wedge \omega_4, \quad \psi_2 = \omega_2 \wedge \omega_3 - \omega_1 \wedge \omega_4, \quad \psi_3 = \omega_3 \wedge \omega_1 - \omega_2 \wedge \omega_4$$

form a basis of the space of 2 forms over  $U$ . In particular,  $\varphi_1, \psi_1, \psi_2$  and  $\psi_3$  give a basis for the space of  $(1, 1)$  forms over  $U$ .

On a Kähler surface, it is well known that its curvature operator has image in the holonomy algebra  $u(2) (\subset so(4))$  spanned by  $(1, 1)$  forms. Thus, the curvature operator  $\mathbf{M}$  in the basis  $\{\varphi_1, \varphi_2, \varphi_3, \psi_1, \psi_2, \psi_3\}$  has the following form,

$$\mathbf{M} = \begin{pmatrix} a & 0 & 0 & b_1 & b_2 & b_3 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ b_1 & 0 & 0 & & & \\ b_2 & 0 & 0 & & A & \\ b_3 & 0 & 0 & & & \end{pmatrix}$$

where  $A$  is a  $3 \times 3$  symmetric matrix.

Let  $V$  be a real tangent vector of the Kähler surface  $\widetilde{M}$ . Denote by  $J$  the complex structure of the Kähler surface  $\widetilde{M}$ . It is clear that the complex 2-plane  $V \wedge JV$  is dual to  $(1, 1)$  form  $u\varphi_1 + v_1\psi_1 + v_2\psi_2 + v_3\psi_3$  satisfying the decomposability condition  $u^2 = v_1^2 + v_2^2 + v_3^2$ . Then after normalizing  $u$  to 1 by scaling, we see that the holomorphic bisectional curvature is nonnegative if and only if

$$a + b \cdot v + b \cdot w + {}^t v A w \geq 0, \quad (3.4)$$

for any unit vectors  $v = (v_1, v_2, v_3)$  and  $w = (w_1, w_2, w_3)$  in  $\mathbf{R}^3$ , where  $b$  is the vector  $(b_1, b_2, b_3)$  in  $\mathbf{M}$ .

Denote by  $a_1 \leq a_2 \leq a_3$  the eigenvalues of  $A$ . Recall that  $\text{tr } A = a$  by the Bianchi identity, so if we choose  $v$  to be the eigenvector of  $A$  with eigenvalue  $a_3$  and choose  $w = -v$ , (3.4) gives

$$a_1 + a_2 \geq 0 . \tag{3.5}$$

In particular, we have  $a_2 \geq 0$ .

To proceed further, we need to adapt the maximum principle for parabolic equations on compact manifold in Hamilton [14] to  $\widetilde{M}$ . Let

$$(a_i)_{\min}(t) = \inf_{x \in \widetilde{M}} a_i(x, t) , \quad i = 1, 2, 3$$

and

$$K = \sup_{(x,t) \in \widetilde{M} \times (-\infty, 0]} |Rm(x, t)| .$$

By assumption, the ancient solution  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  has bounded holomorphic bisectional curvature, hence  $K$  is finite. Thus, by the derivative estimate of Shi [29] (see also Theorem 7.1 in [16]), the higher order derivatives of the curvature are also uniformly bounded. In particular, we can use the maximum principle of Cheng–Yau (see Proposition 1.6 in [8]) and then, as observed in [16], this implies that the maximum principle of Hamilton in [14] actually works for the evolution equations of the curvature of  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  on the complete noncompact manifold  $\widetilde{M}$ . Thus, from [14], we obtain

$$\begin{aligned} \frac{d(a_1)_{\min}}{dt} &\geq ((a_1)_{\min})^2 + 2(a_2)_{\min} \cdot (a_3)_{\min} \\ &\geq 3((a_1)_{\min})^2 \end{aligned}$$

by (3.5). Then, for fixed  $t_0 \in (-\infty, 0)$  and  $t > t_0$ ,

$$\begin{aligned} (a_1)_{\min}(t) &\geq \frac{1}{(a_1)_{\min}^{-1}(t_0) - 3(t - t_0)} \\ &\geq \frac{1}{-K^{-1} - 3(t - t_0)} . \end{aligned}$$



Letting  $t_0 \rightarrow -\infty$ , we get

$$a_1 \geq 0, \quad \text{for all } (x, t) \in \widetilde{M} \times (-\infty, 0] \quad (3.6)$$

i.e.  $A \geq 0$ .

Finally, to prove the nonnegativity of the curvature operator  $\mathbf{M}$ , we recall its corresponding ODE from [14],

$$\frac{d\mathbf{M}}{dt} = \mathbf{M}^2 + \begin{pmatrix} 0 & 0 \\ 0 & A^\# \end{pmatrix},$$

where  $A^\# \geq 0$  is the adjoint matrix of  $A$ .

Let  $m_1$  be the smallest eigenvalue of the curvature operator  $\mathbf{M}$ . By using the maximum principle of Hamilton ([14] or [16]) again, we have

$$\frac{d(m_1)_{\min}}{dt} \geq (m_1)_{\min}^2$$

where  $(m_1)_{\min}(t) = \inf_{x \in \widetilde{M}} m_1(x, t)$ . Therefore, by the same reasoning in the derivation of (3.6), we have

$$m_1 \geq 0, \quad \text{for all } (x, t) \in \widetilde{M} \times (-\infty, 0]. \quad (3.7)$$

So  $\mathbf{M} \geq 0$  and the proof of the lemma is completed.  $\square$

## §4. Time decay estimate on curvature

Let  $(M, g_{\alpha\bar{\beta}}(x))$  be a complete noncompact Kähler surface satisfying all the assumptions in the Main Theorem and  $(M, g_{\alpha\bar{\beta}}(\cdot, t))$ ,  $t \in [0, T_{\max})$  be the maximal solution of the Ricci flow (2.1) with  $g_{\alpha\bar{\beta}}(\cdot)$  as the initial metric. Clearly, the maximal solution is of either one of the following types.

**Type I:**  $T_{\max} < +\infty$  and  $\sup(T_{\max} - t) R_{\max}(t) < +\infty$ ;

**Type II(a):**  $T_{\max} < +\infty$  and  $\sup(T_{\max} - t) R_{\max}(t) = +\infty$ ;

**Type II(b):**  $T_{\max} = +\infty$  and  $\sup tR_{\max}(t) = +\infty$ ;

**Type III:**  $T_{\max} = +\infty$  and  $\sup tR_{\max}(t) < +\infty$ .

In Section 3, we have proved that the maximal solution satisfies the following injectivity radius estimate

$$\text{inj}(M, g_{\alpha\bar{\beta}}(\cdot, t)) \geq \frac{\beta}{\sqrt{R_{\max}(t)}} \quad \text{on } [0, T_{\max}),$$

for some  $\beta > 0$ . By applying a result of Hamilton (Theorems 16.4 and 16.5 in [16]), we know that there exists a sequence of dilations of the solution converging to a singularity model of the corresponding type. Note that since the maximal solution is complete and noncompact, the limit must also be complete and noncompact. The following is the main result of this section which says that this limit must be of Type III or equivalently, the maximal solution must be of Type III.

**Theorem 4.1** Let  $(M, g_{\alpha\bar{\beta}}(x))$  be a complete noncompact Kähler surface as above. Then the Ricci flow (2.1) with  $g_{\alpha\bar{\beta}}(x)$  as the initial metric has a solution  $g_{\alpha\bar{\beta}}(x, t)$  for all  $t \in [0, +\infty)$  and  $x \in M$ . Moreover, the scalar curvature  $R(x, t)$  of the solution satisfies

$$0 \leq R(x, t) \leq \frac{C}{1+t} \quad \text{on } M \times [0, +\infty), \quad (4.1)$$

for some positive constant  $C$ .

**Proof.** We prove by contradiction. Thus, suppose the maximal solution is of Type I or Type II and let  $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$  be the limit of a sequence of dilations of the maximal solution which is then a singularity model of Type I or Type II respectively. After a study of its properties, we can blow down the singularity model and apply a dimension reduction argument to obtain the desired contradiction.

Now, recall that the maximal solution satisfies the maximal volume growth condition (i) by Proposition 2.1. Since condition (i) is also invariant under rescaling, we see that the singularity model  $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$  also satisfies the maximal volume growth condition, i.e.

$$\text{Vol}_t(\widetilde{B}_t(x, r)) \geq C_1 r^4 \quad \text{for all } 0 \leq r < +\infty \quad \text{and } x \in \widetilde{M}, \quad (4.2)$$

where  $\text{Vol}_t(\widetilde{B}_t(x, r))$  denotes the volume of the geodesic ball  $\widetilde{B}_t(x, r)$  of radius  $r$  with center at  $x$  with respect to the metric  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$ .

It is clear that the limit  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  has nonnegative holomorphic bisectional curvature. Thus, from Lemma 3.1, the curvature operator of the metric  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  is nonnegative definite everywhere.

Denote by  $\widetilde{R}(\cdot, t)$  the scalar curvature of  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  and  $\widetilde{d}_t(x, x_0)$  the geodesic distance between two points  $x, x_0 \in \widetilde{M}$  with respect to the metric  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$ . We claim that at time  $t = 0$ , we have

$$\limsup_{\widetilde{d}_0(x, x_0) \rightarrow +\infty} \widetilde{R}(x, 0) \widetilde{d}_0^2(x, x_0) = +\infty \quad (4.3)$$

for any fixed  $x_0 \in \widetilde{M}$ .

Suppose not, that is the curvature of the metric  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, 0)$  has quadratic decay. Now, by applying a result of Shi (see Theorem 8.2 in [32]), we know that the solution  $\widetilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  of the Ricci flow exists for all  $t \in (-\infty, +\infty)$  and satisfies

$$\lim_{t \rightarrow +\infty} \sup \{ \widetilde{R}(x, t) \mid x \in \widetilde{M} \} = 0. \quad (4.4)$$

On the other hand, by the Harnack inequality of Cao [4], we have

$$\frac{\partial \widetilde{R}}{\partial t} \geq 0 \quad \text{on } \widetilde{M} \times (-\infty, +\infty). \quad (4.5)$$

Thus, combining (4.4) and (4.5), we deduce that

$$\widetilde{R} \equiv 0, \quad \text{for all } (x, t) \in \widetilde{M} \times (-\infty, +\infty)$$

and hence  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  is flat for all  $t \in (-\infty, +\infty)$ . But, by definition, a singularity model cannot be flat. This proves our claim (4.3).

With the estimate (4.3), we can then apply a lemma of Hamilton (Lemma 22.2 in [16]) to find a sequence of positive numbers  $\delta_j, j = 1, 2, \dots$ , with  $\delta_j \rightarrow 0$  such that

- (a)  $\tilde{R}(x, 0) \leq (1 + \delta_j)\tilde{R}(x_j, 0)$  for all  $x$  in the ball  $\tilde{B}_0(x_j, r_j)$  of radius  $r_j$  centered at  $x_j$  with respect to the metric  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, 0)$ ;
- (b)  $r_j^2\tilde{R}(x_j, 0) \rightarrow +\infty$ ;
- (c) if  $s_j = \tilde{d}_0(x_j, x_0)$ , then  $\lambda_j = s_j/r_j \rightarrow +\infty$ ;
- (d) the balls  $\tilde{B}_0(x_j, r_j)$  are disjoint.

Denote the minimum of the holomorphic sectional curvature of the metric  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, 0)$  at  $x_j$  by  $h_j$ . We claim that the following holds

$$\varepsilon_j = \frac{h_j}{\tilde{R}(x_j, 0)} \rightarrow 0 \quad \text{as } j \rightarrow +\infty. \quad (4.6)$$

Suppose not, there exists a subsequence  $j_k \rightarrow +\infty$  and some positive number  $\varepsilon > 0$  such that

$$\varepsilon_{j_k} = \frac{h_{j_k}}{\tilde{R}(x_{j_k}, 0)} \geq \varepsilon \quad \text{for all } k = 1, 2, \dots. \quad (4.7)$$

Since the solution  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  is ancient, it follows from the Harnack inequality of Cao [4] that the scalar curvature  $\tilde{R}(x, t)$  is pointwisely nondecreasing in time. Then, by using the local derivative estimate of Shi [29] (or see Theorem 13.1 in [16]) and (a), (b), we have

$$\begin{aligned} \sup_{x \in \tilde{B}_0(x_{j_k}, r_{j_k})} \left| \nabla \tilde{R} m(x, 0) \right|^2 &\leq C_5 \tilde{R}^2(x_j, 0) \left( \frac{1}{r_{j_k}^2} + \tilde{R}(x_j, 0) \right) \\ &\leq 2C_5 \tilde{R}^3(x_j, 0), \end{aligned} \quad (4.8)$$

where  $\tilde{R}m$  is the curvature tensor of  $\tilde{g}_{\alpha\bar{\beta}}$  and  $C_5$  is a positive constant depending only on the dimension.

For any  $x \in \tilde{B}_0(x_{j_k}, r_{j_k})$ , we obtain from (4.7) and (4.8) that the minimum of the holomorphic sectional curvature  $h_{\min}(x)$  at  $x$ , satisfies

$$\begin{aligned} h_{\min}(x) &\geq h_{j_k} - \sqrt{2C_5} \tilde{R}^{3/2}(x_{j_k}, 0) \tilde{d}_0(x, x_{j_k}) \\ &\geq \tilde{R}(x_{j_k}, 0) \left( \varepsilon - \sqrt{2C_5} \cdot \sqrt{\tilde{R}(x_{j_k}, 0)} \cdot \tilde{d}_0(x, x_{j_k}) \right) \\ &\geq \frac{\varepsilon}{2} \tilde{R}(x_{j_k}, 0) \end{aligned} \quad (4.9)$$

if

$$\tilde{d}_0(x, x_{j_k}) \leq \frac{\varepsilon}{2\sqrt{2C_5} \cdot \sqrt{\tilde{R}(x_{j_k}, 0)}}.$$

Thus, from (a) and (4.9), there exists  $k_0 > 0$  such that for any  $k \geq k_0$  and

$$x \in \tilde{B}_0\left(x_{j_k}, \frac{\varepsilon}{2\sqrt{2C_5} \cdot \sqrt{\tilde{R}(x_{j_k}, 0)}}\right),$$

we have

$$\frac{\varepsilon}{2} \tilde{R}(x_{j_k}, 0) \leq \text{holomorphic sectional curvature at } x \leq 2\tilde{R}(x_{j_k}, 0). \quad (4.10)$$

We have proved that the metric  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, 0)$  has nonnegative definite curvature operator. In particular, the sectional curvature is nonnegative. Then, by the generalized Cohn–Vossen inequality in real dimension 4 [12], we have

$$\int_{\tilde{M}} \Theta \leq \chi(\tilde{M}) < +\infty \quad (4.11)$$

where  $\Theta$  is the Gauss–Bonnet–Chern integrand for the metric  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, 0)$  and  $\chi(\tilde{M})$  is the Euler number of the manifold  $\tilde{M}$  which has finite topology type by the soul theorem of Cheeger–Gromoll.

On the other hand, from the proof of Theorem 1.3 of Bishop–Goldberg [2] (see Page 523 of [2]), the inequality (4.10) implies that

$$\Theta(x) \geq C(\varepsilon) \tilde{R}^2(x_{j_k}, 0) \quad \text{for all } x \in \tilde{B}_0\left(x_{j_k}, \frac{\varepsilon}{2\sqrt{2C_5} \cdot \sqrt{\tilde{R}(x_{j_k}, 0)}}\right), \quad (4.12)$$

where  $C(\varepsilon)$  is some positive constant depending only on  $\varepsilon$ . Now, by combining (4.2), (b), (d), (4.11) and (4.12), we get

$$\begin{aligned}
+\infty > \chi(\widetilde{M}) &\geq \sum_{k=k_0}^{\infty} \int_{\widetilde{B}_0(x_{j_k}, \frac{\varepsilon}{2\sqrt{2}C_5 \cdot \sqrt{\widetilde{R}(x_{j_k}, 0)}})} \Theta \\
&\geq C(\varepsilon) \sum_{k=k_0}^{\infty} \widetilde{R}^2(x_{j_k}, 0) \cdot C_1 \left( \frac{\varepsilon}{2\sqrt{2}C_5 \cdot \sqrt{\widetilde{R}(x_{j_k}, 0)}} \right)^4 \\
&= C(\varepsilon) \sum_{k=k_0}^{\infty} \frac{C_1 \varepsilon^4}{64C_5^2} \\
&= +\infty
\end{aligned}$$

which is a contradiction. Hence our claim (4.6) is proved.

Now, we are going to blow down the singularity model  $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$ . For the above chosen  $x_j, r_j$  and  $\delta_j$ , let  $x_j$  be the new origin  $O$ , dilate the space by a factor  $\lambda_j$  so that  $\widetilde{R}(x_j, 0)$  become 1 at the origin at  $t = 0$ , and dilate in time by  $\lambda_j^2$  so that it is still a solution to the Ricci flow. The balls  $\widetilde{B}_0(x_j, r_j)$  are dilated to the balls centered at the origin of radii  $\widetilde{r}_j = r_j^2 \widetilde{R}(x_j, 0) \rightarrow +\infty$  ( by (b) ). Since the scalar curvature of  $\widetilde{g}_{\alpha\bar{\beta}}(x, t)$  is pointwise nondecreasing in time by the Harnack inequality, the curvature bounds on  $\widetilde{B}_0(x_j, r_j)$  also give bounds for previous times in these balls. And the maximal volume growth estimate (4.2) and the local injectivity radius estimate of Cheeger, Gromov and Taylor [5] imply that

$$\text{inj}_{\widetilde{M}}(x_j, \widetilde{g}_{\alpha\bar{\beta}}(\cdot, 0)) \geq \frac{\beta}{\sqrt{\widetilde{R}(x_j, 0)}},$$

for some positive constant  $\beta$  independent of  $j$ .

So we have everything to take a limit for the dilated solutions. By applying the compactness theorem in [15] and combining (4.2), (4.6), (a) and (b), we obtain a complete noncompact solution, still denoted by  $(\widetilde{M}, \widetilde{g}_{\alpha\bar{\beta}}(x, t))$ , for  $t \in (-\infty, 0]$  such that

- (e) the curvature operator is still nonnegative;

- (f)  $\tilde{R}(x, t) \leq 1$ , for all  $x \in \tilde{M}$ ,  $t \in (-\infty, 0]$ , and  $\tilde{R}(0, 0) = 1$ ;
- (g)  $\text{Vol}_t(\tilde{B}_t(x, r)) \geq C_1 r^4$  for all  $x \in \tilde{M}$ ,  $0 \leq r \leq +\infty$ ;
- (h) there exists a complex 2-plane  $V \wedge JV$  at the origin  $O$  so that at  $t = 0$ , the corresponding holomorphic sectional curvature vanishes.

If we consider the universal covering of  $\tilde{M}$ , the induced metric of  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  on the universal covering is clearly still a solution to the Ricci flow and satisfies all of above (e), (f), (g), (h). Thus, without loss of generality, we may assume that  $\tilde{M}$  is simply connected.

Next, by using the strong maximum principle on the evolution equation of the curvature operator of  $\tilde{g}_{\alpha\bar{\beta}}(\cdot, t)$  as in [14] (see Theorem 8.3 of [14]), we know that there exists a constant  $K > 0$  such that on the time interval  $-\infty < t < -K$ , the image of the curvature operator of  $(\tilde{M}, \tilde{g}_{\alpha\bar{\beta}}(\cdot, t))$  is a fixed Lie subalgebra of  $so(4)$  of constant rank on  $\tilde{M}$ . Because  $\tilde{M}$  is Kähler, the possibilities are limited to  $u(2)$ ,  $so(2) \times so(2)$  or  $so(2)$ .

In the case  $u(2)$ , the sectional curvature is strictly positive. Thus, this case is ruled out by (h). In the cases  $so(2) \times so(2)$  or  $so(2)$ , according to [14], the simply connected manifold  $\tilde{M}$  splits as a product  $\tilde{M} = \Sigma_1 \times \Sigma_2$ , where  $\Sigma_1$  and  $\Sigma_2$  are two Riemann surfaces with nonnegative curvature (by (e)), and at least one of them, say  $\Sigma_1$ , has positive curvature (by (f)).

Denote by  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(\cdot, t)$  the corresponding metric on  $\Sigma_1$ . Clearly, it follows from (g) and standard volume comparison that for any  $x \in \Sigma_1$ ,  $t \in (-\infty, -K)$ , we have

$$\text{Vol}B_{\Sigma_1}(x, r) \geq C_6 r^2 \quad \text{for } 0 \leq r < +\infty \quad (4.13)$$

where both the geodesic ball  $B_{\Sigma_1}(x, r)$  and the volume are taken with respect to the metric  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(\cdot, t)$  on  $\Sigma_1$ ,  $C_6$  is a positive constant depending only on  $C_1$ . Also as the curvature of  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$  is positive, it follows from Cohn–Vossen

inequality that

$$\int_{\Sigma_1} \tilde{R}^{(1)}(x, t) d\sigma_t \leq 8\pi, \quad (4.14)$$

where  $\tilde{R}^{(1)}(x, t)$  is the scalar curvature of  $(\Sigma_1, \tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t))$  and  $d\sigma_t$  is the volume element of the metric  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$ .

Now, the metric  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$  is a solution to the Ricci flow on the Riemann surface  $\Sigma_1$  over the ancient time interval  $(-\infty, -K)$ . Thus, (4.13) and (4.14) imply that for each  $t \in (-\infty, -K)$ , the curvature of  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$  has quadratic decay in the average sense of Shi [32] and then the apriori estimate of Shi (see Theorem 8.2 in [32]) implies that the solution  $\tilde{g}_{\alpha\bar{\beta}}^{(1)}(x, t)$  exists for all  $t \in (-\infty, +\infty)$  and satisfies

$$\lim_{t \rightarrow +\infty} \sup \left\{ \tilde{R}^{(1)}(x, t) \mid x \in \Sigma_1 \right\} = 0. \quad (4.15)$$

Again, by the Harnack inequality of Cao [4], we know that  $\tilde{R}^{(1)}(x, t)$  is pointwisely nondecreasing in time. Therefore, we conclude that

$$\tilde{R}^{(1)}(x, t) \equiv 0 \quad \text{on} \quad \Sigma_1 \times (-\infty, +\infty).$$

This contradicts with the fact that  $(\Sigma_1, \tilde{g}_{\alpha\bar{\beta}}(\cdot, t))$  has positive curvature for  $t < -K$ . Hence we have seeked the desired contradiction and have completed the proof of Theorem 4.1.  $\square$

## §5. Topology and Steinness

In this section, we use the estimates obtained in the previous sections to study the topology and the complex structure of the Kähler surface in our Main Theorem. Our result is

**Theorem 5.1** Suppose  $(M, g_{\alpha\bar{\beta}})$  is a complete noncompact Kähler surface satisfying the assumptions in the Main Theorem. Then  $M$  is homeomorphic to  $\mathbf{R}^4$  and is a Stein manifold.



The proof of this theorem is exactly the same as in [7]. For the convenience of the readers, we give a sketch of the arguments and refer to the cited reference for details.

**Sketch of proof.** We evolve the metric  $g_{\alpha\bar{\beta}}(x)$  by the Ricci flow (2.1). From Theorem 4.1, the solution  $g_{\alpha\bar{\beta}}(x, t)$  exists for all  $t \in [0, +\infty)$  and satisfies

$$R(x, t) \leq \frac{C}{1+t} \quad \text{on } M \times [0, +\infty) \quad (5.1)$$

for some positive constant  $C$ . Also, Proposition 2.1 tells us that the volume growth condition (i) is preserved under the Ricci flow. By using the local injectivity radius estimate of Cheeger–Gromov–Taylor, this implies that

$$\text{inj} \left( M, g_{\alpha\bar{\beta}}(\cdot, t) \right) \geq C_7(1+t)^{\frac{1}{2}} \quad \text{for } t \in [0, +\infty) \quad (5.2)$$

with some positive constant  $C_7$ .

Since the Ricci curvature of  $g_{\alpha\bar{\beta}}(x, t)$  is positive for all  $x \in M$  and  $t \geq 0$ , the Ricci flow equation (2.1) implies that the ball  $B_t(x_0, \frac{C_7}{2}(1+t)^{\frac{1}{2}})$  of radius  $\frac{C_7}{2}(1+t)^{\frac{1}{2}}$  with respect to the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$  contains the ball  $B_0(x_0, \frac{C_7}{2}(1+t)^{\frac{1}{2}})$  of the same radius with respect to the initial metric  $g_{\alpha\bar{\beta}}(\cdot, 0)$ . Combining this with (5.2), we deduce that

$$\pi_p(M, x_0) = 0 \quad \text{for any } p \geq 1$$

and

$$\pi_q(M, \infty) = 0 \quad \text{for } 1 \leq q \leq 2,$$

where  $\pi_q(M, \infty)$  is the  $q$ th homotopy group of  $M$  at infinity.

Thus, by the resolution of the generalized Poincaré conjecture on four manifolds by Freedman [11], we know that  $M$  is homeomorphic to  $\mathbf{R}^4$ .

Next, the injectivity radius estimate (5.2) also tells us that, for  $t$  large enough, the exponential maps provide diffeomorphisms between big geodesic balls  $B_t(x_0, \frac{C_7}{2}(1+t)^{\frac{1}{2}})$  of  $M$  with big Euclidean balls on  $\mathbf{C}^2$ . The curvature

estimate (5.1) together with its derivative estimates imply that the difference of the complex structure of  $M$  and the standard complex structure of  $\mathbf{C}^2$  in those geodesic balls can be made arbitrarily small by taking  $t$  large enough. Then, we can use the  $L^2$  estimates of Hörmander [17] to modify the exponential map to a biholomorphism between a domain  $\Omega(t)$  containing  $B_t(x_0, \frac{C_7}{4}(1+t)^{\frac{1}{2}})$  and a Euclidean ball of radius  $\frac{C_7}{3}(1+t)^{\frac{1}{2}}$  in  $\mathbf{C}^2$ . Since the solution  $g_{\alpha\bar{\beta}}(\cdot, t)$  of the Ricci flow is shrinking, the domain  $\Omega(t)$  contains the geodesic ball  $B_0(x_0, \frac{C_7}{4}(1+t)^{\frac{1}{2}})$ . Thus, we can choose a sequence of  $t_k \rightarrow +\infty$ , such that

$$M = \cup_{k=0}^{+\infty} \Omega(t_k) , \quad \Omega(t_1) \subset \Omega(t_2) \subset \cdots \subset \Omega(t_k) \subset \cdots .$$

Since for each  $k$ ,  $\Omega(t_k)$  is biholomorphic to the unit ball of  $\mathbf{C}^2$ ,  $(\Omega(t_k), \Omega(t_l))$  forms a Runge pair for any  $k, l$ . Finally, we can appeal to a theorem of Markeo [19] (see also Siu [33]) to conclude that  $M$  is a Stein manifold.  $\square$

## §6. Space decay estimate on curvature and the Poincaré–Lelong equation

Let  $(M, g_{\alpha\bar{\beta}})$  be a complete noncompact Kähler surface satisfying all the assumptions in the Main Theorem. The main purpose of this section is to establish the existence of a strictly plurisubharmonic function of logarithmic growth on  $M$ . To this end, we first prove a curvature decay estimate at infinity of the metric  $g_{\alpha\bar{\beta}}$ .

**Theorem 6.1** Let  $(M, g_{\alpha\bar{\beta}})$  be a complete noncompact Kähler surface as above. Then there exists a constant  $C > 0$  such that for all  $x \in M$ ,  $r > 0$ , we have

$$\int_{B(x,r)} R(y) \frac{1}{d^2(x,y)} dy \leq C \log(2+r) . \quad (6.1)$$

**Proof.** Let  $g_{\alpha\bar{\beta}}(x, t)$  be the solution of the Ricci flow (2.1) with  $g_{\alpha\bar{\beta}}(x)$  as the initial metric. From Theorem 4.1, we know that the solution exists for

all times and satisfies

$$R(x, t) \leq \frac{C_8}{1+t} \quad \text{on } M \times [0, +\infty) \quad (6.2)$$

for some positive constant  $C_8$ .

Let

$$F(x, t) = \log \frac{\det(g_{\alpha\bar{\beta}}(x, t))}{\det(g_{\alpha\bar{\beta}}(x, 0))}$$

be the function introduced in the proof of Proposition 2.1. Since

$$-\partial_\alpha \bar{\partial}_\beta \log \frac{\det(g_{\gamma\bar{\delta}}(\cdot, t))}{\det(g_{\gamma\bar{\delta}}(\cdot, 0))} = R_{\alpha\bar{\beta}}(\cdot, t) - R_{\alpha\bar{\beta}}(\cdot, 0) ,$$

after taking trace with the initial metric  $g_{\alpha\bar{\beta}}(\cdot, 0)$ , we get

$$R(\cdot, 0) = \Delta_0 F(\cdot, t) + g^{\alpha\bar{\beta}}(\cdot, 0) R_{\alpha\bar{\beta}}(\cdot, t) \quad (6.3)$$

where  $\Delta_0$  is the Laplace operator of the metric  $g_{\alpha\bar{\beta}}(\cdot, 0)$ .

Since  $(M, g_{\alpha\bar{\beta}}(\cdot, 0))$  has positive Ricci curvature and maximal volume growth, it is well known (see [28]) that the Green function  $G_0(x, y)$  of the initial metric  $g_{\alpha\bar{\beta}}(\cdot, 0)$  exists on  $M$  and satisfies the estimates

$$\frac{C_9^{-1}}{d_0^2(x, y)} \leq G_0(x, y) \leq \frac{C_9}{d_0^2(x, y)} \quad (6.4)$$

and

$$|\nabla_y G_0(x, y)|_0 \leq \frac{C_9}{d_0^3(x, y)} \quad (6.5)$$

for some positive constant  $C_9$  depending only on  $C_1$ .

For any fixed  $\bar{x}_0 \in M$  and any  $\alpha > 0$ , we denote

$$\Omega_\alpha = \{ x \in M \mid G_0(\bar{x}_0, x) \geq \alpha \} .$$

By (6.4), it is not hard to see

$$B_0 \left( \bar{x}_0, \left( \frac{C_9^{-1}}{\alpha} \right)^{\frac{1}{2}} \right) \subset \Omega_\alpha \subset B_0 \left( \bar{x}_0, \left( \frac{C_9}{\alpha} \right)^{\frac{1}{2}} \right) . \quad (6.6)$$

Recall that  $F$  evolves by

$$\frac{\partial F(x, t)}{\partial t} = -R(x, t) \quad \text{on } M \times [0, +\infty) .$$

Combining with (6.2), we obtain

$$0 \geq F(x, t) \geq -C_{10} \log(1 + t) \quad \text{on } M \times [0, +\infty) . \quad (6.7)$$

Multiplying (6.3) by  $G_0(\bar{x}_0, x) - \alpha$  and integrating over  $\Omega_\alpha$ , we have

$$\begin{aligned} \int_{\Omega_\alpha} R(x, 0) (G_0(\bar{x}_0, x) - \alpha) dx &= \int_{\Omega_\alpha} (\Delta_0 F(x, t)) (G_0(\bar{x}_0, x) - \alpha) dx \\ &\quad + \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(\cdot, t) (G_0(\bar{x}_0, x) - \alpha) dx \\ &= - \int_{\partial\Omega_\alpha} F(x, t) \frac{\partial G_0(\bar{x}_0, x)}{\partial \nu} d\sigma - F(\bar{x}_0, t) \\ &\quad + \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(\cdot, t) (G_0(\bar{x}_0, x) - \alpha) dx \\ &\leq C_{10} \left( 1 + C_9^{\frac{5}{2}} \alpha^{\frac{3}{2}} \text{Vol}_0(\partial\Omega_\alpha) \right) \log(1 + t) \\ &\quad + \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(\cdot, t) G_0(\bar{x}_0, x) dx , \end{aligned} \quad (6.8)$$

by (6.4) and (6.7). Here, we have used  $\nu$  to denote the outer unit normal of  $\partial\Omega_\alpha$ .

From the coarea formula, we have

$$\begin{aligned} \frac{1}{\alpha} \int_\alpha^{2\alpha} r^{\frac{3}{2}} \text{Vol}_0(\partial\Omega_r) dr &\leq 2^{\frac{3}{2}} \alpha^{\frac{1}{2}} \int_\alpha^{2\alpha} \int_{\partial\Omega_r} |\nabla G_0(\bar{x}_0, x)|_0 d\sigma |d\nu| \\ &\leq 2^{\frac{3}{2}} C_9^{\frac{5}{2}} \alpha^2 \text{Vol}_0(\Omega_\alpha) \\ &\leq 2^{\frac{3}{2}} C_9^{\frac{5}{2}} \alpha^2 \text{Vol}_0 \left( B_0 \left( \bar{x}_0, \left( \frac{C_9}{\alpha} \right)^{\frac{1}{2}} \right) \right) \\ &\leq C_{11} \end{aligned}$$

for some positive constant  $C_{11}$  by the standard volume comparison. Substitute this into (6.8) and integrate (6.8) from  $\alpha$  to  $2\alpha$ , we get

$$\int_{\Omega_{2\alpha}} R(x, 0) (G_0(\bar{x}_0, x) - 2\alpha) dx \leq C_{10} \left( 1 + C_9^{\frac{5}{2}} C_{11} \right) \log(1 + t)$$

$$+ \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(x, t) G_0(\bar{x}_0, x) dx . \quad (6.9)$$

It is easy to see that

$$\int_{\Omega_{4\alpha}} R(x, 0) G_0(\bar{x}_0, x) dx \leq 2 \int_{\Omega_{2\alpha}} R(x, 0) (G_0(\bar{x}_0, x) - 2\alpha) dx$$

and by the equation Ricci flow (2.1), we also have

$$\begin{aligned} & \int_0^t \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) R_{\alpha\bar{\beta}}(x, t) G_0(\bar{x}_0, x) dx dt \\ &= \int_{\Omega_\alpha} g^{\alpha\bar{\beta}}(x, 0) (g_{\alpha\bar{\beta}}(x, 0) - g_{\alpha\bar{\beta}}(x, t)) G_0(\bar{x}_0, x) dx \\ &\leq 2 \int_{\Omega_\alpha} G_0(\bar{x}_0, x) dx . \end{aligned}$$

Thus by integrating (6.9) in time from 0 to  $t$  and combining the above two inequalities, we get for any  $t > 0$ ,

$$\int_{\Omega_{4\alpha}} R(x, 0) G_0(\bar{x}_0, x) dx \leq 2C_{10} \left(1 + C_9^{\frac{5}{2}} C_{11}\right) \log(1+t) + \frac{4}{t} \int_{\Omega_\alpha} G_0(\bar{x}_0, x) dx .$$

Finally, substituting (6.4) and (6.6) into the above inequality, we see that there exists some positive constant  $C_{12}$  such that for any  $\bar{x}_0 \in M$ ,  $t > 0$  and  $r > 0$ ,

$$\int_{B_0(\bar{x}_0, r)} R(x, 0) \frac{1}{d^2(\bar{x}_0, x)} dx \leq C_{12} \left( \log(1+t) + \frac{r^2}{t} \right) . \quad (6.10)$$

Choose  $t = r^2$ , we get the desired estimate.  $\square$

Now we can use the estimate (6.1) to solve the following Poincaré–Lelong equation on  $M$

$$\sqrt{-1}\partial\bar{\partial}u = \text{Ric} \quad (6.11)$$

to get the strictly plurisubharmonic function mentioned at the beginning of this section.

As in [20] or [25], we first study the corresponding Poisson equation on  $M$

$$\Delta u = R. \quad (6.12)$$

After we solve the Poisson equation (6.12) with a solution of logarithmic growth, we will see that it is indeed a solution of the Poincaré–Lelong equation with logarithmic growth.

To solve (6.12), we first construct a family of approximate solutions  $u_r$  as follows.

For a fixed  $x_0 \in M$  and  $r > 0$ , define  $u_r(x)$  on  $B(x_0, r)$  by

$$u_r(x) = \int_{B(x_0, r)} (G(x_0, y) - G(x, y)) R(y) dy$$

where  $G(x, y)$  is the Green function of the metric  $g_{\alpha\bar{\beta}}$  on  $M$ . It is clear that

$$u_r(x_0) = 0 \quad \text{and} \quad \Delta u_r(x) = R(x) \quad \text{on} \quad B(x_0, r).$$

For  $x \in B(x_0, \frac{r}{2})$ , we write

$$\begin{aligned} u_r(x) &= \left( \int_{B(x_0, r) \setminus B(x_0, 2d(x, x_0))} + \int_{B(x_0, 2d(x, x_0))} \right) (G(x_0, y) - G(x, y)) R(y) dy \\ &:= I_1 + I_2 . \end{aligned}$$

From (6.1), we see that

$$|I_2| \leq C_{13} \log(2 + d(x, x_0)) \quad \text{on} \quad B(x_0, \frac{r}{2}) \quad (6.13)$$

for some positive constant  $C_{13}$  independent of  $x_0$ ,  $x$  and  $r$ .

To estimate  $I_1$ , we get from (6.5) that for  $y \in B(x_0, r) \setminus B(x_0, 2d(x, x_0))$ ,

$$\begin{aligned} |G(x_0, y) - G(x, y)| &\leq d(x, x_0) \cdot \sup_{z \in B(x_0, d(x, x_0))} |\nabla_z G(z, y)| \\ &\leq C_9 d(x, x_0) \cdot \sup_{z \in B(x_0, d(x, x_0))} \frac{1}{d^3(z, y)} \\ &\leq 8C_9 \frac{d(x, x_0)}{d^3(y, x_0)} . \end{aligned}$$

Thus by (6.1), we have

$$|I_1| \leq 8C_9 d(x, x_0) \int_{B(x_0, r) \setminus B(x_0, 2d(x, x_0))} \frac{R(y)}{d^3(y, x_0)} dy$$

$$\begin{aligned}
&\leq 8C_9 d(x, x_0) \sum_{k=1}^{\infty} \frac{1}{2^k d(x, x_0)} \cdot \int_{B(x_0, 2^{k+1}d(x, x_0)) \setminus B(x_0, 2^k d(x, x_0))} \frac{R(y)}{d^2(y, x_0)} dy \\
&\leq 8C_9 C \sum_{k=1}^{\infty} \frac{1}{2^k} \log(2 + 2^{k+1}d(x, x_0)) \\
&\leq C_{14} \log(2 + d(x, x_0))
\end{aligned} \tag{6.14}$$

for some positive constant  $C_{14}$ .

Hence, by combining (6.13) and (6.14), we deduce

$$|u_r(x)| \leq (C_{13} + C_{14}) \log(2 + d(x, x_0)) \tag{6.15}$$

for any  $r \geq 2d(x, x_0)$ .

On the other hand, by taking the derivative of  $u_r(x)$ , we get

$$\begin{aligned}
|\nabla u_r(x)| &\leq C_9 \int_M \frac{R(y)}{d^3(x, y)} dy \\
&\leq C_9 \left( \int_{B(x, 1)} \frac{R(y)}{d^3(x, y)} dy + \sum_{k=1}^{\infty} \frac{1}{2^{k-1}} \int_{B(x, 2^k) \setminus B(x, 2^{k-1})} \frac{R(y)}{d^2(x, y)} dy \right) \\
&\leq C_9 \left( C_{15} + \sum_{k=1}^{\infty} \frac{1}{2^{k-1}} C \log(2 + 2^k) \right) \\
&= C_{16} .
\end{aligned} \tag{6.16}$$

Here, we have used (6.1) and (6.5),  $C_{15}$  and  $C_{16}$  are positive constants independent of  $r$ . Therefore, it follows from the Schauder theory of elliptic equations that there exists a sequence of  $r_j \rightarrow +\infty$  such that  $u_{r_j}(x)$  converges uniformly on compact subset of  $M$  to a smooth function  $u$  satisfying

$$\begin{cases} u(x_0) = 0 & \text{and} & \Delta u = R & \text{on } M , \\ |u(x)| \leq (C_{13} + C_{14}) \log(2 + d(x, x_0)) & \text{for } x \in M , \\ |\nabla u(x)| \leq C_{16} & \text{for } x \in M . \end{cases} \tag{6.17}$$

Thus, we have obtained a solution  $u$  of logarithmic growth to the Poisson equation (6.12) on  $M$ . In the following we prove that  $u$  is actually a solution of the Poincaré–Lelong equation (6.11).

Recall the Bochner identity, with  $\Delta u = R$

$$\begin{aligned} \frac{1}{2} \Delta |\nabla u|^2 &= |\nabla^2 u|^2 + \langle \nabla u, \nabla R \rangle + Ric(\nabla u, \nabla u) \\ &\geq |\nabla^2 u|^2 + \langle \nabla u, \nabla R \rangle . \end{aligned} \quad (6.18)$$

For any  $r > 0$  and any  $\bar{x}_0 \in M$ , by multiplying (6.18) by the cutoff function in (2.7) and integrating by parts, we get

$$\begin{aligned} \int_M |\nabla^2 u|^2 \varphi dx &\leq \frac{1}{2} \int_M |\nabla u|^2 \cdot |\Delta \varphi| dx + \int_M |\nabla^2 u| \cdot R \varphi dx \\ &\quad + \int_M |\nabla u| \cdot R \cdot |\nabla \varphi| dx \\ &\leq \frac{C_{16}^2}{2} \cdot \frac{C_3}{r^2} \int_M \varphi dx + \frac{1}{2} \int_M |\nabla^2 u|^2 \varphi dx \\ &\quad + \frac{1}{2} \int_M R^2 \varphi dx + C_{16} \cdot \left( \sup_M R \right) \cdot \frac{C_3}{r} \int_M \varphi dx . \end{aligned}$$

Thus,

$$\int_M |\nabla^2 u|^2 \varphi dx \leq \left( C_3 C_{16}^2 \cdot \frac{1}{r^2} + 2C_{16} C_3 \left( \sup_M R \right) \frac{1}{r} \right) \int_M \varphi dx + \int_M R^2 \varphi dx . \quad (6.19)$$

By (6.1), (2.7) and the standard volume comparison, we have

$$\begin{aligned} \int_M R^2 \varphi dx &\leq \left( \sup_M R \right) \int_M R(x) e^{-\left(1 + \frac{d(x, \bar{x}_0)}{r}\right)} dx \\ &\leq \left( \sup_M R \right) \cdot \\ &\quad \left( \int_{B(\bar{x}_0, r)} R(x) dx + \sum_{k=0}^{\infty} e^{-2^{k-1}} \int_{B(\bar{x}_0, 2^{k+1}r) \setminus B(\bar{x}_0, 2^k r)} R(x) dx \right) \\ &\leq C_{17} r^2 \log(2+r) \end{aligned} \quad (6.20)$$

and

$$\begin{aligned} \int_M \varphi dx &\leq \int_M e^{-\left(1 + \frac{d(x, \bar{x}_0)}{r}\right)} dx \\ &\leq \int_{B(\bar{x}_0, r)} dx + \sum_{k=0}^{\infty} e^{-2^{k-1}} \int_{B(\bar{x}_0, 2^{k+1}r) \setminus B(\bar{x}_0, 2^k r)} dx \\ &\leq C_{17} r^4 \end{aligned} \quad (6.21)$$



for some positive constant  $C_{17}$  independent of  $r$  and  $\bar{x}_0$ .

Substituting these two inequalities into (6.19) we have

$$\frac{1}{r^4} \int_{B(\bar{x}_0, r)} |\nabla^2 u|^2 dx \leq C_{18} \left( \frac{1}{r^2} + \frac{1}{r} + \frac{\log(2+r)}{r^2} \right) \quad (6.22)$$

for some positive constant  $C_{18}$  independent of  $r$  and  $\bar{x}_0$ .

Since the holomorphic bisectional curvature of  $g_{\alpha\bar{\beta}}$  is positive, it was shown in [20] that the function  $|\sqrt{-1}\partial\bar{\partial}u - Ric|^2$  is subharmonic on  $M$ . Then by the mean value inequality and (6.20), (6.22), we have

$$\begin{aligned} |\sqrt{-1}\partial\bar{\partial}u - Ric|^2(\bar{x}_0) &\leq \frac{C_{19}}{r^4} \int_{B(\bar{x}_0, r)} |\sqrt{-1}\partial\bar{\partial}u - Ric|^2(x) dx \\ &\leq \frac{2C_{19}}{r^4} \int_{B(\bar{x}_0, r)} (|\nabla^2 u|^2 + R^2) dx \\ &\leq C_{20} \left( \frac{1}{r^2} + \frac{1}{r} + \frac{\log(2+r)}{r^2} \right) \end{aligned}$$

for some positive constants  $C_{19}$ ,  $C_{20}$  independent of  $r$  and  $\bar{x}_0$ . Since  $\bar{x}_0 \in M$  and  $r > 0$  are arbitrary, by letting  $r \rightarrow +\infty$  we know that

$$\sqrt{-1}\partial\bar{\partial}u = Ric \quad \text{on } M.$$

In summary, we have proved the following result.

**Proposition 6.2** Suppose  $(M, g_{\alpha\bar{\beta}})$  is a complete noncompact Kähler surface satisfying all the assumptions in the Main Theorem. Then there exists a strictly plurisubharmonic function  $u(x)$  on  $M$  satisfying the Poincaré–Lelong equation (6.11) with the estimate

$$|u(x)| \leq C \log(2 + d(x, x_0)) \quad \text{for all } x \in M$$

for some positive constant  $C$ .

## §7. Uniform estimates on multiplicity and the number of components of an “algebraic” divisor

Let  $(M, g_{\alpha\bar{\beta}})$  be a complete noncompact Kähler surface satisfying all the assumptions in the Main Theorem. In this section, we will consider the algebra  $P(M)$  of holomorphic functions of polynomial growth on  $M$ . We first construct  $f_1, f_2$  in  $P(M)$  which are algebraically independent over  $\mathbf{C}$ .

In the previous section, by solving the Poincaré–Lelong equation, we have obtained a strictly plurisubharmonic function  $u$  on  $M$  of logarithmic growth. As shown in [21], the existence of nontrivial functions in the algebra  $P(M)$  then follows readily from the  $L^2$  estimates of the  $\bar{\partial}$  operator on complete Kähler manifold of Andreotti–Vesentini [1] and Hörmander [17]. For completeness, we give the proof as follows.

Let  $x \in M$  and  $\{(z_1, z_2) \mid |z_1|^2 + |z_2|^2 < 1\}$  be local holomorphic coordinates at  $x$  with  $z_1(x) = z_2(x) = 0$ . Let  $\eta$  be a smooth cutoff function on  $\mathbf{C}^2$  with  $\text{Supp } \eta \subset\subset \{|z_1|^2 + |z_2|^2 < 1\}$  and  $\eta \equiv 1$  on  $\{|z_1|^2 + |z_2|^2 < \frac{1}{4}\}$ . Then the function

$$\eta \log |z| = \eta(z_1, z_2) \log \left( |z_1|^2 + |z_2|^2 \right)^{\frac{1}{2}}$$

is globally defined on  $M$  and is smooth except at  $x$ . Furthermore, the  $(1, 1)$  form  $\partial\bar{\partial}(\eta \log |z|)$  is bounded from below. Since  $u$  is strictly plurisubharmonic, we can choose a sufficiently large positive constant  $C$  such that

$$v = Cu + 6\eta \log |z|$$

is strictly plurisubharmonic on  $M$ . Then, for any nonzero tangent vector  $\xi$  of type  $(1, 0)$  on  $M$ , we have

$$\left\langle \sqrt{-1}\partial\bar{\partial}v + \text{Ric}, \xi \wedge \bar{\xi} \right\rangle > 0 .$$

Now  $\bar{\partial}(\eta z_i)$ ,  $i=1,2$ , is a  $\bar{\partial}$  closed  $(0, 1)$  form on the complete Kähler manifold  $M$ . Using the standard  $L^2$ -estimates of  $\bar{\partial}$  operator (c.f. Theorem 2.1 in [21]),

there exists a smooth function  $u_i$  such that

$$\bar{\partial}u_i = \bar{\partial}(\eta z_i) \quad i = 1, 2$$

and

$$\int_M |u_i|^2 e^{-v} dx \leq \frac{1}{c} \int_M |\bar{\partial}(\eta z_i)|^2 e^{-v} dx$$

where  $c$  is a positive constant satisfying

$$\langle \sqrt{-1} \partial \bar{\partial} v + \text{Ric}, \xi \wedge \bar{\xi} \rangle \geq c |\xi|^2$$

whenever  $\xi$  is a tangent vector on  $\text{Supp } \eta$ . First of all, this estimate implies that  $u_i$  is of polynomial growth as the weight function  $v$  is of logarithmic growth. Secondly, because of the singularity of  $6 \log |z|$  at  $x$ , it forces the function  $u_i$  and its first order derivative to vanish at  $x$ . Therefore, the holomorphic functions  $f_1 = u_1 - \eta z_1$  and  $f_2 = u_2 - \eta z_2$  define a local biholomorphism at  $x$ . Clearly, they are algebraically independent over  $\mathbf{C}$ . This concludes our construction.

For later use, we also point out here that, as a consequence of the above argument, the algebra  $P(M)$  separates points on  $M$ . In other words, for any  $x_1, x_2 \in M$  with  $x_1 \neq x_2$ , there exists  $f \in P(M)$  such that  $f(x_1) \neq f(x_2)$ .

Before we can state our main result in this section, we need the following definition. For a holomorphic function  $f \in P(M)$ , we define the degree of  $f$ ,  $\deg(f)$ , to be the infimum of all  $q$  for which the following inequality holds

$$|f(x)| \leq C(q) (1 + d^q(x, x_0)) \quad \text{for all } x \in M,$$

where  $x_0$  is some fixed point in  $M$  and  $C(q)$  is some positive constant depending on  $q$ .

Our main result in this section is the following uniform bound on the multiplicity of the zero divisor of a function  $f \in P(M)$  by its degree.

**Proposition 7.1**      Let  $(M, g_{\alpha\bar{\beta}})$  be a complete noncompact Kähler surface as above. For  $f \in P(M)$ , let

$$[V] = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |f|^2$$

be the zero divisor, counting multiplicity, determined by  $f$ . Then, there exists a positive constant  $C$ , independent of  $f$ , such that

$$\text{mult}([V], x) \leq C \deg(f)$$

holds for all  $x \in M$ .

**Proof.** Recall that the Ricci flow (2.1) with  $g_{\alpha\bar{\beta}}(x)$  as initial metric has a solution  $g_{\alpha\bar{\beta}}(x, t)$  for all times  $t \in [0, +\infty)$  and satisfies the following estimates

$$R(x, t) \leq \frac{C}{1+t} \quad (7.1)$$

and

$$\text{inj}(M, g_{\alpha\bar{\beta}}(\cdot, t)) \geq C_7(1+t)^{\frac{1}{2}} \quad (7.2)$$

on  $M \times [0, +\infty)$ .

Let  $d_t$  be the distance function from an arbitrary fixed point  $\bar{x}_0 \in M$  with respect to the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$ . By the standard Hessian comparison theorem (see [28]), we have, for any unit real vector  $v$  orthogonal to the radial direction  $\partial/\partial d_t$ ,

$$\frac{\sqrt{\frac{\alpha_1}{t}}d_t}{\tan\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)} \leq \text{Hess}\left(d_t^2\right)(v, v) \leq \frac{\sqrt{\frac{\alpha_1}{t}}d_t}{\tanh\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)} \quad \text{when } d_t \leq \frac{\pi}{4}\sqrt{\frac{t}{\alpha_1}}.$$

Here,  $\alpha_1$  is some positive constant depending only on the constants  $C$  and  $C_7$  in (7.1) and (7.2). Hence, for any unit vector  $\tilde{v}$ , we have

$$\frac{\sqrt{\frac{\alpha_1}{t}}d_t}{\tan\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)} \leq \text{Hess}\left(d_t^2\right)(\tilde{v}, \tilde{v}) + \text{Hess}\left(d_t^2\right)(J\tilde{v}, J\tilde{v}) \leq \frac{2\sqrt{\frac{\alpha_1}{t}}d_t}{\tanh\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)}$$

whenever  $d_t \leq \frac{\pi}{4}\sqrt{\frac{t}{\alpha_1}}$ . Since  $M$  is Kähler, the above expression is equivalent to

$$\frac{\sqrt{\frac{\alpha_1}{t}}d_t}{\tan\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)}\omega_t \leq \sqrt{-1}\partial\bar{\partial}d_t^2 \leq \frac{2\sqrt{\frac{\alpha_1}{t}}d_t}{\tanh\left(\sqrt{\frac{\alpha_1}{t}}d_t\right)}\omega_t.$$

In particular, we have

$$\frac{1}{2}\omega_t \leq \sqrt{-1}\partial\bar{\partial}d_t^2 \leq 4\omega_t \quad \text{whenever} \quad d_t \leq \frac{\pi}{4}\sqrt{\frac{t}{\alpha_1}}. \quad (7.3)$$

Here,  $\omega_t$  is the Kähler form of the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$ .

We next claim that \*

$$\sqrt{-1}\partial\bar{\partial} \log \tan \left( \sqrt{\frac{\alpha_1}{t}} \frac{d_t}{2} \right) \geq 0, \quad \text{whenever} \quad d_t \leq \frac{\pi}{4}\sqrt{\frac{t}{\alpha_1}}. \quad (7.4)$$

In fact, after recaling, we may assume that the sectional curvature of  $g_{\alpha\bar{\beta}}(\cdot, t)$  is less than 1 and  $\sqrt{\frac{\alpha_1}{t}} = 1$ . Then by the standard Hessian comparison, we have

$$\text{Hess}(d_t)(v, v) \geq \frac{1}{\tan d_t} \left( |v|_t^2 - \left\langle v, \frac{\partial}{\partial d_t} \right\rangle_t^2 \right)$$

for any vector  $v$  and  $d_t \leq \frac{\pi}{4}$ . Thus, by a direct computation,

$$\begin{aligned} & \text{Hess} \left( \log \tan \left( \frac{d_t}{2} \right) \right) (v, v) + \text{Hess} \left( \log \tan \left( \frac{d_t}{2} \right) \right) (Jv, Jv) \\ & \geq \frac{1}{(\tan d_t) \tan \left( \frac{d_t}{2} \right)} (1 + \tan d_t) |v|_t^2 \\ & \geq 0, \end{aligned}$$

which is our claim (7.4).

Now for any  $0 < b < a < \frac{\pi}{8}\sqrt{\frac{t}{\alpha_1}}$ , it follows from Stoke's theorem that

$$\begin{aligned} 0 & \leq \sqrt{-1} \int_{\{b \leq d_t \leq a\}} [V] \wedge \partial\bar{\partial} \log \tan \left( \sqrt{\frac{\alpha_1}{t}} \frac{d_t}{2} \right) \\ & = \sqrt{-1} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} \log \tan \left( \sqrt{\frac{\alpha_1}{t}} \frac{d_t}{2} \right) \\ & \quad - \sqrt{-1} \int_{\{d_t=b\}} [V] \wedge \bar{\partial} \log \tan \left( \sqrt{\frac{\alpha_1}{t}} \frac{d_t}{2} \right). \end{aligned}$$

---

\*we are grateful to Professor L.F.Tam for this suggestion

Then, it is no hard to see that for  $0 < b < a < \frac{\pi}{8}\sqrt{\frac{t}{\alpha_1}}$ ,

$$\frac{\sqrt{-1}}{a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) \geq \frac{1}{2} \cdot \frac{\sqrt{-1}}{b^2} \int_{\{d_t=b\}} [V] \wedge \bar{\partial} (d_t^2) . \quad (7.5)$$

Using Stoke's theorem on the right hand side of (7.5) and letting  $b \rightarrow 0$  it follows from the inequality of Bishop–Lelong that

$$\frac{\sqrt{-1}}{a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) \geq \alpha_2 \text{mult} ([V], \bar{x}_0) , \quad (7.6)$$

for some positive absolute constant  $\alpha_2$ .

Then by (7.3), (7.5), (7.6) and Stoke's theorem, we have

$$\begin{aligned} & \frac{1}{a^2} \int_{B_t(\bar{x}_0, a) \setminus B_t(\bar{x}_0, \frac{a}{2})} [V] \wedge \omega_t \\ \geq & \frac{1}{4a^2} \int_{B_t(\bar{x}_0, a) \setminus B_t(\bar{x}_0, \frac{a}{2})} [V] \wedge \sqrt{-1} \partial \bar{\partial} (d_t^2) \\ = & \frac{\sqrt{-1}}{4} \left( \frac{1}{a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) - \frac{1}{4 \cdot (\frac{a}{2})^2} \int_{\{d_t=\frac{a}{2}\}} [V] \wedge \bar{\partial} (d_t^2) \right) \\ = & \frac{\sqrt{-1}}{8} \left( \frac{1}{a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) - \frac{1}{2 \cdot (\frac{a}{2})^2} \int_{\{d_t=\frac{a}{2}\}} [V] \wedge \bar{\partial} (d_t^2) \right) \\ & + \frac{\sqrt{-1}}{8a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) \\ \geq & \frac{\sqrt{-1}}{8a^2} \int_{\{d_t=a\}} [V] \wedge \bar{\partial} (d_t^2) \\ \geq & \frac{\alpha_2}{8} \text{mult} ([V], \bar{x}_0) \end{aligned} \quad (7.7)$$

for  $0 < a < \frac{\pi}{8}\sqrt{\frac{t}{\alpha_1}}$ .

For the function  $f \in P(M)$ , let  $\tilde{x}_0$  be a point close to  $\bar{x}_0$  such that  $f(\tilde{x}_0) \neq 0$ . By definition, for any  $\delta > 0$ , there exists a constant  $C(\delta) > 0$  such that

$$|f(x)| \leq C(\delta) \left( 1 + d_0^{\deg(f)+\delta}(x, \tilde{x}_0) \right) \quad \text{on } M . \quad (7.8)$$

By equation (2.1) and estimate (7.1), we have

$$\begin{aligned} \frac{\partial g_{\alpha\bar{\beta}}(\cdot, t)}{\partial t} &\geq -R(\cdot, t)g_{\alpha\bar{\beta}}(\cdot, t) \\ &\geq -\frac{C}{1+t}g_{\alpha\bar{\beta}}(\cdot, t), \end{aligned}$$

which implies that

$$g_{\alpha\bar{\beta}}(\cdot, 0) \leq (1+t)^C g_{\alpha\bar{\beta}}(\cdot, t) \quad \text{for any } t > 0.$$

Hence, (7.8) becomes

$$|f(x)| \leq C(\delta) \left\{ 1 + \left[ (1+t)^{\frac{C}{2}} d_t(x, \tilde{x}_0) \right]^{\deg(f)+\delta} \right\} \quad \text{on } M. \quad (7.9)$$

We now fix  $t = \frac{\alpha_1}{\pi^2} 4^{K+8}$  for each positive interger  $K$ . Set

$$v_K(x) = \int_{B_t(\tilde{x}_0, 2^K)} -G_t^{(K)}(x, y) \Delta_t \log |f(y)|^2 \cdot \omega_t^2(y),$$

where  $G_t^{(K)}$  is the positive Green function with value zero on the boundary  $\partial B_t(\tilde{x}_0, 2^K)$  with respect to the metric  $g_{\alpha\bar{\beta}}(\cdot, t)$ . The function  $\log |f|^2 - v_K$  is then harmonic on  $B_t(\tilde{x}_0, 2^K)$ . From the maximum principle and (7.9), we have

$$\begin{aligned} \log \left( |f(\tilde{x}_0)|^2 \right) - v_K(\tilde{x}_0) &\leq \sup_{x \in \partial B_t(\tilde{x}_0, 2^K)} \log |f(x)|^2 \\ &\leq C_{19} K (\deg(f) + \delta) + C'(\delta) \end{aligned} \quad (7.10)$$

for some positive constants  $C_{19}, C'(\delta)$  independent of  $K, f$ , and  $\bar{x}_0$ .

On the other hand, since the volume growth condition (i) is preserved for all times, by virtue of (6.4) (c.f. Proposition 1.1 in [21]), we have

$$\begin{aligned} -v_K(\tilde{x}_0) &\geq \frac{1}{C_9} \int_{B_t(\tilde{x}_0, 2^K)} \frac{1}{d_t^2(x, \tilde{x}_0)} \Delta_t \log |f(x)|^2 \cdot \omega_t^2(x) \\ &\geq \frac{1}{C_9} \sum_{j=1}^K \left( \frac{1}{2^j} \right)^2 \int_{B_t(\tilde{x}_0, 2^j) \setminus B_t(\tilde{x}_0, 2^{j-1})} \Delta_t \log |f(x)|^2 \cdot \omega_t^2(x). \end{aligned}$$

Then, by (7.7) and the fact that  $\tilde{x}_0$  is arbitrarily close to  $\bar{x}_0$ ,

$$-v_K(\tilde{x}_0) \geq C_{20} K \text{mult}([V], \bar{x}_0) \quad (7.11)$$

for some positive constant  $C_{20}$  independent of  $K$ ,  $f$  and  $\bar{x}_0$ .

Therefore, by combining (7.10) and (7.11) and letting  $K \rightarrow +\infty$  and then  $\delta \rightarrow 0$ , we obtain

$$\text{mult}([V], \bar{x}_0) \leq C_{21} \deg(f)$$

where  $C_{21}$  is some positive constant independent of  $f$  and  $\bar{x}_0$ .  $\square$

A modified version of the proof of Proposition 7.1 gives the uniform bound on the number of irreducible components of  $[V]$ .

**Proposition 7.2** Suppose  $(M, g_{\alpha\bar{\beta}})$  is a complete noncompact Kähler surface as assumed in Proposition 7.1. Let  $f$  be a holomorphic function of polynomial growth,

$$[V] = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log |f|^2$$

be the corresponding zero divisor determined by  $f$ . Then the number of irreducible components of  $[V]$  is not bigger than  $C \deg(f)$  for the same positive constant  $C$  as in Proposition 7.1.

**Proof.** Let  $g_{\alpha\bar{\beta}}(\cdot, t)$  be the evolving metric to the Ricci flow with  $g_{\alpha\bar{\beta}}(\cdot)$  as the initial metric and  $[V_1], [V_2], \dots, [V_l]$  be any  $l$  distinct irreducible components of  $[V]$ . Fix a constant  $a > 0$  such that the intersection of the smooth points of  $[V_i]$  with  $B_0(\bar{x}_0, a)$  is nonempty for each  $0 \leq i \leq l$ .

Choose  $t = \frac{\alpha_1}{\pi^2} 4^{K+8} a^2$  for each positive integer  $K$ . As the manifold  $M$  is Stein by Theorem 5.1, each  $[V_i]$  must be noncompact. Hence, for  $j = 1, 2, \dots, K$ , we have

$$[V_i] \cap \left( B_t(\bar{x}_0, 2^j a) \setminus B_t(\bar{x}_0, 2^{j-1} a) \right) \neq \emptyset$$

and there exists a point  $x_j \in [V_i]$  with  $d_t(x_j, \bar{x}_0) = \frac{3}{2} 2^{j-1} a$  in the middle of  $B_t(\bar{x}_0, 2^j a) \setminus B_t(\bar{x}_0, 2^{j-1} a)$ . The triangle inequality says

$$B_t(x_j, 2^{j-2} a) \subset \left( B_t(\bar{x}_0, 2^j a) \setminus B_t(\bar{x}_0, 2^{j-1} a) \right) .$$



Applying a slight variant of (7.7) to  $[V_i]$ , we have

$$\begin{aligned} \frac{1}{(2^{j-2}a)^2} \int_{B_t(x_j, 2^{j-2}a)} [V_i] \wedge \omega_t &\geq \frac{\alpha_2}{8} \text{mult}([V_i], x_j) \\ &\geq \frac{\alpha_2}{8} . \end{aligned}$$

Since  $\sum_{i=1}^l [V_i]$  is only a part of the divisor  $[V]$ , we get

$$\frac{1}{(2^{j-2}a)^2} \int_{B_t(\bar{x}_0, 2^j a) \setminus B_t(\bar{x}_0, 2^{j-1}a)} \Delta_t \log |f(x)|^2 \cdot \omega_t^2(x) \geq \frac{\alpha_2}{8} l .$$

The subsequent argument is then exactly as in the proof of Proposition 7.1. In the end, we have

$$C_{20}K \cdot l \leq -\log(|f(\tilde{x}_0)|^2) + C_{19}K(\deg(f) + \delta) + C'(\delta) .$$

Letting  $K \rightarrow +\infty$  and then  $\delta \rightarrow 0$ , we get the desired estimate.  $\square$

## §8. Proof of the main theorem

In this section, we will basically follow the approach of Mok in [21], [23] to accomplish the proof of the main theorem. Let  $M$  be a Kähler surface as assumed in the Main Theorem. Recall that  $P(M)$  stands for the algebra of holomorphic functions of polynomial growth on  $M$ . Let  $R(M)$  be the quotient field of  $P(M)$ . By an abuse of terminology, we will call it the field of rational functions on  $M$ .

In the previous section, we showed that there exist two functions  $f_1, f_2 \in P(M)$  giving local holomorphic coordinates at any given point  $x \in M$ , and that the algebra  $P(M)$  separates points on  $M$ . Moreover, we obtained the following basic multiplicity estimate

$$\text{mult}([V], x) \leq C \deg(f) \tag{8.1}$$

for all  $x \in M$  and  $f \in P(M)$ , where

$$[V] = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |f|^2$$

is the zero divisor of  $f$  and  $C$  is a constant independent of  $f$  and  $x$ . Thus, by combining these facts with the classical arguments of Poincaré and Siegel, we have (c.f. the proof of Proposition 5.1 in [21])

$$\dim_{\mathbf{C}} H_p \leq 10^3 C p^2, \quad (8.2)$$

where  $H_p$  denotes the vector space of all holomorphic functions with degree  $\leq p$ , and the field of rational functions  $R(M)$  is a finite extension field over  $\mathbf{C}(f_1, f_2)$  for some algebraically independent holomorphic functions  $f_1, f_2 \in P(M)$  over  $\mathbf{C}$ . By the primitive element theorem, we can then write

$$R(M) = \mathbf{C} \left( f_1, f_2, \frac{f_3}{f_4} \right)$$

for some  $f_3, f_4 \in P(M)$ .

Now, consider the mapping  $F : M \rightarrow \mathbf{C}^4$  defined by

$$F = (f_1, f_2, f_3, f_4) .$$

Since  $R(M)$  is a finite extension field of  $\mathbf{C}(f_1, f_2)$ ,  $f_3$  and  $f_4$  satisfy equations of the form

$$f_3^p + \sum_{j=0}^{p-1} P_j(f_1, f_2) f_3^j = 0 ,$$

$$f_4^q + \sum_{j=0}^{q-1} Q_j(f_1, f_2) f_4^j = 0 ,$$

where  $P_j(w_1, w_2), Q_j(w_1, w_2)$  are rational functions of  $w_1, w_2$ . After clearing denominators, we see that  $f_1, f_2, f_3, f_4$  satisfy polynomial equations

$$P(f_1, f_2, f_3, f_4) = 0 \quad \text{and} \quad Q(f_1, f_2, f_3, f_4) = 0.$$

Let  $Z_0$  be the subvariety of  $\mathbf{C}^4$  defined by

$$Z_0 = \left\{ (w_1, w_2, w_3, w_4) \in \mathbf{C}^4 \left| \begin{array}{l} P(w_1, w_2, w_3, w_4) = 0 \\ Q(w_1, w_2, w_3, w_4) = 0 \end{array} \right. \right\},$$

and let  $Z$  be the connected component of  $Z_0$  containing  $F(M)$ . It is clear that  $\dim_{\mathbf{C}} Z = 2$ .

In the following we will show that  $F$  is an "almost injective" and "almost surjective" map to  $Z$  and we can desingularize  $F$  to obtain a biholomorphic map from  $M$  onto a quasi-affine algebraic variety by adjoining a finite number of holomorphic functions of polynomial growth.

First of all, we claim that  $Z$  is irreducible and  $F$  is "almost injective", i.e., there exists a subvariety  $V$  of  $M$  such that  $F|_{M \setminus V} : M \setminus V \rightarrow Z$  is an injective locally biholomorphic mapping. Indeed, take  $V$  to be the union of  $F^{-1}(\text{Sing}(Z))$  and the branching locus of  $F$ , here  $\text{Sing}(Z)$  denotes the singular set of  $Z$ . It is clear that  $F$  is locally biholomorphic on  $M \setminus V$ . That  $F$  is also injective there follows from the fact that  $P(M)$  separates points and  $f_1, \dots, f_4$  generate  $P(M)$ . To see the irreducibility of  $Z$ , note that  $M \setminus F^{-1}(\text{Sing}(Z))$  is connected and hence  $\overline{F(M \setminus F^{-1}(\text{Sing}(Z)))}$  is irreducible (as its set of smooth points is connected). Since  $F(M) \subset \overline{F(M \setminus F^{-1}(\text{Sing}(Z)))}$ , by the definition of  $Z$ , it must be irreducible.

Next, we come to the "almost surjectivity" of  $F$ , i.e., there exists an algebraic subvariety  $T$  of  $Z$  such that  $F(M)$  contains  $Z \setminus T$ . The method of Mok[21] in proving the almost surjectivity of  $F$  is to solve an ideal problem for each  $x \in Z \setminus T_0$  missed by  $F$ , where  $T_0$  is some fixed algebraic subvariety of  $Z$  containing the singular set of  $Z$ . The solution of the ideal problem gives a holomorphic function  $f_x \in P(M)$  with degree bounded independent of  $x$  which will correspond to a rational function on  $\mathbf{C}^4$  with pole set passes through  $x$ . Then, the almost surjectivity of  $F$  follows. Otherwise, one could select an infinite number of linearly independent  $f_x$ 's contradicting the finite

dimensionality of the space of holomorphic functions with polynomial growth of some fixed degree, c.f. (8.2).

In [21], Mok used the solution  $u$  of the Poincaré–Lelong equation as the weight function in the Skoda’s estimates for solving the ideal problem. In his case, because of his curvature quadratic decay condition, the growth of  $u$  is bounded both from above and below by the logarithm of the distance function on  $M$ . This does not work in our case because we do not have the luxury of the lower bound of  $u$ . However, thanks to the Steinness of  $M$  by Theorem 5.1, we can adapt the argument of Mok in [23] to choose another weight function by resorting to Oka’s theory of pseudoconvex Riemann domains.

Before we carry out the above procedures in proving the almost surjectivity of  $F$ . We first need to construct a nontrivial holomorphic  $(2, 0)$  vector field of polynomial growth on  $M$ .

Consider the anticanonical line bundle,  $\mathbf{K}^{-1}$ , on  $M$  equipped with the induced Hermitian metric, its curvature form  $\Omega(\mathbf{K}^{-1})$  is then simply the Ricci form of  $M$ . Let  $u$  be the strictly plurisubharmonic function of logarithmic growth obtained in Proposition 6.2. For any given point  $\bar{x}_0 \in M$ , let  $\{z_1, z_2\}$  be local holomorphic coordinates at  $\bar{x}_0$ . Choose a smooth cutoff function  $\eta$  supporting in this local holomorphic coordinate chart with value 1 in a neighborhood of  $\bar{x}_0$ . We study the following  $\bar{\partial}$  equation for the sections of  $\mathbf{K}^{-1}$  on  $M$ ,

$$\bar{\partial}S = \bar{\partial} \left( \eta \frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_2} \right). \quad (8.3)$$

Clearly, we can choose  $k > 0$  large enough such that

$$k\sqrt{-1}\partial\bar{\partial}u + \Omega(\mathbf{K}^{-1}) + 3\sqrt{-1}\partial\bar{\partial} \left( \eta \log (|z_1|^2 + |z_2|^2) \right) > 0 .$$

Then by the standard  $L^2$  estimate of  $\bar{\partial}$  operator on Hermitian holomorphic line bundles (c.f. Theorem 1.2 in [23]), equation (8.3) has a smooth solution

$S(x)$  satisfying the estimate

$$\begin{aligned}
& \int_M |S|^2 e^{-ku-3\eta \log(|z_1|^2+|z_2|^2)} \omega^2 \\
& \leq C \int_M \left| \bar{\partial} \left( \eta \frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_2} \right) \right|^2 e^{-ku-3\eta \log(|z_1|^2+|z_2|^2)} \omega^2 \\
& < +\infty
\end{aligned} \tag{8.4}$$

for some positive constant  $C$ . Recall the Poincaré–Lelong equation for the section  $S(x)$  of the anticanonical line bundle  $\mathbf{K}^{-1}$ ,

$$\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |S|^2 = [V] - \frac{1}{2\pi} \text{Ric} \quad \text{on } M,$$

where  $[V]$  is the zero divisor of  $S(x)$  (c.f. [23]). Thus,  $\log |S|^2 + u$  is subharmonic and so is  $|S|^2 e^u = \exp(\log |S|^2 + u)$ . Since  $M$  has positive Ricci curvature and maximal volume growth, we can apply the mean value inequality of subharmonic functions, (8.4) and the fact that  $u$  has logarithmic growth to show that  $S(x)$  is of polynomial growth. Set

$$v = \eta \left( \frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_2} \right) - S.$$

Then,  $v$  is a nontrivial holomorphic  $(2, 0)$  vector field over  $M$  with polynomial growth we desired.

Now, for any  $f_i, f_j \in \{f_1, f_2, f_3, f_4\}$  with  $df_i \wedge df_j \not\equiv 0$ , we can choose the point  $\bar{x}_0$  in the above construction of  $v$  so that the holomorphic function  $f_{ij}$  defined by

$$f_{ij} = \langle v, df_i \wedge df_j \rangle \tag{8.5}$$

is a nontrivial holomorphic function of polynomial growth. Here, we have used the fact that  $\|df_i \wedge df_j\|$  grows at most polynomially by the gradient estimate of harmonic functions of Yau [36]. It is obvious that the zero divisor of  $df_i \wedge df_j$  is contained in the zero divisor of  $f_{ij}$ , for which we denote by  $V_0$ . Since  $M$  is Stein, the same is also true for  $M \setminus V_0$ .

Denote by  $\pi_{ij} : Z \rightarrow \mathbf{C}^2$  the projection map given by  $(w_1, w_2, w_3, w_4) \mapsto (w_i, w_j)$ . Then, the map

$$\rho = \pi_{ij} \circ F : M \setminus V_0 \rightarrow \mathbf{C}^2$$

realises the Stein manifold  $M \setminus V_0$  as a Riemann domain of holomorphy over  $\mathbf{C}^2$ . Let  $\delta(x)$  be the Euclidean distance to the boundary as in Oka [26]. Then,  $-\log \delta$  is a plurisubharmonic function on  $M \setminus V_0$  by a theorem of Oka [26].  $\delta(x)$  will be used in the weight function of the Skoda's estimate mentioned above. It is essential to estimate it from below in terms of the intrinsic distance  $d(x, x_0)$  on  $M$ .

**Lemma 8.1**      There exist positive constants  $p$  and  $C$  such that

$$\delta(x) \geq C |f_{ij}(x)|^2 (d(x, x_0) + 1)^{-p} .$$

**Proof.**      Let  $v_i, v_j$  be two holomorphic vector fields on  $M \setminus V_0$  defined by

$$\langle v_k, df_l \rangle = \delta_{kl} , \quad k, l = i, j .$$

By the Cramer's rule, we have

$$|v_k| \leq \frac{|df_i| + |df_j|}{|df_i \wedge df_j|} \leq \frac{|v| (|df_i| + |df_j|)}{|f_{ij}|} \leq C_{22} \frac{(d(x, x_0) + 1)^{k_1}}{|f_{ij}(x)|} \quad \text{on } M \setminus V_0,$$

for  $k = i, j$  and some positive constants  $C_{22}, k_1$ .

Since  $f_{ij}$  is of polynomial growth,  $|\nabla f_{ij}|$  is also of polynomial growth by the gradient estimate of Yau, i.e.,

$$\max \{ |f_{ij}(x)|, |\nabla f_{ij}(x)| \} \leq C_{23} (d(x, x_0) + 1)^{k_2} \quad \text{on } M ,$$

for some positive constants  $C_{23}$  and  $k_2$ . Take  $x \in M \setminus V_0$ , then for any  $y \in B(x, |f_{ij}(x)| / 3C_{23} (d(x, x_0) + 1)^{k_2})$ , we have

$$\begin{aligned} |f_{ij}(y)| &\geq |f_{ij}(x)| - C_{23} (d(x, x_0) + 2)^{k_2} \cdot \frac{|f_{ij}(x)|}{3C_{23} (d(x, x_0) + 1)^{k_2}} \\ &\geq \frac{1}{2} |f_{ij}(x)| . \end{aligned} \tag{8.6}$$

This implies

$$B\left(x, |f_{ij}(x)| / 3C_{23} (d(x, x_0) + 1)^{k_2}\right) \subset M \setminus V_0$$

and

$$|v_k(y)| \leq 2C_{22} \frac{(d(x, x_0) + 1)^{k_1}}{|f_{ij}(x)|}, \quad (8.7)$$

for all  $y \in B\left(x, |f_{ij}(x)| / 3C_{23} (d(x, x_0) + 1)^{k_2}\right)$ ,  $k = i, j$ .

By the definition of  $\delta(x)$ , it suffices to prove

$$\begin{aligned} & \rho\left(B\left(x, |f_{ij}(x)| / 6C_{23} (d(x, x_0) + 1)^{k_2}\right)\right) \\ & \supset B_{\mathbf{C}^2}\left(\rho(x), C_{24} |f_{ij}(x)|^2 / (d(x, x_0) + 1)^{k_1+k_2}\right), \end{aligned} \quad (8.8)$$

for some positive constant  $C_{24}$ . Here,  $B_{\mathbf{C}^2}(a, r)$  denotes the Euclidean ball in  $\mathbf{C}^2$  with center  $a$  and radius  $r$ .

Consider the real vector field

$$\begin{aligned} \xi &= \alpha_i (2\operatorname{Re}(v_i)) + \alpha_j (2\operatorname{Re}(v_j)) + \beta_i (2\operatorname{Im}(v_i)) + \beta_j (2\operatorname{Im}(v_j)) \\ &= (\alpha_i - \sqrt{-1}\beta_i) v_i + (\alpha_j - \sqrt{-1}\beta_j) v_j + (\alpha_i + \sqrt{-1}\beta_i) \bar{v}_i \\ &\quad + (\alpha_j + \sqrt{-1}\beta_j) \bar{v}_j \end{aligned}$$

with  $|\alpha_i|^2 + |\alpha_j|^2 + |\beta_i|^2 + |\beta_j|^2 = 1$ . Clearly  $\xi$  also satisfies (8.7). Let  $\gamma_\xi(\tau)$  be the integral curve in  $M$  defined by  $\xi$  and passes through  $x$ , i.e.,

$$\begin{cases} \frac{d\gamma_\xi(\tau)}{d\tau} = \xi \\ \gamma_\xi(0) = x. \end{cases} \quad (8.9)$$

We have

$$\begin{aligned} \frac{d(f_i \circ \gamma_\xi(\tau))}{d\tau} &= \langle \xi, df_i \rangle = \alpha_i - \sqrt{-1}\beta_i, \\ \frac{d(f_j \circ \gamma_\xi(\tau))}{d\tau} &= \langle \xi, df_j \rangle = \alpha_j - \sqrt{-1}\beta_j, \end{aligned}$$

and

$$|f_i \circ \gamma_\xi(\tau) - f_i(x)|^2 + |f_j \circ \gamma_\xi(\tau) - f_j(x)|^2 = \tau^2. \quad (8.10)$$

Note that (8.10) implies that  $\gamma_\xi(\tau)$  cannot always stay in

$$B\left(x, |f_{ij}(x)| / 6C_{23} (d(x, x_0) + 1)^{k_2}\right),$$

otherwise  $F = (f_1, f_2, f_3, f_4)$  would become unbounded in this ball. Denote by  $\tau_0$  the first time when  $\gamma_\xi(\tau)$  touches the boundary

$$\partial B\left(x, |f_{ij}(x)| / 6C_{23} (d(x, x_0) + 1)^{k_2}\right),$$

it is easy to see that

$$\begin{aligned} \frac{|f_{ij}(x)|}{6C_{23} (d(x, x_0) + 1)^{k_2}} &\leq \text{the length of } \gamma_\xi \text{ on } [0, \tau_0] \\ &\leq 2C_{22} \int_0^{\tau_0} \frac{(d(x, x_0) + 1)^{k_1}}{|f_{ij}(x)|} d\tau \quad (\text{by (8.7)}) \\ &= 2C_{22}\tau_0 \frac{(d(x, x_0) + 1)^{k_1}}{|f_{ij}(x)|}. \end{aligned}$$

Thus,

$$\tau_0 \geq \frac{|f_{ij}(x)|^2}{2C_{22}C_{23}6 (d(x, x_0) + 1)^{k_1+k_2}}. \quad (8.11)$$

Note that the integral curve  $\gamma_\xi$  projects to straight line passing through  $f(x)$  by  $\rho$ . Thus, when  $(\alpha_i, \alpha_j, \beta_i, \beta_j)$  runs through the unit sphere in  $\mathbf{C}^2$ , the collection of integral curves  $\gamma_\xi$  inside

$$B\left(x, |f_{ij}(x)| / 6C_{23} (d(x, x_0) + 1)^{k_2}\right)$$

will project, by  $\rho$ , onto the Euclidean ball

$$B_{\mathbf{C}^2}\left(\rho(x), |f_{ij}(x)|^2 / 2C_2 2C_2 6 (d(x, x_0) + 1)^{k_1+k_2}\right).$$

This proves (8.8) and hence the lemma.  $\square$

Now, we are ready to prove the almost surjectivity of the holomorphic map  $F : M \rightarrow \mathbf{C}^4$ . For each  $1 \leq i, j \leq 4$ , since  $f_{ij}$  is a holomorphic function of polynomial growth and  $R(M)$  is generated by  $f_1, \dots, f_4$ , we can write

$$f_{ij}(x) = H_{ij}(f_1(x), f_2(x), f_3(x), f_4(x)) \quad \text{on } M,$$



for some rational function  $H_{ij}$  on  $\mathbf{C}^4$ . Let  $T_0$  be the union of the singular set of  $Z$  and the zero and pole sets of all  $H_{ij}$ ,  $1 \leq i, j \leq 4$ . For any  $b \in Z \setminus (F(M) \cup T_0)$ , there exist fixed  $\{i, j\} \subset \{1, 2, 3, 4\}$  such that the projection  $\pi_{ij} : Z \rightarrow \mathbf{C}^2$  is nondegenerate at  $b$ . Since  $Z$  is algebraic, the number of points contained in  $\pi_{ij}^{-1} \circ \pi_{ij}(b)$  is less than some fixed integer  $K$  depending only on  $Z$ . By interpolation, there is a polynomial  $h_b$  of degree  $\leq K$  on  $\mathbf{C}^4$  such that  $h_b(b) = 1$ , and  $h_b(w) = 0$  for all  $w \in (\pi_{ij}^{-1} \circ \pi_{ij}(b)) \setminus \{b\}$ . We now solve on  $M \setminus V_0$  the ideal problem with unknown holomorphic functions  $g_i$  and  $g_j$ ,

$$(f_i - b_i) g_i + (f_j - b_j) g_j = (h_b \circ F)^4, \quad (8.12)$$

where  $b = (b_1, b_2, b_3, b_4)$ .

Let

$$\psi = -n_1 \log \delta + n_2 \log(1 + |f_i|^2 + |f_j|^2),$$

where the integers  $n_1, n_2 > 0$  will be determined later. Clearly,  $\psi$  is a strictly plurisubharmonic function on  $M \setminus V_0$ . By the estimate of Skoda (c.f. Theorem 1.3 in [23]), given any  $\alpha > 1$ , there exists a solution  $\{g_i, g_j\}$  to (8.12) such that

$$\begin{aligned} & \int_{M \setminus V_0} \frac{(|g_i|^2 + |g_j|^2) e^{-\psi}}{(|f_i - b_i|^2 + |f_j - b_j|^2)^{2\alpha}} \rho^* dV_E \\ & \leq C_\alpha \int_{M \setminus V_0} \frac{(h_b \circ F)^8 e^{-\psi}}{(|f_i - b_i|^2 + |f_j - b_j|^2)^{2\alpha+1}} \rho^* dV_E, \end{aligned} \quad (8.13)$$

provided the right hand side is finite. Recall that  $\rho = \pi_{ij} \circ F$  and here

$$\rho^* dV_E = \pm \left( \frac{\sqrt{-1}}{2} \right)^2 df_i \wedge \bar{df}_i \wedge df_j \wedge \bar{df}_j$$

denotes the pull back of the Euclidean volume element of  $\mathbf{C}^4$ .

Let  $\{\zeta_1, \zeta_2, \dots, \zeta_m\} = \pi_{ij}^{-1} \circ \pi_{ij}(b)$  ( $m < K$ ) be the preimages of  $\pi_{ij}(b)$  with  $\zeta_1 = b$ . And let  $U_k$  ( $1 \leq k \leq m$ ) be disjoint small neighborhoods of  $\zeta_k$  ( $1 \leq$

$k \leq m$ ). The integral on the right hand side of (8.13) can be decomposed into three parts

$$\begin{aligned} RHS &= \left( \int_{F^{-1}(U_1)} + \sum_{k=2}^m \int_{F^{-1}(U_k)} + \int_{(M \setminus V_0) \setminus \cup_{k=1}^m F^{-1}(U_k)} \right) \\ &\quad \frac{(h_b \circ F)^8 e^{-\psi}}{\left(|f_i - b_i|^2 + |f_j - b_j|^2\right)^{2\alpha+1}} \rho^* dV_E \\ &= I_1 + I_2 + I_3 . \end{aligned}$$

For  $I_1$ , since  $h_b(b) = 1$  and  $\delta(x) \leq \left(|f_i - b_i|^2 + |f_j - b_j|^2\right)^{\frac{1}{2}}$ , we can choose  $n_1 \geq 2(2\alpha + 1)$  and  $U_1$  small enough so that the integral  $I_1$  is finite.

For  $I_2$ , since  $h_b(\zeta_k) = 0$  for  $2 \leq k \leq m$ , we can choose  $\alpha$  such that  $2(2\alpha + 1) < 8$  (e.g.  $\alpha = 1.4$ ). Then the integral  $I_2$  is also finite.

For  $I_3$ , we choose  $n_2 \geq 10 + 8K + n_1$ , where  $h_b$  is of degree  $\leq K$ . Then  $I_3$  can be estimated as

$$I_3 \leq C_{25} \int_{\mathbf{C}^2} \frac{1}{(1 + |w|^2)^{10}} dV_E < +\infty .$$

Hence, we have obtained a solution  $\{g_i, g_j\}$  of the ideal problem (8.12) such that

$$\int_{M \setminus V_0} \frac{\left(|g_i|^2 + |g_j|^2\right) e^{-\psi}}{\left(|f_i - b_i|^2 + |f_j - b_j|^2\right)^{2\alpha}} \rho^* dV_E < +\infty . \quad (8.14)$$

Recall from Lemma 8.1 and (8.5), we have

$$\delta(x) \geq C |f_{ij}(x)|^2 (d(x, x_0) + 1)^{-p}$$

and

$$\begin{aligned} \rho^* dV_E &= \pm \left(\frac{\sqrt{-1}}{2}\right)^2 df_i \wedge df_j \wedge \overline{df}_i \wedge \overline{df}_j \\ &\geq \frac{|f_{ij}|^2}{|v \wedge \overline{v}|} \omega^2 \\ &\geq C_{26} \frac{|f_{ij}(x)|^2}{(d(x, x_0) + 1)^{k_3}} \omega^2 \end{aligned}$$

for some positive constants  $C_{26}$  and  $k_3$ . Substituting these two inequalities into (8.14) we get

$$\int_{M \setminus V_0} \frac{(|g_i|^2 + |g_j|^2) |f_{ij}|^{2+2n_1}}{(d(x, x_0) + 1)^{k_4}} \omega^2 < +\infty \quad (8.15)$$

where  $k_4$  is some positive constant independent of  $b$  and  $i, j$ . Then, both  $g_i f_{ij}^{n_1+1}$  and  $g_j f_{ij}^{n_1+1}$  are locally square integrable. They can thus be extended holomorphically from  $M \setminus V_0$  to  $M$ . By the mean value inequality of subharmonic functions, we deduce also that they are of polynomial growth with degree bounded by some positive number  $k_5$  independent of  $b$ .

Now, recall that  $R(M) = \mathbf{C}(f_1, f_2, f_3/f_4)$ , the holomorphic functions  $g_i f_{ij}^{n_1+1}$  and  $g_j f_{ij}^{n_1+1}$  are thus rational functions of  $f_1, f_2, f_3$  and  $f_4$ . Hence, we can regard the equation (8.12) as an equation on the variety  $Z \subset \mathbf{C}^4$ , namely

$$(w_i - b_i) g_i f_{ij}^{n_1+1} + (w_j - b_j) g_j f_{ij}^{n_1+1} = H_{ij}^{n_1+1} \cdot h_b^4.$$

Since  $h_b$  is a polynomial with  $h_b(b) = 1$  and the point  $b$  lies outside of the zero and pole sets of  $H_{ij}$ , either  $g_i f_{ij}^{n_1+1}$  or  $g_j f_{ij}^{n_1+1}$ , when regarded as rational function on  $\mathbf{C}^4$ , must have a pole at  $b$ . Denote this function by  $G^0$ . Thus,  $G^0$  is a rational function on  $Z$  with  $G^0(b) = \infty$  and  $G^0 \circ F$  is a holomorphic function on  $M$  with degree  $\leq k_5$ . If  $Z \setminus (F(M) \cup T_0 \cup \text{pole sets of } G^0)$  is empty, then we are done. Otherwise, pick any  $b_1 \in Z \setminus (F(M) \cup T_0 \cup \text{pole sets of } G^0)$  and repeat the same procedure to obtain a rational function  $G^1$  on  $Z$  with  $G^1(b_1) = \infty$  and  $G^1 \circ F$  a holomorphic function on  $M$  with degree  $\leq k_5$ . Proceeding this way, we obtain a sequence of points  $\{b, b_1, b_2, \dots\}$  and rational functions  $\{G^0, G^1, G^2, \dots\}$  such that  $G^k(b_k) = \infty$  and  $G^l$  regular at  $b_k$  for  $l < k$ . So,  $\{G^0, G^1, G^2, \dots\}$  must be linearly independent over  $\mathbf{C}$ . Moreover, all of  $G^k \circ F$  are holomorphic functions with degree  $\leq k_5$ . Hence, by (8.2), the above procedure must terminate in a finite number of steps. In other words, there exists an algebraic subvariety  $T$  of  $Z$  such that  $F(M) \supset Z \setminus T$ .

Moreover,  $F$  establishes a quasi embedding from  $M$  to a quasi-affine algebraic variety. Indeed, let  $W = F^{-1}(T)$ . By the definition of  $T_0$  and the construction of  $T$ , we know that  $W \supset V$ , where  $V$  is the union of the branching locus of  $F$  and  $F^{-1}(\text{Sing}(Z))$ , and  $W$  is the zero divisor of finitely many holomorphic functions of polynomial growth. Therefore,  $F$  maps  $M \setminus W$  biholomorphically onto  $Z \setminus T$ .

Finally, to complete the proof of our Main Theorem, we have to show that the mapping  $F$  can be desingularized by adjoining a finite number of holomorphic functions of polynomial growth and taking normalization of the image.

We have constructed the mapping  $F : M \rightarrow Z$  into an affine algebraic variety which maps  $M \setminus W$  biholomorphically onto  $Z \setminus T$ . Now, we use normalization of the affine algebraic variety  $Z$  to resolve the codimension 1 singularities of  $F$ . Let  $\text{Reg}(Z)$  denote the Zariski dense subset of  $Z$  consisting of its regular points. It is well known that the normalization  $\tilde{Z}$  of  $Z$  can be obtained by taking  $\tilde{Z}$  to be the closure of the graph of  $\{Q_1, Q_2, \dots, Q_m\}$  on  $\text{Reg}(Z)$  where  $Q_i$  is a rational function which is holomorphic (or regular in the terminology of algebraic geometry) on  $\text{Reg}(Z)$ . The lifting of  $F : M \rightarrow Z$  to  $\tilde{F} : M \rightarrow \tilde{Z}$  is then given by  $\{f_1, f_2, f_3, f_4, Q_1 \circ F, \dots, Q_m \circ F\}$  where, as was shown in proposition 8.1 of Mok [21], for each  $i$ ,  $Q_i \circ F$  can be holomorphically extended to the whole manifold  $M$  as a holomorphic function of polynomial growth.

Write  $F_0 = F : M \rightarrow Z$  and denote  $\tilde{F}_0 : M \rightarrow \tilde{Z}$  the normalization of  $F_0$ . For any smooth point  $x$  on the subvariety  $W$ , by using the  $L^2$  estimates of the  $\bar{\partial}$  operator as in Section 7, one can find two holomorphic functions  $g_x^1, g_x^2$  of polynomial growth which give local holomorphic coordinates at  $x$ . Adding  $g_x^1, g_x^2$  to the map  $\tilde{F}_0$ , we get a new map  $F_1 = (\tilde{F}_0, g_x^1, g_x^2) : M \rightarrow Z_1 \subset \mathbf{C}^{6+m}$ , which is nondegenerate at  $x$ . Write the normalization of  $F_1$  as  $\tilde{F}_1 : M \rightarrow \tilde{Z}_1$  and continue in this way to get holomorphic mappings  $F_i : M \rightarrow Z_i$  and their

normalizations  $\tilde{F}_i : M \rightarrow \tilde{Z}_i$  such that

$$\tilde{W}_0 \supsetneq \tilde{W}_1 \supsetneq \dots \supsetneq \tilde{W}_i \supsetneq \dots,$$

where  $\tilde{W}_i$  is the locus of ramification of  $\tilde{F}_i$ .

Note that  $\tilde{W}_i$  contains no isolated point because  $\tilde{Z}_i$  is normal. Moreover, by Proposition 7.2,  $W$  has only finite number of irreducible components because  $W$  is the zero divisor of finitely many holomorphic functions of polynomial growth. This implies that the above procedure must terminate in a finite number of steps, say  $l$ . Thus, we get a biholomorphism  $\tilde{F}_l$  from  $M$  onto its image  $\tilde{F}_l(M) \subset \tilde{Z}_l$ . The argument in our proof of the almost surjectivity shows that  $\tilde{F}_l(M)$  can miss at most finitely many irreducible subvarieties of  $\tilde{Z}_l$ , say  $\tilde{T}_1^{(l)}, \dots, \tilde{T}_q^{(l)}$ . If  $\tilde{F}_l(M) \cap \tilde{T}_i^{(l)} \neq \emptyset$ , then it must intersect  $\tilde{T}_i^{(l)}$  in a nonempty open set because  $\tilde{F}_l$  is open. We arrange  $\tilde{T}_i^{(l)}$  so that  $\tilde{F}_l(M) \cap \tilde{T}_i^{(l)} = \emptyset$  for  $1 \leq i \leq p$  and  $\tilde{F}_l(M) \cap \tilde{T}_i^{(l)} \neq \emptyset$  for  $p + 1 \leq i \leq q$ . Note that  $\tilde{F}_l(M)$  is a Stein subset of  $\tilde{Z}_l$  because  $M$  is Stein by Theorem 5.1 and  $\tilde{F}_l$  maps  $M$  biholomorphically onto its image. By Hartog's extension theorem, every holomorphic function on  $\tilde{Z}_l \setminus \cup_{1 \leq i \leq q} \tilde{T}_i^{(l)}$  extends to  $\tilde{Z}_l \setminus \cup_{1 \leq i \leq p} \tilde{T}_i^{(l)}$ . Hence, we get a biholomorphic map from  $M$  onto a quasi-affine algebraic variety. Finally, recall that a classical theorem of Ramanujam [27] in affine algebraic geometry says that an algebraic variety homeomorphic to  $\mathbf{R}^4$  is biregular to  $\mathbf{C}^2$ . Combining this result of Ramanujam with Theorem 5.1, we deduce that  $M$  is actually biholomorphic to  $\mathbf{C}^2$ . Therefore we have completed the proof of the Main Theorem.

## References

- [1] Andreotti, A. and Vesentini, E., *Carleman estimates for the Laplacian–Beltrami operator on complex manifolds*, Publ. Math. Inst. Hantes Études Sci. **25** (1965), 81-130.

- [2] Bishop, R. L., and Goldberg, S. I., *Some implications of the generalized Gauss–Bennet theorem*, Trans. Amer. Math. Soc. **112**(1964), 508–535.
- [3] Burns, D., Shnider, S. and Wells, R. O., *On deformations of strictly pseudoconvex domain*, Invent. Math. **46** (1978), 237-253.
- [4] Cao, H. D., *On Harnack’s inequalities for the Kähler–Ricci flow*, Invent. Math. **109** (1992), 247-263.
- [5] Cheeger, J., Gromov, M., and Taylor, M., *Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds*, J. Differential Geometry **17** (1982), 15-53.
- [6] Chen, B. L. and Zhu, X. P., *Complete Riemannian manifolds with pointwise pinched curvature*, Invent. Math. **140** (2000), 423-452.
- [7] Chen, B. L. and Zhu, X. P., *On complete noncompact Kähler manifolds with positive bisectional curvature*, preprint.
- [8] Cheng, S. Y. and Yau, S. T., *On the existence of a complete Kähler metric on non–compact complex manifolds and the regularity of Fefferman’s equation*, Comm. Pure Appl. Math. **33** (1980), 507-544.
- [9] Fefferman, C., *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. **26** (1974), 1-65.
- [10] Frankel, T., *Manifolds with positive curvature*, Pacific J. Math. **11** (1968), 157-170.
- [11] Freedman, M. H., *The topology of four–dimensional manifolds*, J. Differential Geom. **17** (1982), 357-453.

- [12] Greene, R. E., and Wu, H.,  *$C^\infty$  convex functions and manifolds of positive curvature*, Acta. Math. **137** (1976), 209-245.
- [13] Greene, R. E., and Wu, H., *Analysis on non-compact Kähler manifolds*, Proc. Symp. Pure. Math. Vol **30**, Part **II**, Amer. Math. Soc. (1977).
- [14] Hamilton, R. S., *Four-manifolds with positive curvature operator*, J. Differential Geom. **24** (1986), 153-179.
- [15] Hamilton, R. S., *A compactness property for solution of the Ricci flow*, Amer. J. Math. **117** (1995), 545-572.
- [16] Hamilton, R. S., *The formation of singularities in the Ricci flow*, Surveys in Differential Geometry **2** (1995), 7-136, International Press.
- [17] Hörmander, L.,  *$L^2$ -estimates and existence theorems for the  $\bar{\partial}$ -operator*, Acta. Math. **113** (1965), 89-152.
- [18] Mabuchi, T.,  *$\mathbf{C}^3$ -actions and algebraic threefolds with ample tangent bundle*, Nagoya Math. J. **69** (1978), 33-64.
- [19] Markeo, A., *Runge families and increasing unions of Stein spaces*, Bull. Amer. Math. Soc. **82** (1976), 787-788.
- [20] Mok, N., Siu, Y. T. and Yau, S. T., *The Poincaré–Lelong equation on complete Kähler manifolds*, Compositio Math. **44** (1981), 183-218.
- [21] Mok, N., *An embedding theorem of complete Kähler manifolds of positive bisectional curvature onto affine algebraic varieties*, Bull. Soc. Math. France **112** (1984), 197-258.
- [22] Mok, N., *The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature*, J. Differential Geom. **27** (1988), 179-214.

- [23] Mok, N., *An embedding theorem of complete Kähler manifolds of positive Ricci curvature onto quasi-projective varieties*, Math. Ann. **286** (1990), 377-408.
- [24] Mori, S., *Projective manifolds with ample tangent bundles*, Ann. of Math. **100** (1979), 593-606.
- [25] Ni, L., Shi, Y. G., and Tam, L. F., *Poisson equation, Poincaré–Lelong equation and curvature decay on complete Kähler manifolds*, preprint.
- [26] Oka, K., *Domaines finis sans point critique intérieur*, Jap. J. Math. **27** (1953), 97-155.
- [27] Ramanujam, C. P., *A topological characterization of the affine plane as an algebraic variety*, Ann. of Math. **94** (1971), 69-88.
- [28] Schoen, R. and Yau, S. T., *Lectures on differential geometry*, in conference proceedings and Lecture Notes in Geometry and Topology, Volume **1**, International Press Publications, 1994.
- [29] Shi, W. X., *Deforming the metric on complete Riemannian manifold*, J. Differential Geometry **30** (1989), 223-301.
- [30] Shi, W. X., *Complete noncompact Kähler manifolds with positive holomorphic bisectional curvature*, Bull. Amer Math. Soc. **23** (1990), 437-440.
- [31] Shi, W. X., *Ricci deformation of the metric on complete noncompact Kähler manifolds*, Ph. D. thesis Harvard University, 1990.
- [32] Shi, W. X., *Ricci flow and the uniformization on complete noncompact Kähler manifolds*, J. Differential Geom. **45** (1997), 94-220.
- [33] Siu, Y. T., *Pseudoconvexity and the problem of Levi*, Bull. Amer Math. Soc. **84** (1978), 481-512.



- [34] Siu, Y. T. and Yau, S. T., *Compact Kähler manifolds of positive bisectional curvature*, Invent. Math. **59** (1980), 189-204.
- [35] To, W. K., *Quasi-projective embeddings of noncompact complete Kähler manifolds of positive Ricci curvature and satisfying certain topological conditions*, Duke Math. J. **63** (1991), no. **3**, 745-789.
- [36] Yau, S. T., *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math. **28** (1975), 201-228.
- [37] Yau, S. T., *Problem Section*, Seminar on Diff. Geom. edit by S. T. Yau, Princeton Univ. Press, 1982.
- [38] Yau, S. T., *A review of complex differential geometry*, Proc. Symp. Pure Math., Vol **52**, Part **II**, Amer. Math. Soc. (1991).