

RELATIVE 2-SEGAL SPACES

MATTHEW B. YOUNG

ABSTRACT. We introduce a relative version of the 2-Segal simplicial spaces defined by Dyckerhoff and Kapranov and Gálvez-Carrillo, Kock and Tonks. Examples of relative 2-Segal spaces include the categorified unoriented cyclic nerve, real pseudoholomorphic polygons in almost complex manifolds and the \mathcal{R}_\bullet -construction from Grothendieck-Witt theory. We show that a relative 2-Segal space defines a categorical representation of the Hall algebra associated to the base 2-Segal space. In this way, after decategorification we recover a number of known constructions of Hall algebra representations. We also describe some higher categorical interpretations of relative 2-Segal spaces.

CONTENTS

Introduction	1
1. Higher Segal spaces	4
1.1. Simplicial objects	4
1.2. 1-Segal spaces	5
1.3. 2-Segal spaces	5
1.4. The Waldhausen \mathcal{S}_\bullet -construction	6
2. Relative higher Segal spaces	8
2.1. Relative 1-Segal spaces	8
2.2. Relative 2-Segal spaces	11
2.3. Relative polyhedral subdivisions	13
2.4. The stable framed \mathcal{S}_\bullet -construction	15
2.5. Real pseudoholomorphic polygons	17
3. Relative higher Segal spaces from categories with dualities	20
3.1. Categories with duality	20
3.2. Unoriented categorified nerves	21
3.3. Unoriented categorified twisted cyclic nerves	23
3.4. The \mathcal{R}_\bullet -construction	26
4. Applications to Hall algebras	29
4.1. Categorical Hall algebra representations	29
4.2. Hall monoidal module categories	34
4.3. Modules over multivalued categories	36
4.4. Presheaves on Hall 2-categories	39
References	44

INTRODUCTION

Motivated by Segal's notion of a Γ -space [38], Rezk introduced Segal spaces in his study of the homotopy theory of $(\infty, 1)$ -categories [32]. Generalizing these ideas, for

Date: November 16, 2017.


2010 Mathematics Subject Classification. Primary: 18G30; Secondary 19G38, 16G20.

Key words and phrases. Higher Segal spaces. Categorified Hall algebra representations. Categories with duality. Grothendieck-Witt theory.

each integer $k \geq 1$, Dyckerhoff and Kapranov introduced k -Segal spaces [7]. Very roughly, a simplicial topological space X_\bullet is called k -Segal if it satisfies a collection of locality conditions governed by polyhedral subdivisions of k -dimensional cyclic polytopes. When $k = 1$, so that the locality conditions are governed by subdivisions of the interval, the 1-Segal conditions state that, for each $n \geq 2$, the canonical map to the homotopy fibre product

$$X_n \rightarrow \overbrace{X_1 \times_{X_0}^R \cdots \times_{X_0}^R X_1}^{n \text{ factors}}$$

is a weak homotopy equivalence. Hence 1-Segal spaces reduce to Rezk's Segal spaces. The 2-Segal spaces, which were introduced independently by Gálvez-Carrillo, Kock and Tonks [12] under the name decomposition spaces, obey locality conditions governed by subdivisions of convex plane polygons. The first non-trivial conditions derive from the two triangulations of the square,



and state that the induced morphisms

$$X_{\{0,1,3\}} \times_{X_{\{1,3\}}}^R X_{\{1,2,3\}} \leftarrow X_{\{0,1,2,3\}} \rightarrow X_{\{0,1,2\}} \times_{X_{\{0,2\}}}^R X_{\{0,2,3\}} \quad (1)$$

are weak homotopy equivalences. A large number of examples of 2-Segal spaces from a diverse range of subjects are described in [7] and [12].

One motivation to study 2-Segal spaces is the theory of Hall algebras. Indeed, as exploited by both Dyckerhoff and Kapranov [7] and Gálvez-Carrillo, Kock and Tonks [12], the 2-Segal conditions admit a natural interpretation as higher coherence conditions on a multiplication defined on the 1-simplices X_1 , the weak equivalences (1) imposing weak associativity. From the Hall algebra point of view, the most important example of a 2-Segal space is the Waldhausen \mathcal{S}_\bullet -construction applied to an exact category [46], or more generally an exact ∞ -category [7]. Applying suitable realization functors to this 2-Segal space recovers various familiar incarnations of the Hall algebra [33], [26], [20], [24]. However, these realizations use only the lowest 2-Segal conditions, namely (1). Taking into account the remaining conditions leads to higher categorical structures and thus to categorical Hall algebras. Applications of 2-Segal spaces to other areas, including combinatorics, topological field theories and Fukaya categories, can be found in [8], [9], [12].

Partially motivated by the representation theory of Hall algebras, in this paper we introduce relative higher Segal spaces. For an integer $k \geq 1$, a relative k -Segal space over a k -Segal space X_\bullet is a $(k-1)$ -Segal space Y_\bullet together with a morphism $Y_\bullet \rightarrow X_\bullet$ which satisfies k -dimensional locality conditions involving both X_\bullet and Y_\bullet ; by convention the 0-Segal conditions are vacuous. The simplest case is that of right relative 1-Segal spaces, the locality conditions reducing to the condition that the map $Y_\bullet \rightarrow X_\bullet$ be a right fibration of Segal spaces in the sense of [22], [3]. More interesting is the case of relative 2-Segal spaces. The relative 2-Segal conditions on a morphism $Y_\bullet \rightarrow X_\bullet$ are governed by polyhedral subdivisions of convex plane polygons with a distinguished vertex ∞ , the most basic of which are again the two

mirror symmetry admits an orientifold enhancement, relating the relative 2-Segal spaces arising from the \mathcal{R}_\bullet - and real pseudoholomorphic polygon constructions.

Finally, we describe some applications of relative higher Segal spaces to higher category theory, thus lifting some of the results of [32], [19], [7]. It is a classical fact that 1-Segal simplicial sets can be characterized as the essential image of the fully faithful nerve functor $N_\bullet : \text{Cat} \rightarrow \text{Set}_\Delta$. In a similar vein, right relative 1-Segal simplicial sets are the essential image of the relative nerve construction applied to the category of discrete right fibrations which, via the Grothendieck construction, can be interpreted as presheaves on small categories; see Proposition 2.2. Using the work of several authors [17], [25], [3], we explain a quasicategorical generalization of these statements by considering instead right relative 1-Segal combinatorial simplicial spaces and $(\infty, 1)$ -presheaves. Secondly, in [7] it is proved that the category of 2-Segal simplicial sets is equivalent to both the category of multivalued categories and to the category of \sqcup -semisimple semibicategories. Pursuing an interpretation in terms of actions of categories as in the relative 1-Segal case, we lift of these statements to the relative setting, establishing equivalences of the category of relative 2-Segal simplicial sets with both the category of modules over multivalued categories and with the category of Cat^{\sqcup} -valued presheaves on \sqcup -semisimple semibicategories; see Theorems 4.5 and 4.8, respectively.

Remark. After the first version of this paper was completed, a preprint by Tashi Walde [45] was posted to the arXiv which also aims at developing a theory of modules over higher Segal spaces. We comment where appropriate on the overlap.

Acknowledgements. The author would like to thank Ajneet Dhillon, Tobias Dyckerhoff, Joachim Kock, Karol Szumilo and Pal Zsamboki for interesting discussions regarding the subject of this paper. The author is also grateful for the comments and suggestions of an anonymous referee. The author was supported by the Direct Grants and Research Fellowship Scheme from the Chinese University of Hong Kong.

1. HIGHER SEGAL SPACES

In this section we recall, following closely [7], some required background material from the theory of higher Segal spaces.

1.1. Simplicial objects. Let Δ be the category whose objects are the non-empty finite ordinals $[n] = \{0 < \dots < n\}$, $n \geq 0$, and whose morphisms are weakly monotone set maps. Let also Δ_{aug} be the category of all finite non-empty ordinals, of which Δ is a skeleton. Denote by $\Delta_{\text{inj}} \subset \Delta$ the subcategory of injective morphisms. We sometimes consider the object $[n] \in \Delta$ as a category itself. Explicitly, the objects of $[n]$ are labelled by integers $0 \leq i \leq n$ and the morphism set $\text{Hom}_{[n]}(i, j)$ is empty if $i > j$ and consists of a single element if $i \leq j$.

A simplicial object of a category \mathcal{C} is a functor $X_\bullet : \Delta^{\text{op}} \rightarrow \mathcal{C}$. We write X_n for $X_{[n]} \in \mathcal{C}$ if it will not lead to confusion. The face and degeneracy maps of X_\bullet are denoted by

$$\partial_i : X_n \rightarrow X_{n-1}, \quad s_i : X_n \rightarrow X_{n+1}, \quad 0 \leq i \leq n.$$

More generally, a functor $X_\bullet : \Delta_{\text{inj}}^{\text{op}} \rightarrow \mathcal{C}$ is a semi-simplicial object of \mathcal{C} . A simplicial object X_\bullet admits a canonical extension to a functor $\Delta_{\text{aug}}^{\text{op}} \rightarrow \mathcal{C}$, which we continue to denote by X_\bullet . For $I \in \Delta_{\text{aug}}$, we write Δ^I for the simplicial set $\text{Hom}_\Delta(-, I)$. In particular, $\Delta^n = \Delta^{[n]}$.

Given categories \mathcal{C} and \mathcal{D} , with \mathcal{C} small, denote by $[\mathcal{C}, \mathcal{D}]$ or $\mathcal{D}^{\mathcal{C}}$ the category of functors $\mathcal{C} \rightarrow \mathcal{D}$. Let Set , Grpd and Top be the categories of sets, small groupoids and compactly generated topological spaces, respectively. Objects of the categories

$[\Delta^{\text{op}}, \text{Top}]$ and $[\Delta^{\text{op}}, \text{Grpd}]$ are called simplicial spaces and groupoids while objects of $\mathbb{S} = [\Delta^{\text{op}}, \text{Set}]$ and $\mathbb{S}_\Delta = [\Delta^{\text{op}}, \mathbb{S}]$ are called simplicial sets and combinatorial simplicial spaces, respectively.

1.2. 1-Segal spaces. Segal spaces (called 1-Segal spaces below) were introduced by Rezk [32, §4]; see also [38, §1]. The definition below is slightly different, omitting a fibrancy condition. Write $-\times_U^R-$ for the homotopy fibre product over a topological space U .

Definition. A semi-simplicial space $X_\bullet : \Delta_{\text{inj}}^{\text{op}} \rightarrow \text{Top}$ is called 1-Segal if for every $n \geq 2$ the map

$$X_n \rightarrow X_1 \times_{X_0}^R \cdots \times_{X_0}^R X_1$$

induced by the inclusions $\{i, i+1\} \hookrightarrow [n]$, $i = 0, \dots, n-1$, is a weak homotopy equivalence.

It is straightforward to verify that a semi-simplicial space X_\bullet is 1-Segal if and only if one of the following two conditions hold:

- (i) For every $n \geq 2$ and every $0 \leq i_1 < \cdots < i_l \leq n$ the map

$$X_n \rightarrow X_{i_1} \times_{X_{\{i_1\}}}^R \cdots \times_{X_{\{i_l\}}}^R X_{n-i_l}$$

induced by the inclusions $\{0, \dots, i_1\}, \dots, \{i_l, \dots, n\} \hookrightarrow [n]$ is a weak equivalence.

- (ii) For every $n \geq 2$ and every $0 \leq i \leq n$ the map

$$X_n \rightarrow X_{\{0, \dots, i\}} \times_{X_{\{i\}}}^R X_{\{i, \dots, n\}}$$

induced by the inclusions $\{0, \dots, i\}, \{i, \dots, n\} \hookrightarrow [n]$ is a weak equivalence.

With only minor changes one can formulate the theory of 1-Segal objects (along with their higher and relative variants defined below) of a combinatorial model category, in which case $-\times^R-$ becomes a homotopy limit; see [7, §5]. In particular, we can speak of 1-Segal simplicial sets, groupoids or combinatorial simplicial spaces. For simplicity we will state results in terms of simplicial spaces.

Example. The nerve $N_\bullet(\mathcal{C})$ of a small category \mathcal{C} is the simplicial set which assigns to $[n] \in \Delta$ the set underlying the category $\mathcal{C}^{[n]}$. It is well-known that $N_\bullet(\mathcal{C})$ is a 1-Segal simplicial set. In fact, the nerve functor $N_\bullet : \text{Cat} \rightarrow \mathbb{S}$ is fully faithful with essential image the 1-Segal simplicial sets. The category \mathcal{X} associated to a 1-Segal simplicial set X_\bullet has objects $\text{Ob}(\mathcal{X}) = X_0$ and morphisms

$$\text{Hom}_{\mathcal{X}}(x_0, x_1) = \{x_0\} \times_{X_{\{0\}}} X_{\{0,1\}} \times_{X_{\{1\}}} \{x_1\}.$$

Composition of morphisms is defined using the lowest 1-Segal conditions while associativity follows from the higher 1-Segal conditions. \triangleleft

Example. The categorified nerve $\mathcal{N}_\bullet(\mathcal{C})$ of a small category \mathcal{C} is the 1-Segal simplicial groupoid which assigns to $[n] \in \Delta$ the maximal groupoid of the category $\mathcal{C}^{[n]}$ [32, §3.5]. Passing to classifying spaces gives a 1-Segal simplicial space $B\mathcal{N}_\bullet(\mathcal{C})$. In Rezk's framework the categorified nerve is preferred to the ordinary nerve as the former is a complete 1-Segal space. \triangleleft

1.3. 2-Segal spaces. The 1-Segal spaces are the first in an infinite tower of higher Segal spaces introduced in [7]. In this section we focus on 2-Segal spaces, the next step in this tower. See [12] for a second approach to (unital) 2-Segal spaces.

For each integer $n \geq 2$ let $P_n \subset \mathbb{R}^2$ be a convex $(n+1)$ -gon with a total order on its vertices which is consistent with the counterclockwise orientation of \mathbb{R}^2 . The total order induces a canonical identification of the set of vertices of P_n with $[n]$. Let \mathcal{P} be a polyhedral subdivision of P_n . Associating to each polygon of \mathcal{P} its set of

vertices defines a collection of subsets of $[n]$ and hence a simplicial subset $\Delta^{\mathcal{P}} \subset \Delta^n$. Let X_{\bullet} be a semi-simplicial space. The polyhedral subdivision \mathcal{P} induces a map

$$f_{\mathcal{P}} : X_n \simeq (\Delta^n, X_{\bullet})_R \rightarrow (\Delta^{\mathcal{P}}, X_{\bullet})_R$$

where, following [7, §2.2], for a semi-simplicial set D the derived space of D -membranes of X_{\bullet} is defined to be

$$(D, X_{\bullet})_R = \operatorname{holim}_{\leftarrow \{\Delta^{\mathcal{P}} \hookrightarrow D\} \in \Delta_{\text{inj}}/D} \operatorname{Top} X_{\mathcal{P}}.$$

Definition. A semi-simplicial space X_{\bullet} is called 2-Segal if for every $n \geq 3$ and every triangulation \mathcal{T} of P_n the map $f_{\mathcal{T}} : X_n \rightarrow (\Delta^{\mathcal{T}}, X_{\bullet})_R$ is a weak equivalence.

As in the case of 1-Segal spaces, the 2-Segal conditions can be verified using coarser subdivisions. Indeed, it is proved in [7, Proposition 2.3.2] that a semi-simplicial space X_{\bullet} is 2-Segal if and only if one of the following conditions holds:

- (i) For every $n \geq 3$ and every polyhedral subdivision \mathcal{P} of P_n the map $f_{\mathcal{P}} : X_n \rightarrow (\Delta^{\mathcal{P}}, X_{\bullet})_R$ is a weak equivalence.
- (ii) For every $n \geq 3$ and every $0 \leq i < j \leq n$ the map

$$f_{\{i,j\}} : X_n \rightarrow X_{\{i,\dots,j\}} \times_{X_{\{i,j\}}}^R X_{\{0,\dots,i,j,\dots,n\}} \quad (3)$$

induced by the inclusions $\{i, \dots, j\}, \{0, \dots, i, j, \dots, n\} \hookrightarrow [n]$ is a weak equivalence.

- (iii) For every $n \geq 3$ the map (3) is a weak equivalence if $i = 0$ or $j = n$.

The following definition uses degeneracy maps and so can only be formulated in the simplicial setting.

Definition. A 2-Segal simplicial space X_{\bullet} is called unital 2-Segal if for every $n \geq 2$ and every $0 \leq i \leq n-1$ the map

$$\partial_{\{i\}} \times s_i : X_{n-1} \rightarrow X_{\{i\}} \times_{X_{\{i,i+1\}}}^R X_n$$

is a weak equivalence.

One simple construction of 2-Segal spaces is the following.

Proposition 1.1 ([7, Propositions 2.3.3, 2.5.3], [12, Proposition 3.5]). *Let X_{\bullet} be a 1-Segal semi-simplicial space. Then X_{\bullet} is 2-Segal. If in fact X_{\bullet} is a simplicial space, then X_{\bullet} is unital 2-Segal.*

1.4. The Waldhausen \mathcal{S}_{\bullet} -construction. We recall a motivating example of a unital 2-Segal space. We will work with proto-exact categories, a not necessarily additive generalization of exact categories in the sense of Quillen [31].

Definition ([7, §2.4]). *A proto-exact category is a pointed category \mathcal{C} , with zero object 0 , together with two classes of morphisms, \mathfrak{I} and \mathfrak{D} , called inflations and deflations and denoted by \rightarrow and \twoheadrightarrow , respectively, which have the following properties:*

- (i) Any morphism $0 \rightarrow U$ is in \mathfrak{I} and any morphism $U \rightarrow 0$ is in \mathfrak{D} .
- (ii) The classes \mathfrak{I} and \mathfrak{D} are closed under composition and contain all isomorphisms.
- (iii) A commutative square of the form

$$\begin{array}{ccc} U & \rightarrow & V \\ \downarrow & & \downarrow \\ W & \rightarrow & X \end{array} \quad (4)$$

is Cartesian if and only if it is coCartesian.

- (iv) Any diagram $W \rightarrow X \leftarrow V$ can be completed to a biCartesian diagram of the form (4).

(v) Any diagram $W \leftarrow U \rightarrow V$ can be completed to a biCartesian diagram of the form (4).

BiCartesian squares of the form

$$\begin{array}{ccc} U & \twoheadrightarrow & V \\ \downarrow & & \downarrow \\ 0 & \twoheadrightarrow & X \end{array}$$

are called conflations and play the role of short exact sequences in \mathcal{C} . Familiar examples of proto-exact categories include abelian and, more generally, exact categories. A more exotic example is given by the category of representations of a quiver over \mathbb{F}_1 , as described in [42].

The Waldhausen \mathcal{S}_\bullet -construction associates to a proto-exact category \mathcal{C} a simplicial groupoid $\mathcal{S}_\bullet(\mathcal{C})$ as follows [46, §1.3], [7, §2.4]. Let $\text{Ar}_n = [[1], [n]]$ be the arrow category of $[n]$. The assignment $[n] \mapsto \text{Ar}_n$ defines a cosimplicial category. An object $\{(i \rightarrow j) \mapsto A_{\{i,j\}}\}_{0 \leq i \leq j \leq n}$ of the functor category $[\text{Ar}_n, \mathcal{C}]$ is a commutative diagram in \mathcal{C} of the form

$$\begin{array}{ccccccc} A_{\{0,0\}} & \longrightarrow & A_{\{0,1\}} & \longrightarrow & \cdots & \longrightarrow & A_{\{0,n-1\}} & \longrightarrow & A_{\{0,n\}} \\ & & \downarrow & & & & \downarrow & & \downarrow \\ & & A_{\{1,1\}} & \longrightarrow & \cdots & \longrightarrow & A_{\{1,n-1\}} & \longrightarrow & A_{\{1,n\}} \\ & & & & \ddots & & \downarrow & & \downarrow \\ & & & & & & \downarrow & & \downarrow \\ & & & & & & A_{\{n-1,n-1\}} & \longrightarrow & A_{\{n-1,n\}} \\ & & & & & & & & \downarrow \\ & & & & & & & & A_{\{n,n\}}. \end{array}$$

Let $\mathcal{W}_n(\mathcal{C}) \subset [\text{Ar}_n, \mathcal{C}]$ be the full subcategory consisting of diagrams which have the following properties:

- (i) For each $0 \leq i \leq n$ the object $A_{\{i,i\}}$ is isomorphic to $0 \in \mathcal{C}$.
- (ii) All horizontal morphisms are inflations and all vertical morphisms are deflations.
- (iii) Each square that can be formed in the diagram is biCartesian.

Let $\mathcal{S}_n(\mathcal{C})$ be the maximal groupoid of $\mathcal{W}_n(\mathcal{C})$. Then $\mathcal{S}_\bullet(\mathcal{C})$ is a simplicial groupoid, the degeneracy map $s_i : \mathcal{S}_n(\mathcal{C}) \rightarrow \mathcal{S}_{n+1}(\mathcal{C})$ inserting a row/column of identity morphisms after the i th row/column and the face map $\partial_i : \mathcal{S}_n(\mathcal{C}) \rightarrow \mathcal{S}_{n-1}(\mathcal{C})$ deleting the i th row/column and composing the obvious morphisms.

Theorem 1.2 ([7, Proposition 2.4.8], [12, Theorem 10.14]). *For any proto-exact category \mathcal{C} , the simplicial groupoid $\mathcal{S}_\bullet(\mathcal{C})$ is unital 2-Segal.*

When \mathcal{C} is an exact category the simplicial space $B\mathcal{S}_\bullet(\mathcal{C})$ plays a fundamental role in the higher algebraic K -theory of \mathcal{C} . Indeed, we have $K_i(\mathcal{C}) = \pi_i \Omega |B\mathcal{S}_\bullet(\mathcal{C})|$, $i \geq 0$, where the basepoint of $|B\mathcal{S}_\bullet(\mathcal{C})|$ is taken to be $0 \in \mathcal{C}$. See [43], [46].

Remark. A variation of the Waldhausen \mathcal{S}_\bullet -construction was defined in [1], giving a functor from the category of augmented stable double categories to the category of simplicial sets. It was proved that this functor is fully faithful with essential image the unital 2-Segal simplicial sets.

2. RELATIVE HIGHER SEGAL SPACES

2.1. Relative 1-Segal spaces. Before introducing and studying relative 2-Segal spaces, which will be the main objects of interest in this paper, we study the more basic relative 1-Segal spaces.

Definition. Let X_\bullet be a 1-Segal semi-simplicial space. A morphism $F_\bullet : Y_\bullet \rightarrow X_\bullet$ of semi-simplicial spaces is called *right relative 1-Segal* if for every $n \geq 1$ and every $0 \leq i \leq n$ the outside square of the diagram

$$\begin{array}{ccc}
 Y_n & \longrightarrow & Y_{\{i, \dots, n\}} \\
 \downarrow & & \downarrow \\
 Y_{\{0, \dots, i\}} & \longrightarrow & Y_{\{i\}} \\
 F_{\{0, \dots, i\}} \downarrow & & \downarrow F_{\{i\}} \\
 X_{\{0, \dots, i\}} & \longrightarrow & X_{\{i\}}
 \end{array} \tag{5}$$

is homotopy Cartesian.

Similarly, a left relative 1-Segal space is a morphism $F_\bullet : Y_\bullet \rightarrow X_\bullet$, with X_\bullet being 1-Segal, for which the outside square of all diagrams of the form

$$\begin{array}{ccccc}
 Y_n & \longrightarrow & Y_{\{i, \dots, n\}} & \xrightarrow{F_{\{i, \dots, n\}}} & X_{\{i, \dots, n\}} \\
 \downarrow & & \downarrow & & \downarrow \\
 Y_{\{0, \dots, i\}} & \longrightarrow & Y_{\{i\}} & \xrightarrow{F_{\{i\}}} & X_{\{i\}}
 \end{array}$$

is homotopy Cartesian. All results below will be formulated for right relative 1-Segal spaces; analogous results hold for left relative 1-Segal spaces.

Example. Let X_\bullet be a 1-Segal semi-simplicial space. Then the identity morphism $1_{X_\bullet} : X_\bullet \rightarrow X_\bullet$ is both left and right relative 1-Segal. \triangleleft

Example. Let x_* be an object of a small category \mathcal{C} and let $\mathcal{C}/_{x_*}$ be the corresponding overcategory. The forgetful functor $\mathcal{C}/_{x_*} \rightarrow \mathcal{C}$ induces a simplicial morphism $N_\bullet(\mathcal{C}/_{x_*}) \rightarrow N_\bullet(\mathcal{C})$ which is right relative 1-Segal. Using instead the undercategory $_{x_*}/\mathcal{C}$ we obtain a left relative 1-Segal simplicial set $N_\bullet(_{x_*}/\mathcal{C}) \rightarrow N_\bullet(\mathcal{C})$. \triangleleft

Example. Suppose that a group G acts on a set E . Then G acts diagonally on the Cartesian product E^n , $n \geq 1$. The action groupoid $G \backslash\!\!\! \backslash E^{n+1}$ is the category with objects E^{n+1} and morphisms $\text{Hom}_{G \backslash\!\!\! \backslash E^{n+1}}(e_\bullet, e'_\bullet) = \{g \in G \mid g \cdot e_\bullet = e'_\bullet\}$. The assignment $[n] \mapsto G \backslash\!\!\! \backslash E^{n+1}$ defines a 1-Segal simplicial groupoid $\mathcal{S}_\bullet(G, E)$, the face (resp. degeneracy) maps omitting (resp. repeating) the appropriate entries of $E^{\bullet+1}$. Passing to classifying spaces yields a 1-Segal space $BS_\bullet(G, E)$, called the Hecke-Waldhausen space of (G, E) [7, §2.6].

Let $H \leq G$ be a subgroup. The inclusion $H \hookrightarrow G$ defines a simplicial morphism $\mathcal{S}_\bullet(H, E) \rightarrow \mathcal{S}_\bullet(G, E)$. At the level of classifying spaces we obtain a morphism $BS_\bullet(H, E) \rightarrow BS_\bullet(G, E)$ which is both left and right relative 1-Segal; this can be verified in much the same way as the 1-Segal property of $BS_\bullet(G, E)$. At the level of geometric realizations this map is homotopy equivalent to the induced morphism $BH \rightarrow BG$ of classifying spaces (cf. [7, Proposition 2.6.7]). \triangleleft

We have the following alternative characterization of right relative 1-Segal spaces.

Proposition 2.1. *A semi-simplicial morphism $F_\bullet : Y_\bullet \rightarrow X_\bullet$, with X_\bullet being 1-Segal, is right relative 1-Segal if and only if Y_\bullet is 1-Segal and the map*

$$(F_1, \partial_0) : Y_1 \rightarrow X_1 \times_{X_0}^R Y_0 \quad (6)$$

is a weak equivalence.

Proof. Suppose that F_\bullet is right relative 1-Segal. Taking $i = n = 1$ in diagram (5) implies that the map (6) is a weak equivalence. For arbitrary i and n , both the outside square and the bottom square (which is a degenerate version of the outside square) of diagram (5) are homotopy Cartesian. By the 2-out-of-3 property of weak equivalences the top square is therefore also homotopy Cartesian. Hence Y_\bullet is 1-Segal.

Conversely, suppose that Y_\bullet is 1-Segal and that the map (6) is a weak equivalence. Then we have the following sequence of weak equivalences:

$$\begin{aligned} Y_n &\xrightarrow{\text{w.e.}} Y_{\{0, \dots, i\}} \times_{Y_{\{i\}}}^R Y_{\{i, \dots, n\}} \\ &\xrightarrow{\text{w.e.}} Y_{\{0, 1\}} \times_{Y_{\{1\}}}^R Y_{\{1, \dots, i\}} \times_{Y_{\{i\}}}^R Y_{\{i, \dots, n\}} \\ &\xrightarrow{\text{w.e.}} X_{\{0, 1\}} \times_{X_{\{1\}}}^R Y_{\{1\}} \times_{Y_{\{1\}}}^R Y_{\{1, \dots, i\}} \times_{Y_{\{i\}}}^R Y_{\{i, \dots, n\}} \\ &\quad \vdots \\ &\xrightarrow{\text{w.e.}} X_{\{0, 1\}} \times_{X_{\{1\}}}^R \cdots \times_{X_{\{i-1\}}}^R X_{\{i-1, i\}} \times_{X_{\{i\}}}^R Y_{\{i, \dots, n\}} \\ &\xleftarrow{\text{w.e.}} X_{\{0, \dots, i\}} \times_{X_{\{i\}}}^R Y_{\{i, \dots, n\}}. \end{aligned}$$

The map $Y_n \rightarrow X_{\{0, \dots, i\}} \times_{X_{\{i\}}}^R Y_{\{i, \dots, n\}}$ makes this chain of maps commute and is thus a weak equivalence. Hence F_\bullet is right relative 1-Segal. \square

Kazhdan and Varshavsky [22] and de Brito [3] define a right Segal fibration to be a morphism of 1-Segal spaces $F_\bullet : Y_\bullet \rightarrow X_\bullet$ for which the map (6) is a weak equivalence. It follows from Proposition 2.1 that right relative 1-Segal spaces and right Segal fibrations are the same objects. Using [3, Proposition 1.10] we obtain a model category theoretic interpretation of the relative 1-Segal conditions. Namely, right relative 1-Segal objects in \mathbb{S}_Δ are the fibrant objects of a natural left Bousfield localization of $(\mathbf{Seg}_1)_{/X_\bullet}$, the overcategory model structure on $(\mathbb{S}_\Delta)_{/X_\bullet}$ induced by Rezk's 1-Segal model structure \mathbf{Seg}_1 on \mathbb{S}_Δ [32, Theorem 7.1].

Right relative 1-Segal simplicial sets admit a simple nerve theoretic characterization, analogous to that of 1-Segal simplicial sets; in Section 4.3 we will prove a similar result in the 2-Segal setting. To formulate this, let $\mathbf{DRFib} \subset \mathbf{Cat}^{[1]}$ be the full subcategory of discrete right fibrations and let $\mathbf{1-SegRelS} \subset \mathbb{S}^{[1]}$ be the full subcategory of right relative 1-Segal simplicial sets. Let also \mathbf{Psh} be the category whose objects are presheaves on small categories and whose morphisms are pairs

$$(\phi, \theta) : (\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}) \rightarrow (\mathcal{F}' : \mathcal{C}'^{\text{op}} \rightarrow \mathbf{Set}) \quad (7)$$

consisting of a functor $\phi : \mathcal{C} \rightarrow \mathcal{C}'$ and a natural transformation $\theta : \mathcal{F} \Rightarrow \mathcal{F}' \circ \phi^{\text{op}}$. The Grothendieck construction defines a functor $\int : \mathbf{Psh} \rightarrow \mathbf{Cat}^{[1]}$, assigning to the presheaf \mathcal{F} the functor $F : \int_{\mathcal{C}} \mathcal{F} \rightarrow \mathcal{C}$ and assigning to a morphism (7) the diagram

$$\begin{array}{ccc} \int_{\mathcal{C}} \mathcal{F} & \xrightarrow{\tilde{\theta}} & \int_{\mathcal{C}'} \mathcal{F}' \\ F \downarrow & & \downarrow F' \\ \mathcal{C} & \xrightarrow{\phi} & \mathcal{C}'. \end{array}$$

Recall that $\int_{\mathcal{C}} \mathcal{F}$ is the category whose objects are pairs (c, \tilde{c}) , with $c \in \mathcal{C}$ and $\tilde{c} \in \mathcal{F}(c)$, and whose morphisms $(c_1, \tilde{c}_1) \rightarrow (c_2, \tilde{c}_2)$ are morphisms $c_1 \xrightarrow{x} c_2$ which

satisfy $\mathcal{F}(x)(\tilde{c}_2) = \tilde{c}_1$. The functor $\tilde{\theta}$ assigns to $(c, \tilde{c}) \in \int_c \mathcal{F}$ the object $(\phi(c), \theta_c(\tilde{c}))$ and assigns to $x : (c_1, \tilde{c}_1) \rightarrow (c_2, \tilde{c}_2)$ the morphism $\phi(x)$.

Proposition 2.2. *The relative nerve functor*

$$N_{\bullet}^{[1]} : \text{Cat}^{[1]} \rightarrow \mathbb{S}^{[1]}, \quad (\mathcal{Y} \xrightarrow{F} \mathcal{X}) \mapsto (N_{\bullet}(\mathcal{Y}) \xrightarrow{N_{\bullet}(F)} N_{\bullet}(\mathcal{X}))$$

is fully faithful and fits into the commutative diagram of functors

$$\begin{array}{ccccc} \text{Psh} & \xrightarrow{f} & \text{Cat}^{[1]} & \xrightarrow{N_{\bullet}^{[1]}} & \mathbb{S}^{[1]} \\ & \searrow \sim & \uparrow & & \uparrow \\ & & \text{DRFib} & \xrightarrow{\sim} & \text{1-SegRelS} \end{array}$$

with indicated equivalences. In particular, there is an equivalence of categories $\text{1-SegRelS} \simeq \text{Psh}$.

Proof. That $N_{\bullet}^{[1]}$ is fully faithful is well-known; see for example [37, Proposition 2.1]. Commutativity of the triangle in the above diagram and the fact that f induces an equivalence $\text{Psh} \simeq \text{DRFib}$ are both standard. To see that $N_{\bullet}^{[1]}$ restricts as claimed, let $(F : \mathcal{Y} \rightarrow \mathcal{X}) \in \text{DRFib}$. Then $N_{\bullet}(F) : N_{\bullet}(\mathcal{Y}) \rightarrow N_{\bullet}(\mathcal{X})$ is a morphism of 1-Segal simplicial sets. The condition that F be a discrete right fibration is precisely the condition that the map $N_1(\mathcal{Y}) \rightarrow N_1(\mathcal{X}) \times_{N_{\{1\}}(\mathcal{X})} N_{\{1\}}(\mathcal{Y})$ be a bijection. Proposition 2.1 therefore implies that $N_{\bullet}(F)$ is right relative 1-Segal. The construction of a quasi-inverse $\text{1-SegRelS} \rightarrow \text{DRFib}$ is similar. \square

To end this section we explain a quasicategorical generalization of Proposition 2.2. Suppose that $X_{\bullet} \in \mathbb{S}_{\Delta}$ is a complete 1-Segal combinatorial simplicial space. The quasicategory \mathcal{X} modelled by X_{\bullet} (see [32, §5], [19, §4]) has object set the 0-simplices of X_0 and has mapping spaces

$$\text{map}_{\mathcal{X}}(x_0, x_1) = \{x_0\} \times_{X_{\{0\}}}^R X_{\{0,1\}} \times_{X_{\{1\}}}^R \{x_1\}.$$

Here $\{x\}$ is regarded as the simplicial set Δ^0 . The lowest 1-Segal conditions define, up to homotopy, a composition law

$$\text{map}_{\mathcal{X}}(x_0, x_1) \times \text{map}_{\mathcal{X}}(x_1, x_2) \rightarrow \text{map}_{\mathcal{X}}(x_0, x_2)$$

which, by the remaining 1-Segal conditions, is coherently associative. Suppose now that we are given a right relative 1-Segal morphism $F_{\bullet} : Y_{\bullet} \rightarrow X_{\bullet}$. For each object $x \in \mathcal{X}$ define a Kan complex by $\mathcal{F}(x) = \{x\} \times_{X_0}^R Y_0 \in \mathbb{S}$. The diagram

$$\begin{array}{ccc} \{x_0\} \times_{X_{\{0\}}}^R Y_{\{0,1\}} & \longrightarrow & \{x_0\} \times_{X_{\{0\}}}^R Y_{\{0\}} \\ \text{w.e.} \downarrow & & \\ \{x_0\} \times_{X_{\{0\}}}^R X_{\{0,1\}} \times_{X_{\{1\}}}^R Y_{\{1\}} & & \\ \uparrow & & \\ (\{x_0\} \times_{X_{\{0\}}}^R X_{\{0,1\}} \times_{X_{\{1\}}}^R \{x_1\}) \times (\{x_1\} \times_{X_{\{1\}}}^R Y_{\{1\}}), & & \end{array}$$

whose indicated arrow is a weak equivalence by the lowest right relative 1-Segal condition, defines up to homotopy an action map

$$\text{map}_{\mathcal{X}}(x_0, x_1) \times \mathcal{F}(x_1) \rightarrow \mathcal{F}(x_0).$$

The remaining right relative 1-Segal conditions ensure that this action is coherently associative. In this way we obtain an $(\infty, 1)$ -presheaf on \mathcal{X} . Conversely, by combining [25, Proposition 5.1.1.1] (see also [17]) and [3, Theorem 1.22] we see that, up to weak equivalence, any $(\infty, 1)$ -presheaf on \mathcal{X} arises in this way.

2.2. Relative 2-Segal spaces. In this section we give a direct definition of relative 2-Segal spaces. In Section 2.3 we will describe a second approach using polyhedral subdivisions, in line with the definition of 2-Segal spaces.

Definition. Let X_\bullet be a 2-Segal semi-simplicial space. A morphism $F_\bullet : Y_\bullet \rightarrow X_\bullet$ of semi-simplicial spaces is called relative 2-Segal if

- (1) for every $n \geq 2$ and every $0 \leq i < j \leq n$ the outside square of the diagram

$$\begin{array}{ccc}
 Y_n & \longrightarrow & Y_{\{0, \dots, i, j, \dots, n\}} \\
 \downarrow & & \downarrow \\
 Y_{\{i, \dots, j\}} & \longrightarrow & Y_{\{i, j\}} \\
 F_{\{i, \dots, j\}} \downarrow & & \downarrow F_{\{i, j\}} \\
 X_{\{i, \dots, j\}} & \longrightarrow & X_{\{i, j\}}
 \end{array} \tag{8}$$

is homotopy Cartesian, and

- (2) the simplicial space Y_\bullet is 1-Segal.

Remarks.

- (1) Morphisms $F_\bullet : Y_\bullet \rightarrow X_\bullet$ of simplicial spaces for which the first condition above holds play an important role in [12], where they are called unique lifting of factorization (ULF) functors. If, in addition, F_\bullet preserves units (see below), then F_\bullet is called a conservative ULF functor.
- (2) The above notion of relative 2-Segal space coincides with that of Walde; see [45, Proposition 3.5.10].
- (3) Rather than considering diagrams of the form (8), one could require that the outside square of all diagrams of the form

$$\begin{array}{ccccc}
 Y_n & \longrightarrow & Y_{\{0, \dots, i, j, \dots, n\}} & \xrightarrow{F_{\{0, \dots, i, j, \dots, n\}}} & X_{\{0, \dots, i, j, \dots, n\}} \\
 \downarrow & & \downarrow & & \downarrow \\
 Y_{\{i, \dots, j\}} & \longrightarrow & Y_{\{i, j\}} & \xrightarrow{F_{\{i, j\}}} & X_{\{i, j\}}
 \end{array}$$

be homotopy Cartesian. However, morphisms F_\bullet satisfying these conditions are less interesting from our perspective since, for example, they do not lead to categorified Hall algebra representations.

The following result will be helpful in verifying the relative 2-Segal conditions.

Proposition 2.3. Let $F_\bullet : Y_\bullet \rightarrow X_\bullet$ be a morphism of semi-simplicial spaces. Assume that X_\bullet is 2-Segal. The following statements are equivalent:

- (1) For every $n \geq 1$ and every $0 \leq i < j \leq n$ the outside square of diagram (8) is homotopy Cartesian.
- (2) Y_\bullet is 2-Segal and for every $n \geq 1$ and the outside square of diagram (8) is homotopy Cartesian if $i = 0$ or $j = n$.
- (3) Y_\bullet is 2-Segal and for every $n \geq 1$ the square

$$\begin{array}{ccc}
 Y_{\{0, \dots, n\}} & \longrightarrow & Y_{\{0, n\}} \\
 F_{\{0, \dots, n\}} \downarrow & & \downarrow F_{\{0, n\}} \\
 X_{\{0, \dots, n\}} & \longrightarrow & X_{\{0, n\}}
 \end{array} \tag{9}$$

is homotopy Cartesian.

Proof. Assume that the first condition holds. Since the bottom square of (8) is a degenerate version of the outer square, the 2-out-of-3 property of weak equivalences implies that Y_\bullet is 2-Segal. Hence the second condition holds. It is clear that the second condition implies the third. Assume that the third condition holds. Then the bottom square of diagram (8) is homotopy Cartesian by assumption while the top square is homotopy Cartesian since Y_\bullet is 2-Segal. It follows that the outside square is also homotopy Cartesian, showing that the first condition holds. \square

We will often show that a morphism $F_\bullet : Y_\bullet \rightarrow X_\bullet$ is relative 2-Segal by first proving that Y_\bullet is 1-Segal, applying Proposition 1.1 and then verifying the third condition of Proposition 2.3.

We briefly describe a model theoretic interpretation of relative 2-Segal spaces. Denote by \mathcal{R} the Reedy model structure on \mathbb{S}_Δ . Following [7, §5.2], let $\mathcal{S}\text{eg}_2$ be the left Bousfield localization of \mathcal{R} along the maps

$$\mathcal{S}\text{eg}_2 = \{\Delta^\mathcal{T} \hookrightarrow \Delta^n \mid n \geq 3, \mathcal{T} \text{ is a triangulation of } P_n\}.$$

Fibrant objects of $(\mathbb{S}_\Delta, \mathcal{S}\text{eg}_2)$ are then (Reedy fibrant) 2-Segal combinatorial simplicial spaces. For $X_\bullet \in \mathbb{S}_\Delta$ write $(\mathcal{S}\text{eg}_2)_{/X_\bullet}$ for the induced model structure on the overcategory $(\mathbb{S}_\Delta)_{/X_\bullet}$ and let $(\mathcal{S}\text{eg}_2)_{/X_\bullet}^{\text{fib}}$ be its left Bousfield localization along

$$\mathcal{S}\text{eg}_2^{\text{fib}} = \{\Delta^{\{0,n\}} \hookrightarrow \Delta^n \xrightarrow{x_n} X_\bullet \mid x_n \in X_n, n \geq 2\}.$$

Assuming that X_\bullet is 2-Segal, Proposition 2.3 implies that the fibrant objects of $((\mathbb{S}_\Delta)_{/X_\bullet}, (\mathcal{S}\text{eg}_2)_{/X_\bullet}^{\text{fib}})$ are the morphisms $F_\bullet : Y_\bullet \rightarrow X_\bullet$ for which the outside square of the diagrams (8) is homotopy Cartesian, that is, the ULF functors. From this point of view, a relative 2-Segal combinatorial simplicial space is a fibrant object of $((\mathbb{S}_\Delta)_{/X_\bullet}, (\mathcal{S}\text{eg}_2)_{/X_\bullet}^{\text{fib}})$ whose total space is in addition 1-Segal. It is important to note that the main results of this paper do not hold for arbitrary fibrant objects of $((\mathbb{S}_\Delta)_{/X_\bullet}, (\mathcal{S}\text{eg}_2)_{/X_\bullet}^{\text{fib}})$.

We now formulate the relative analogue of the unital conditions.

Definition. A relative 2-Segal simplicial space $F_\bullet : Y_\bullet \rightarrow X_\bullet$ is called unital relative 2-Segal if for every $n \geq 2$ and every $0 \leq i \leq n-1$ the map

$$(F_{\{i\}} \circ \partial_{\{i\}}) \times s_i : Y_{n-1} \rightarrow X_{\{i\}} \times_{X_{\{i,i+1\}}}^R Y_n$$

is a weak equivalence.

We have the following relative analogue of Proposition 1.1.

Proposition 2.4. Let $F_\bullet : Y_\bullet \rightarrow X_\bullet$ be a right relative 1-Segal semi-simplicial space. Then F_\bullet is relative 2-Segal. If, moreover, F_\bullet is a morphism of simplicial spaces, then F_\bullet is unital relative 2-Segal.

Proof. Proposition 2.1 implies that Y_\bullet is 1-Segal. The relative 2-Segal morphism $Y_{\{0,\dots,n\}} \rightarrow X_{\{0,\dots,n\}} \times_{X_{\{0,n\}}}^R Y_{\{0,n\}}$ factors as the composition

$$\begin{aligned} Y_{\{0,\dots,n\}} &\rightarrow X_{\{0,\dots,n\}} \times_{X_{\{n\}}}^R Y_{\{n\}} \\ &\rightarrow X_{\{0,\dots,n\}} \times_{X_{\{0,n\}}}^R X_{\{0,n\}} \times_{X_{\{n\}}}^R Y_{\{n\}} \\ &\rightarrow X_{\{0,\dots,n\}} \times_{X_{\{0,n\}}}^R Y_{\{0,n\}}. \end{aligned}$$

The first and third morphisms are weak equivalences by the right relative 1-Segal conditions on F_\bullet while the second morphism is a weak equivalence for trivial reasons. Hence the composition is a weak equivalence. The unital condition is verified in a similar way. \square

Example. For any 1-Segal semi-simplicial space X_\bullet , the identity morphism $\mathbf{1}_{X_\bullet}$ is relative 2-Segal. If X_\bullet is 2-Segal but not 1-Segal, then $\mathbf{1}_{X_\bullet}$ is not relative 2-Segal. \triangleleft

We end this section with a construction of relative 2-Segal spaces which can be seen as the 2-Segal analogue of the fact that the morphisms $N_\bullet(\mathcal{C}/_{x_*}) \rightarrow N_\bullet(\mathcal{C})$ and $N_\bullet(x_*/\mathcal{C}) \rightarrow N_\bullet(\mathcal{C})$, defined in Section 2.1, are right and left relative 1-Segal, respectively. Recall the left join functor

$$l : \Delta \rightarrow \Delta, \quad \{0, \dots, n\} \mapsto \{0', 0, \dots, n\}.$$

The left path space of a simplicial space X_\bullet is the composition

$$P^{\triangleleft} X_\bullet : \Delta^{\text{op}} \xrightarrow{l^{\text{op}}} \Delta^{\text{op}} \xrightarrow{X_\bullet} \text{Top}.$$

After identifying $P^{\triangleleft} X_m$ with X_{m+1} for each $m \geq 0$, the face map $\partial_i^{\triangleleft} : P^{\triangleleft} X_n \rightarrow P^{\triangleleft} X_{n-1}$ is identified with ∂_{i+1} . The remaining face maps $\partial_{0'}$ assemble to a simplicial morphism $F_\bullet^{\triangleleft} : P^{\triangleleft} X_\bullet \rightarrow X_\bullet$. Using instead the right join functor

$$r : \Delta \rightarrow \Delta, \quad \{0, \dots, n\} \mapsto \{0, \dots, n, n'\}$$

we obtain the right path space $F_\bullet^{\triangleright} : P^{\triangleright} X_\bullet \rightarrow X_\bullet$.

Proposition 2.5. *Assume that X_\bullet is (unital) 2-Segal. Then the left and right path spaces*

$$F_\bullet^{\triangleleft} : P^{\triangleleft} X_\bullet \rightarrow X_\bullet, \quad F_\bullet^{\triangleright} : P^{\triangleright} X_\bullet \rightarrow X_\bullet$$

are (unital) relative 2-Segal.

Proof. Since X_\bullet is 2-Segal, the path space criterion [7, Theorem 6.3.2], [12, Theorem 4.11] implies that $P^{\triangleleft} X_\bullet$ and $P^{\triangleright} X_\bullet$ are 1-Segal. That the remaining relative 2-Segal conditions hold is also proved in [12, Theorem 4.11]. Explicitly, the relative 2-Segal map

$$P^{\triangleleft} X_n = X_{\{0', 0, \dots, n\}} \rightarrow X_n \times_{X_{\{0, n\}}}^R X_{\{0', 0, n\}} = X_n \times_{X_{\{0, n\}}}^R P^{\triangleleft} X_{\{0, n\}}$$

and the relative unit map

$$P^{\triangleleft} X_{n-1} = X_{\{0', 0, \dots, n-1\}} \rightarrow X_{\{i\}} \times_{X_{\{i, i+1\}}}^R X_{\{0', 0, \dots, n\}} = X_{\{i\}} \times_{X_{\{i, i+1\}}}^R P^{\triangleleft} X_n$$

are 2-Segal and unit maps for X_\bullet and so are weak equivalences by assumption. \square

2.3. Relative polyhedral subdivisions. In this section we use combinatorial geometry to give a uniform treatment of the two conditions which define relative 2-Segal spaces. The construction below was suggested by the referee (see also [45, §3.2]) and replaces an earlier construction of the author in terms of polyhedral subdivisions of reflection symmetric cyclic polygons.

We require some minor modifications of the definitions from Section 1.3. For each $n \geq 1$ let $P_n^\infty \subset \mathbb{R}^2$ be a convex $(n+2)$ -gon with a total order on its vertices which is consistent with the counterclockwise orientation of \mathbb{R}^2 . There is then a canonical identification of the vertices of P_n^∞ with $[n]_\infty$, the totally ordered set $\{0 < 1 < \dots < n < \infty\}$. Note that there is a unique isomorphism $[n]_\infty \simeq [n+1]$ in Δ .

Let $F_\bullet : Y_\bullet \rightarrow X_\bullet$ be a morphism of semi-simplicial spaces. Associated to a polyhedral subdivision \mathcal{P} of P_n^∞ is the overcategory $\Delta_{\text{inj}}/\Delta^{\mathcal{P}}$. Define a functor

$$F_{\mathcal{P}} : (\Delta_{\text{inj}}/\Delta^{\mathcal{P}})^{\text{op}} \rightarrow \text{Top}$$

as follows. For each injective morphism $\sigma : \Delta^k \hookrightarrow \Delta^{\mathcal{P}}$, whose image we identify with a $(k+1)$ -element subset of $[n]_\infty$, put

$$F_{\mathcal{P}}(\sigma) = \begin{cases} X_k & \text{if } \infty \notin \sigma(\Delta^k), \\ Y_{k-1} & \text{if } \infty \in \sigma(\Delta^k). \end{cases}$$

By convention we take Y_{-1} to be a point, although Y_{-1} will not play any significant role in what follows. Given a morphism

$$\begin{array}{ccc} \Delta^k & \xrightarrow{\phi} & \Delta^{k'} \\ & \searrow \sigma & \swarrow \sigma' \\ & \Delta^{\mathcal{P}} & \end{array}$$

in $\Delta_{\text{inj}}/\Delta^{\mathcal{P}}$ define a morphism $F_{\mathcal{P}}(\phi) : F_{\mathcal{P}}(\sigma') \rightarrow F_{\mathcal{P}}(\sigma)$ of topological spaces as follows. If the images σ and σ' both contain ∞ (resp. both do not contain ∞), then $F_{\mathcal{P}}(\phi)$ is defined by the simplicial structure of Y_{\bullet} (resp. X_{\bullet}), while if only the image of σ' contains ∞ , then $F_{\mathcal{P}}(\phi)$ is defined by the morphism F_{\bullet} .

With the above notation, the relative derived membrane space is defined to be the homotopy limit

$$(\Delta^{\mathcal{P}}, F_{\bullet})_R = \text{holim}_{\leftarrow \sigma \in \Delta_{\text{inj}}/\Delta^{\mathcal{P}}}^{\text{Top}} F_{\mathcal{P}}(\sigma).$$

As in [7, Proposition 2.2.19], the definition is unchanged if instead the homotopy limit is taken over the full subcategory of $\Delta_{\text{inj}}/\Delta^{\mathcal{P}}$ on objects $\Delta^k \hookrightarrow \Delta^{\mathcal{P}}$ with $k \geq 1$. A refinement \mathcal{P} of \mathcal{P}' defines a simplicial subset $\Delta^{\mathcal{P}} \subset \Delta^{\mathcal{P}'}$ and thus a morphism $(\Delta^{\mathcal{P}'}, F_{\bullet})_R \rightarrow (\Delta^{\mathcal{P}}, F_{\bullet})_R$ of topological spaces. In particular, taking \mathcal{P}' to be the trivial subdivision and noting that $(\Delta^{[n]_{\infty}}, F_{\bullet})_R \simeq Y_n$, we obtain a morphism

$$f_{\mathcal{P}}^{F_{\bullet}} : Y_n \rightarrow (\Delta^{\mathcal{P}}, F_{\bullet})_R.$$

Proposition 2.6. *Let $F_{\bullet} : Y_{\bullet} \rightarrow X_{\bullet}$ be a morphism of semi-simplicial spaces. Assume that X_{\bullet} is 2-Segal. Then the following statements are equivalent:*

- (1) *The morphism F_{\bullet} is relative 2-Segal.*
- (2) *For every $n \geq 2$ and every polyhedral subdivision \mathcal{P} of P_n^{∞} the morphism $f_{\mathcal{P}}^{F_{\bullet}}$ is a weak equivalence.*
- (3) *For every $n \geq 2$ and every triangulation \mathcal{T} of P_n^{∞} the morphism $f_{\mathcal{T}}^{F_{\bullet}}$ is a weak equivalence.*

Proof. Assume that the second statement holds. If \mathcal{P} is the polyhedral subdivision of P_n^{∞} consisting only of the diagonal $\{i, \infty\}$, then

$$(\Delta^{\mathcal{P}}, F_{\bullet})_R \simeq Y_{\{0, \dots, i\}} \times_{Y_{\{i\}}}^R Y_{\{i, \dots, n\}}$$

while if \mathcal{P}' consists only of the diagonal $\{0, n\}$, then

$$(\Delta^{\mathcal{P}'}, F_{\bullet})_R \simeq X_{\{0, \dots, n\}} \times_{X_{\{0, n\}}}^R Y_{\{0, n\}}.$$

Hence $f_{\mathcal{P}}^{F_{\bullet}}$ and $f_{\mathcal{P}'}^{F_{\bullet}}$ reduce to the 1-Segal maps for Y_{\bullet} and the relative 2-Segal maps for F_{\bullet} , respectively, proving that F_{\bullet} is relative 2-Segal.

Conversely, assume that F_{\bullet} is relative 2-Segal and let \mathcal{P} be a polyhedral subdivision of P_n^{∞} . There exists a sequence $\mathcal{P}_0, \mathcal{P}_1, \dots, \mathcal{P}_k = \mathcal{P}$ of polyhedral subdivisions such that \mathcal{P}_0 is the trivial subdivision and \mathcal{P}_j is obtained from \mathcal{P}_{j-1} by adding a single diagonal. Then $f_{\mathcal{P}}^{F_{\bullet}}$ factors as the composition

$$Y_n \simeq (\Delta^{\mathcal{P}_0}, F_{\bullet})_R \rightarrow (\Delta^{\mathcal{P}_1}, F_{\bullet})_R \rightarrow \dots \rightarrow (\Delta^{\mathcal{P}_{k-1}}, F_{\bullet})_R \rightarrow (\Delta^{\mathcal{P}}, F_{\bullet})_R.$$

The construction of the \mathcal{P}_j ensures that each morphism in this composition is a weak equivalence, being induced by a 1-Segal map for Y_{\bullet} or a relative 2-Segal map for F_{\bullet} . It follows that $f_{\mathcal{P}}^{F_{\bullet}}$ is a weak equivalence and the second statement holds.

The equivalence of the second and third statements is proved similarly. \square

2.4. The stable framed \mathcal{S}_\bullet -construction. Motivated by the Hall algebra representations of [41], [11], in this section we modify the Waldhausen \mathcal{S}_\bullet -construction so as to construct relative 2-Segal groupoids from an abelian category together with a choice of stability condition.

Fix a field k . Let \mathcal{C} be a k -linear abelian category with Grothendieck group $K_0(\mathcal{C})$. A stability function on \mathcal{C} is a group homomorphism $Z : K_0(\mathcal{C}) \rightarrow \mathbb{C}$ such that

$$Z(A) \in \mathbb{H}_+ = \{m \exp(\sqrt{-1}\pi\phi) \mid m > 0, \phi \in (0, 1]\} \subset \mathbb{C}$$

for all non-zero objects $A \in \mathcal{C}$ [2, §2]. Let $\phi(A) \in (0, 1]$ be the phase of $Z(A)$. A non-zero object $A \in \mathcal{C}$ is called Z -semistable if $\phi(A') \leq \phi(A)$ for all non-trivial subobjects $A' \subset A$. For each $\phi \in (0, 1]$ the full subcategory $\mathcal{C}_\phi^{Z\text{-ss}} \subset \mathcal{C}$ consisting of the zero object together with all objects which are Z -semistable of phase ϕ is again abelian. We can therefore form the Waldhausen simplicial groupoid $\mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}})$, which is 2-Segal by Theorem 1.2.

We formulate a notion of framing following [41, §4]. Fix a left exact functor $\Phi : \mathcal{C} \rightarrow \mathbf{Vect}_k$ with values in the category of finite dimensional vector spaces over k . A framed object of \mathcal{C} is then a pair (M, s) consisting of an object $M \in \mathcal{C}$ and a section $s \in \Phi(M)$. A morphism of framed objects $(M, s) \rightarrow (M', s')$ is a pair $(\pi, \lambda) \in \text{Hom}_{\mathcal{C}}(M, M') \times k$ which satisfies $\Phi(\pi)(s) = \lambda s'$. A framed object (M, s) is called stable framed if M is Z -semistable and $\phi(A) < \phi(M)$ for all proper subobjects $A \subset M$ for which $s \in \Phi(A) \subset \Phi(M)$.

Example. We recall two standard examples of framings.

- (1) Let $\text{Rep}_k(Q)$ be the category of finite dimensional representations of a quiver Q . Sending a representation to its dimension vector defines a surjective group homomorphism $K_0(\text{Rep}_k(Q)) \rightarrow \mathbb{Z}^{Q_0}$. A tuple $\zeta = (\zeta_i)_{i \in Q_0} \in \mathbb{H}_+^{Q_0}$ defines a function $\mathbb{Z}^{Q_0} \rightarrow \mathbb{C}, d \mapsto \sum_{i \in Q_0} \zeta_i d_i$ which induces a stability function Z on $\text{Rep}_k(Q)$. For any $f \in \mathbb{Z}_{\geq 0}^{Q_0}$ the functor $\Phi : U \mapsto \bigoplus_{i \in Q_0} \text{Hom}_k(k^{f_i}, U_i)$ is a framing.
- (2) Let $\text{Coh}(X)$ be the category of coherent sheaves on a smooth projective curve X . Then $Z = -\text{deg} + \sqrt{-1} \cdot \text{rk}$ defines a stability function. The global sections functor $\Phi = H^0(X, -)$ is a framing. \triangleleft

Define a stable framed modification of the \mathcal{S}_\bullet -construction as follows. For each $n \geq 0$ let $\mathcal{S}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ be the maximal groupoid of the category of diagrams of the form

$$\begin{array}{ccccccc}
 A_{\{0,0\}} & \longrightarrow & A_{\{0,1\}} & \longrightarrow & \cdots & \longrightarrow & A_{\{0,n\}} & \longrightarrow & (M_0, s_0) \\
 & & \downarrow & & & & \downarrow & & \downarrow \\
 & & A_{\{1,1\}} & \longrightarrow & \cdots & \longrightarrow & A_{\{1,n\}} & \longrightarrow & (M_1, s_1) \\
 & & & & & & \downarrow & & \downarrow \\
 & & & & & & \vdots & & \vdots \\
 & & & & \ddots & & \downarrow & & \downarrow \\
 & & & & & & A_{\{n,n\}} & \longrightarrow & (M_n, s_n) \\
 & & & & & & & & \downarrow \\
 & & & & & & & & (M_{n+1}, s_{n+1})
 \end{array}$$

where each (M_i, s_i) is a framed object and which have the following properties:

- (i) Upon forgetting the framing data s_0, \dots, s_{n+1} the resulting diagram is an object of $\mathcal{S}_{n+1}(\mathcal{C}_\phi^{Z\text{-ss}})$.
- (ii) Each pair (M_i, s_i) , $i = 0, \dots, n$, is a stable framed object.

In the above diagram a morphism $A_{\{i,n\}} \rightarrow (M_i, s_i)$ is simply a morphism of the underlying objects of \mathcal{C} . The groupoids $\mathcal{S}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ assemble to a simplicial groupoid. There is a canonical simplicial morphism $F_\bullet : \mathcal{S}_\bullet^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}})$ which forgets the rightmost column and bottom row of a diagram.

Theorem 2.7. *Let \mathcal{C} be a k -linear abelian category with stability function Z and framing Φ . For each $\phi \in (0, 1)$ the map $F_\bullet : \mathcal{S}_\bullet^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}})$ is unital relative 2-Segal.*

Proof. The proof that $\mathcal{S}_\bullet^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ is 1-Segal reduces to the 2-Segal property of $\mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}})$, so we omit it. To verify the second of the relative 2-Segal conditions we need to show that the functor

$$\Psi_n : \mathcal{S}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{S}_n(\mathcal{C}_\phi^{Z\text{-ss}}) \times_{\mathcal{S}_{\{0,n\}}(\mathcal{C}_\phi^{Z\text{-ss}})}^{(2)} \mathcal{S}_{\{0,n\}}^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$$

is an equivalence. Here $- \times^{(2)} -$ denotes the 2-pullback of groupoids.

Let $\mathcal{F}_n(\mathcal{C}_\phi^{Z\text{-ss}})$ be the groupoid of n -flags in $\mathcal{C}_\phi^{Z\text{-ss}}$, that is, diagrams in $\mathcal{C}_\phi^{Z\text{-ss}}$ of the form

$$0 \rightrightarrows A_1 \rightrightarrows \dots \rightrightarrows A_{n-1} \rightrightarrows A_n.$$

The forgetful functor $\mu_n : \mathcal{S}_n(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{F}_n(\mathcal{C}_\phi^{Z\text{-ss}})$ is an equivalence of groupoids [46, §1.3]. Similarly, let $\mathcal{F}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ be the groupoid of n -flags in a stable framed object. Explicitly, an object of $\mathcal{F}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ is a diagram of the form

$$0 \rightrightarrows A_1 \rightrightarrows \dots \rightrightarrows A_{n-1} \rightrightarrows A_n \rightrightarrows (M, s) \quad (10)$$

with $A_1, \dots, A_n, M \in \mathcal{C}_\phi^{Z\text{-ss}}$ and (M, s) stable framed. We claim that the forgetful functor $\nu_n : \mathcal{S}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{F}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}})$ is an equivalence. A quasi-inverse η_n of ν_n can be constructed as follows. Given a flag (10) set $A_{\{0,k\}} = A_k$, $1 \leq k \leq n$, and $(M_0, s_0) = (M, s)$. Define $A_{\{1,k\}}$, $1 \leq k \leq n$, and M_1 as the pushouts

$$\begin{array}{ccc} A_{\{0,1\}} & \rightrightarrows & A_{\{0,k\}} \\ \downarrow & & \downarrow \\ 0 & \dashrightarrow & A_{\{1,k\}} \end{array} \quad \begin{array}{ccc} A_{\{0,1\}} & \rightrightarrows & M_0 \\ \downarrow & & \downarrow \\ 0 & \dashrightarrow & M_1. \end{array}$$

Since $\mathcal{C}_\phi^{Z\text{-ss}}$ is abelian, it is automatic that each $A_{\{1,k\}}$ and M_1 are Z -semistable of phase ϕ . Let s_1 be the image of s_0 under the morphism $\Phi(M_0) \rightarrow \Phi(M_1)$. We need to show that (M_1, s_1) is stable framed. Note that s_1 is non-zero. Indeed, if $s_1 = 0$, then $s_0 \in \Phi(A_{\{0,1\}}) \subset \Phi(M_0)$ and $\phi(A_{\{0,1\}}) = \phi(M_0)$, contradicting the assumed framed stability of (M_0, s_0) . Let then $B_1 \subset M_1$ be a proper subobject with $s_1 \in \Phi(B_1) \subset \Phi(M_1)$. There exists a unique object $B_0 \in \mathcal{C}$ which fits into the exact commutative diagram

$$\begin{array}{ccccc} A_{\{0,1\}} & \rightrightarrows & B_0 & \longrightarrow & B_1 \\ \text{id} \downarrow & & \downarrow & & \downarrow \\ A_{\{0,1\}} & \rightrightarrows & M_0 & \longrightarrow & M_1 \\ & & \downarrow & & \downarrow \\ & & M_0/B_0 & \xrightarrow{\sim} & M_1/B_1. \end{array}$$

Since $s_1 \in \Phi(B_1)$, using the left exactness of Φ we see that s_1 lies in the kernel of $\Phi(M_1) \rightarrow \Phi(M_1/B_1)$. It follows that s_0 is in the kernel of $\Phi(M_0) \rightarrow \Phi(M_0/B_0)$. Hence $s_0 \in \Phi(B_0) \subset \Phi(M_0)$. Since (M_0, s_0) is stable framed we have $\phi(B_0) < \phi(M_0)$. The equalities $\phi(A_{\{0,1\}}) = \phi(M_0) = \phi(M_1)$ combined with the standard See-Saw properties of stability functions then give $\phi(M_1) > \phi(B_1)$, as desired. This procedure defines the top two rows of $\eta_n(A_{\{0,\bullet\}} \rightarrow (M_0, s_0))$ and can be iterated to define the remaining $n - 2$ rows. This defines the desired quasi-inverse.

We can now prove the theorem. Consider the following commutative diagram:

$$\begin{array}{ccc} \mathcal{S}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) & \xrightarrow{\Psi_n} & \mathcal{S}_n(\mathcal{C}_\phi^{Z\text{-ss}}) \times_{\mathcal{S}_{\{0,n\}}(\mathcal{C}_\phi^{Z\text{-ss}})}^{(2)} \mathcal{S}_{\{0,n\}}^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \\ \nu_n \downarrow & & \downarrow \mu_n \times \nu_{\{0,n\}} \\ \mathcal{F}_n^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) & \xrightarrow{\tilde{\Psi}_n} & \mathcal{F}_n(\mathcal{C}_\phi^{Z\text{-ss}}) \times_{\mathcal{F}_{\{0,n\}}(\mathcal{C}_\phi^{Z\text{-ss}})}^{(2)} \mathcal{F}_{\{0,n\}}^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}). \end{array}$$

Noting that the groupoids $\mathcal{S}_1(\mathcal{C}_\phi^{Z\text{-ss}})$ and $\mathcal{F}_1(\mathcal{C}_\phi^{Z\text{-ss}})$ are canonically equivalent, the discussion above shows that the vertical functors are equivalences. That Ψ_n is an equivalence therefore reduces to the statement that $\tilde{\Psi}_n$ is an equivalence, which is obvious. We omit the verification of the relative unital condition. \square

Remark. The notion of quotient datum on an abelian category \mathcal{A} is introduced in [45, §5.5], where it is shown to define a relative 2-Segal groupoid over $\mathcal{S}_\bullet(\mathcal{A})$. From this point of view, Theorem 2.7 can be rephrased as the statement that a stability function Z and framing Φ on an abelian category \mathcal{C} defines a (mild generalization of a) quotient datum on the Artin mapping cylinder \mathcal{C}_Φ .

2.5. Real pseudoholomorphic polygons. In this section we give a relative variant of the pseudoholomorphic polygon construction of [7, §3.8].

Recall that an almost complex structure on a smooth manifold M is an endomorphism $J : TM \rightarrow TM$ of the tangent bundle which satisfies $J \circ J = -\mathbf{1}_{TM}$. A continuous map $u : (\Sigma, j) \rightarrow (M, J)$ of almost complex manifolds is called pseudoholomorphic if it is smooth and satisfies the equation

$$du + J \circ du \circ j = 0.$$

Let \mathbb{H} be the Lobachevsky plane, realized as the open unit disk in \mathbb{C} with centre the origin and metric

$$ds^2 = \frac{dz \wedge d\bar{z}}{(1 - |z|^2)^2}.$$

The ideal boundary $\partial\mathbb{H}$ is the unit circle in \mathbb{C} . Denote by $j_{\mathbb{H}}$ the complex structure on \mathbb{H} . The group $\text{SL}_2(\mathbb{R})$ acts on \mathbb{H} by orientation preserving isometries.

We require some basic definitions from Teichmüller theory [29]. Given $b, b' \in \partial\mathbb{H}$, let (b, b') be the oriented geodesic with limiting points b at infinite negative time and b' at infinite positive time. For each $n \geq 0$ let $P(b_0, \dots, b_n)$ be the ideal $(n+1)$ -gon in \mathbb{H} with vertices $b_0, \dots, b_n \in \partial\mathbb{H}$ numbered compatibly with the canonical orientation of $\partial\mathbb{H}$. A decoration of $P(b_0, \dots, b_n)$ is the data of a horocycle $\xi_i \in \text{Hor}_{b_i}$ with hyperbolic centre b_i for each $i = 0, \dots, n$. The group $\text{SL}_2(\mathbb{R}) \times \mathbb{R}$ acts on the set of decorated ideal $(n+1)$ -gons by the formula

$$(g, a) \cdot (P(b_0, \dots, b_n), \xi_0, \dots, \xi_n) = (P(g \cdot b_0, \dots, g \cdot b_n), g \cdot \xi_0 + a, \dots, g \cdot \xi_n + a)$$

where $g \in \text{SL}_2(\mathbb{R})$ and $a \in \mathbb{R}$ and we have used the canonical identification of \mathbb{R} -torsors $g : \text{Hor}_{b_i} \rightarrow \text{Hor}_{g \cdot b_i}$. We will often omit the decoration from the notation if it will not cause confusion.

Given a decorated ideal $(n+1)$ -gon, for each $0 \leq i < j \leq n$ let $m(\xi_i, \xi_j)$ be the midpoint of the unique geodesic connecting the points $\xi_i \cap (b_i, b_j)$ and $\xi_j \cap (b_i, b_j)$. This trivializes the \mathbb{R} -torsor (b_i, b_j) via

$$\mathbb{R} \rightarrow (b_i, b_j), \quad 0 \mapsto m(\xi_i, \xi_j). \quad (11)$$

This trivialization is invariant under the action of $\mathbb{R} \subset \mathrm{SL}_2(\mathbb{R}) \times \mathbb{R}$.

Following [7, §3.8] we construct from an almost complex manifold (M, J) a semi-simplicial set $\tilde{\mathbb{T}}_\bullet(M)$. Let $\tilde{\mathbb{T}}_0(M) = M$ and, for each $n \geq 1$, let $\tilde{\mathbb{T}}_n(M)$ be the set of equivalence classes of pairs consisting of a decorated ideal $(n+1)$ -gon P together with a continuous map $u : (P, j_{\mathbb{H}}) \rightarrow (M, J)$ which is pseudoholomorphic on the two dimensional interior of P . The equivalence relation is generated by the action of $\mathrm{SL}_2(\mathbb{R}) \times \mathbb{R}$ on decorated ideal polygons. Using the trivialization (11), we can identify $\tilde{\mathbb{T}}_1(M)$ with the set of continuous maps $[-\infty, \infty] \rightarrow M$. The face map $\partial_i : \tilde{\mathbb{T}}_n(M) \rightarrow \tilde{\mathbb{T}}_{n-1}(M)$ omits the ideal boundary point b_i and forms the resulting decorated n -gon together with the restricted morphism to M . It is proved in [7, Theorem 3.8.6] that $\tilde{\mathbb{T}}_\bullet(M)$ is a 2-Segal semi-simplicial set.

Consider now the relative setting. Recall that a real structure on (M, J) is a smooth involution $\tau : M \rightarrow M$ which satisfies

$$d\tau - J \circ d\tau \circ J = 0.$$

A map $u : (\Sigma, j, \sigma) \rightarrow (M, J, \tau)$ of almost complex manifolds with real structures is called real pseudoholomorphic if it is pseudoholomorphic and satisfies $u \circ \sigma = \tau \circ u$.

The real structure $\sigma : z \mapsto -\bar{z}$ on \mathbb{C} induces a real structure on \mathbb{H} , again denoted by σ . The subgroup $\mathrm{SL}_2(\mathbb{R})^\sigma \leq \mathrm{SL}_2(\mathbb{R})$ which commutes with σ is isomorphic to $\mathbb{R}^\times \rtimes \mathbb{Z}_2$, the generator of \mathbb{Z}_2 being rotation of \mathbb{H} through an angle π . Define a real ideal boundary point of \mathbb{H} to be either a point of the real locus $\partial\mathbb{H}^\sigma = \{\sqrt{-1}, -\sqrt{-1}\}$ or a pair $\{b, \sigma(b)\}$ of distinct σ -conjugate ideal boundary points.

A real ideal $(n+1)$ -gon $Q = Q(b_0, \dots, b_n, \dots)$ is an ideal polygon which has exactly $n+1$ real ideal boundary points, labelled so that only b_0 and b_n may lie in $\partial\mathbb{H}^\sigma$. It follows that Q is of one of the following four types:

- (i) $Q(b_0, \dots, b_n, \sigma(b_n), \sigma(b_{n-1}), \dots, \sigma(b_1), \sigma(b_0))$
- (ii) $Q(b_0, \dots, b_n, \sigma(b_n), \sigma(b_{n-1}), \dots, \sigma(b_1))$
- (iii) $Q(b_0, \dots, b_n, \sigma(b_{n-1}), \dots, \sigma(b_1), \sigma(b_0))$
- (iv) $Q(b_0, \dots, b_n, \sigma(b_{n-1}), \dots, \sigma(b_1))$.

Then Q has exactly zero, one, one and two ideal vertices in $\partial\mathbb{H}^\sigma$ in cases (i)-(iv), respectively. A decoration of Q is a decoration of its underlying ideal polygon for which the horocycle at a vertex b is equal to that at $\sigma(b)$ under the canonical identification $\mathrm{Hor}_b \simeq \mathrm{Hor}_{\sigma(b)}$. The group $\mathrm{SL}_2(\mathbb{R})^\sigma \times \mathbb{R}$ acts on the set of real decorated ideal polygons.

With the above notation in place, we define a semi-simplicial set $\tilde{\mathbb{T}}_\bullet^\tau(M)$ as follows. For each $n \geq 0$ let $\tilde{\mathbb{T}}_n^\tau(M)$ be the set of equivalence classes of pairs (Q, v) consisting of a real decorated ideal $(n+1)$ -gon Q together with a real continuous map $v : (Q, j_{\mathbb{H}}, \sigma) \rightarrow (M, J, \tau)$ which is pseudoholomorphic on the two dimensional interior of Q . The equivalence relation is generated by the action of $\mathrm{SL}_2(\mathbb{R})^\sigma \times \mathbb{R}$ on real decorated ideal polygons. In particular, we have

$$\tilde{\mathbb{T}}_0^\tau(M) \simeq M^\tau \sqcup C^0([-\infty, \infty], M)^\tau \quad (12)$$

where $C^0([-\infty, \infty], M)^\tau$ denotes the set of real continuous maps $[-\infty, \infty] \rightarrow M$ with the compactified real line $\overline{\mathbb{R}} = [-\infty, \infty]$ given the \mathbb{Z}_2 -action $x \mapsto -x$. Define face maps $\partial_i : \tilde{\mathbb{T}}_n^\tau(M) \rightarrow \tilde{\mathbb{T}}_{n-1}^\tau(M)$ by omitting the real ideal boundary point $\{b_i, \sigma(b_i)\}$. Then $\tilde{\mathbb{T}}_\bullet^\tau(M)$ forms a semi-simplicial set.




FIGURE 1. The map $\Xi_2 : \tilde{\mathbb{T}}_2^\tau(M) \rightarrow \tilde{\mathbb{T}}_{\{0,1\}}^\tau(M) \times_{\tilde{\mathbb{T}}_{\{1\}}^\tau(M)} \tilde{\mathbb{T}}_{\{1,2\}}^\tau(M)$, omitting the data of the maps to M .

Given $(Q, v) \in \tilde{\mathbb{T}}_n^\tau(M)$ write $Q = Q(b_0, \dots, b_n, \dots)$ as above and let $\tilde{Q} = \tilde{Q}(b_0, \dots, b_n)$ be the decorated ideal $(n+1)$ -gon obtained from Q by omitting the indicated vertices and their decorations. This defines a semi-simplicial morphism

$$F_\bullet : \tilde{\mathbb{T}}_\bullet^\tau(M) \rightarrow \tilde{\mathbb{T}}_\bullet(M), \quad (Q, v) \mapsto (\tilde{Q}, v|_{\tilde{Q}}).$$

We can now state the main result of this section.

Theorem 2.8. *Let (M, J, τ) be an almost complex manifold with real structure. The morphism $F_\bullet : \tilde{\mathbb{T}}_\bullet^\tau(M) \rightarrow \tilde{\mathbb{T}}_\bullet(M)$ is a relative 2-Segal semi-simplicial set.*

Proof. For each $n \geq 2$ and $0 < i < n$ we construct an inverse of the 1-Segal map

$$\Xi_n : \tilde{\mathbb{T}}_n^\tau(M) \rightarrow \tilde{\mathbb{T}}_{\{0, \dots, i\}}^\tau(M) \times_{\tilde{\mathbb{T}}_{\{i\}}^\tau(M)} \tilde{\mathbb{T}}_{\{i, \dots, n\}}^\tau(M).$$

Let $(Q', v') \in \tilde{\mathbb{T}}_{\{0, \dots, i\}}^\tau(M)$ and $(Q'', v'') \in \tilde{\mathbb{T}}_{\{i, \dots, n\}}^\tau(M)$ with equal image in $\tilde{\mathbb{T}}_{\{i\}}^\tau(M)$. Since $0 < i < n$ these images lie in the component $C^0([-\infty, \infty], M)^\tau$ in the decomposition (12). There exists a unique $g \in \mathrm{SL}_2(\mathbb{R})^\sigma$ with the property that $g \cdot (b'_i, \sigma(b'_i)) = (b''_i, \sigma(b''_i))$ and the restrictions

$$g \cdot v', v'' : [-\infty, \infty] \rightarrow M$$

are equal. Applying Morera's theorem, we see that $g \cdot (Q', v')$ and (Q'', v'') can be glued in a unique way so as to obtain a continuous map

$$g \cdot v' \cup v'' : g \cdot Q' \cup_{(b'_i, \sigma(b'_i))} Q'' \rightarrow M$$

which is pseudoholomorphic on the two dimensional interior. Since both Q' and Q'' (resp. v' and v'') are real, so too is $g \cdot Q' \cup_{(b'_i, \sigma(b'_i))} Q''$ (resp. $g \cdot v' \cup v''$). See Figure 1. This defines the required inverse of Ξ_n , showing that $\tilde{\mathbb{T}}_\bullet^\tau(M)$ is 1-Segal.

To verify the second of the relative 2-Segal conditions, consider the map

$$\Psi_n : \tilde{\mathbb{T}}_n^\tau(M) \rightarrow \tilde{\mathbb{T}}_n(M) \times_{\tilde{\mathbb{T}}_{\{0, n\}}^\tau(M)} \tilde{\mathbb{T}}_{\{0, n\}}^\tau(M), \quad n \geq 1.$$

Let $(P', u') \in \tilde{\mathbb{T}}_n(M)$ and $(Q'', v'') \in \tilde{\mathbb{T}}_{\{0, n\}}^\tau(M)$ with equal image in $\tilde{\mathbb{T}}_{\{0, n\}}^\tau(M)$. Choose $g \in \mathrm{SL}_2(\mathbb{R})$ so that $g \cdot (b'_0, b'_n) = (b''_0, b''_n)$. As above, the reality condition on (Q'', v'') implies that, up to the action of $\mathrm{SL}_2(\mathbb{R})^\sigma$, the triple

$$\{g \cdot (P', u'), (Q'', v''), g \cdot (\sigma(P'), \tau \circ u' \circ \sigma^{-1})\}$$

can be glued in a unique way so as to obtain an element of $\tilde{\mathbb{T}}_n^\tau(M)$. See Figure 2. This defines the desired inverse of Ψ_n . \square




FIGURE 2. The map $\Psi_2 : \widetilde{\mathbb{T}}_2^\tau(M) \rightarrow \widetilde{\mathbb{T}}_2(M) \times_{\widetilde{\mathbb{T}}_{\{0,2\}}(M)} \widetilde{\mathbb{T}}_{\{0,2\}}^\tau(M)$, omitting the data of the maps to M .

3. RELATIVE HIGHER SEGAL SPACES FROM CATEGORIES WITH DUALITIES

3.1. Categories with duality. We recall some basics about categories with duality. For further details the reader is referred to [35].

Definition. A category with (strong) duality is a triple (\mathcal{C}, P, Θ) consisting of a category \mathcal{C} , a functor $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$ and a natural isomorphism $\Theta : \mathbf{1}_{\mathcal{C}} \Rightarrow P \circ P^{\text{op}}$ such that $P(\Theta_U) \circ \Theta_{P(U)} = \mathbf{1}_{P(U)}$ for all objects $U \in \mathcal{C}$.

When there is no risk of confusion we will omit (P, Θ) from the notation and simply refer to \mathcal{C} as a category with duality.

A form functor $(T, \varphi) : (\mathcal{C}, P, \Theta) \rightarrow (\mathcal{D}, Q, \Xi)$ between categories with duality consists of a functor $T : \mathcal{C} \rightarrow \mathcal{D}$ and a natural transformation $\varphi : T \circ P \Rightarrow Q \circ T^{\text{op}}$ for which the diagram

$$\begin{array}{ccc} T(U) & \xrightarrow{\Xi_{T(U)}} & Q^2 T(U) \\ T(\Theta_U) \downarrow & & \downarrow Q(\varphi_U) \\ TP^2(U) & \xrightarrow{\varphi_{P(U)}} & QTP(U) \end{array}$$

commutes for all objects $U \in \mathcal{C}$. In particular, let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a duality preserving functor, that is, a functor for which $T \circ P = Q \circ T^{\text{op}}$ and $T\Theta = \Xi T$, where $T\Theta$ (resp. ΞT) is the left (resp. right) whiskering of T with Θ (resp. Ξ). Then $(T, \mathbf{1}_T)$ is a form functor. Write CatD for the category of small categories with duality with form functors as morphisms.

A (nonsingular) symmetric form in \mathcal{C} is a pair (N, ψ_N) , often just denoted by N , consisting of an object $N \in \mathcal{C}$ and an isomorphism $\psi_N : N \rightarrow P(N)$ which satisfies $P(\psi_N) \circ \Theta_N = \psi_N$. An isometry of symmetric forms $\phi : (M, \psi_M) \rightarrow (N, \psi_N)$ is an isomorphism $\phi : M \rightarrow N$ which satisfies $P(\phi) \circ \psi_N \circ \phi = \psi_M$. The groupoid of symmetric forms and their isometries is denoted by \mathcal{C}_h and is called the Hermitian groupoid of \mathcal{C} .

For each integer $n \geq 0$ let \mathbf{n} be the ordered set $\{0 < \dots < n < n' < \dots < 0'\}$. There is a unique isomorphism $\mathbf{n} \simeq [2n + 1]$ in Δ . The morphisms $[n] \rightarrow \mathbf{n}$, $i \mapsto i$, define the subdivision functor $\text{sd} : \Delta \rightarrow \Delta$. The edgewise subdivision of a simplicial object X_\bullet of a category \mathcal{C} is then defined to be the composition

$$X_\bullet^e : \Delta^{\text{op}} \xrightarrow{\text{sd}^{\text{op}}} \Delta^{\text{op}} \xrightarrow{X_\bullet} \mathcal{C}.$$

Explicitly, $X_n^e = X_{\mathbf{n}} = X_{2n+1}$ with face maps

$$\partial_i^e : X_n^e = X_{2n+1} \xrightarrow{\partial_i \circ \partial_{2n+1-i}} X_{2n-1} = X_{\mathbf{n}-1}^e,$$

the degeneracy maps admitting a similar description. The morphisms $[n] \rightarrow \mathbf{n}$ define a simplicial morphism $X_\bullet^e \rightarrow X_\bullet$.

In the sections below we will repeatedly apply the following result (*cf.* [15, §1.5]).

Lemma 3.1. *Let X_\bullet be a simplicial object of Cat . For each $n \geq 0$ let (P_n, Θ_n) , $n \geq 0$ be duality structures on X_n . Suppose that*

$$P_{n-1} \circ \partial_i^{\text{op}} = \partial_{n-i} \circ P_n, \quad P_{n+1} \circ s_i^{\text{op}} = s_{n-i} \circ P_n$$

and

$$\partial_i \Theta_n = \Theta_{n-1} \partial_i, \quad s_i \Theta_n = \Theta_{n+1} s_i$$

for all $n \geq 1$ and $0 \leq i \leq n$. Then $(X_\bullet^e, P_\bullet^e, \Theta_\bullet^e)$ is a simplicial object of CatD .

Proof. It is clear that $(X_n^e, P_n^e, \Theta_n^e) = (X_{2n+1}, P_{2n+1}, \Theta_{2n+1})$ is a category with duality. The equality $\partial_i \Theta_n = \Theta_{n-1} \partial_i$ together with the calculation

$$\begin{aligned} \partial_i^e \circ P_n^e &= \partial_i \circ \partial_{2n+1-i} \circ P_{2n+1} \\ &= \partial_i \circ P_{2n} \circ \partial_i^{\text{op}} \\ &= P_{2n-1} \circ \partial_{2n-i}^{\text{op}} \circ \partial_i^{\text{op}} \\ &= P_{2n-1} \circ \partial_i^{\text{op}} \circ \partial_{2n-1-i}^{\text{op}} = P_{n-1}^e \circ (\partial_i^e)^{\text{op}} \end{aligned}$$

shows that the face map $(\partial_i^e, \mathbf{1}_{\partial_i^e}) : X_n^e \rightarrow X_{n-1}^e$ is a form functor. A similar calculation shows that $(s_i^e, \mathbf{1}_{s_i^e}) : X_n^e \rightarrow X_{n+1}^e$ is a form functor. \square

3.2. Unoriented categorified nerves. As a warm-up for Section 3.3 we adapt the categorified nerve construction to the relative setting.

Let \mathcal{C} be a small category. Its categorified nerve $\mathcal{N}_\bullet(\mathcal{C})$ (see Section 1.2) is a 1-Segal simplicial groupoid. Suppose that (P, Θ) is a duality structure on \mathcal{C} . Then the groupoids $\mathcal{N}_n(\mathcal{C})$, $n \geq 0$, inherit duality structures which are compatible in the sense of Lemma 3.1. The edgewise subdivision $\mathcal{N}_\bullet^e(\mathcal{C})$ is therefore a simplicial groupoid with duality.

Definition. *The Hermitian groupoid $\mathcal{N}_\bullet^e(\mathcal{C})_h$, denoted by $\mathcal{U}_\bullet(\mathcal{C})$, is called the categorified unoriented nerve of (\mathcal{C}, P, Θ) .*

Slightly abusively, write $(x_\bullet, \psi_\bullet) \in \mathcal{U}_n(\mathcal{C})$ for the object consisting of the diagram

$$x_0 \rightarrow \cdots \rightarrow x_n \rightarrow x_{n'} \rightarrow \cdots \rightarrow x_{0'}$$

in \mathcal{C} together with isomorphisms $\psi_i : x_i \rightarrow P(x_{i'})$, $0 \leq i \leq 0'$, which satisfy $P(\psi_i) \circ \Theta_{x_{i'}} = \psi_{i'}$ and make the obvious squares commute. Here we use the convention that $i'' = i$. For later use, note that an object $(x_0 \xrightarrow{f} x_{0'}, \{\psi_0, \psi_{0'}\})$ of $\mathcal{U}_0(\mathcal{C})$ is determined by the pair (f, ψ_0) and that $P(f) \circ \psi_0$ defines a possibly singular symmetric form on x_0 . In this way, $\mathcal{U}_0(\mathcal{C})$ can be interpreted as the groupoid of presented symmetric forms in \mathcal{C} , that is, possibly singular symmetric forms $\psi : x \rightarrow P(x)$ with the additional data of a factorization $\psi = P(f_x) \circ \psi_x$, where $\psi_x : x \rightarrow P(x')$ an isomorphism.

There is a canonical forgetful morphism $F_\bullet : \mathcal{U}_\bullet(\mathcal{C}) \rightarrow \mathcal{N}_\bullet(\mathcal{C})$ given by

$$\mathcal{U}_n(\mathcal{C}) \ni (x_\bullet, \psi_\bullet) \mapsto (x_0 \rightarrow \cdots \rightarrow x_n) \in \mathcal{N}_n(\mathcal{C}).$$

Proposition 3.2. *For any small category with duality (\mathcal{C}, P, Θ) , the morphism $F_\bullet : \mathcal{U}_\bullet(\mathcal{C}) \rightarrow \mathcal{N}_\bullet(\mathcal{C})$ is unital right relative 1-Segal.*

Proof. Fix $0 < i < n$. An object of the 2-pullback $\mathcal{N}_{\{0, \dots, i\}}(\mathcal{C}) \times_{\mathcal{N}_{\{i\}}(\mathcal{C})}^{(2)} \mathcal{U}_{\{i, \dots, n\}}(\mathcal{C})$ is a triple

$$(x_0 \rightarrow \cdots \rightarrow x_i, (x'_i \rightarrow \cdots \rightarrow x'_n \rightarrow x'_{n'} \rightarrow \cdots \rightarrow x'_{i'}, \psi_\bullet); \alpha) \quad (13)$$

with $\alpha : x_i \rightarrow x'_i$ an isomorphism. A morphism from the object (13) to a second object, say

$$(y_0 \rightarrow \cdots \rightarrow y_i, (y'_i \rightarrow \cdots \rightarrow y'_n \rightarrow y'_{n'} \rightarrow \cdots \rightarrow y'_{i'}, \mu_\bullet); \beta),$$

is a tuple $((p_0, \dots, p_i), (q_i, \dots, q_n, q_{n'}, \dots, q_{i'}))$ where $p_j : x_j \rightarrow y_j$, $1 \leq j \leq i$, and $q_j : x'_j \rightarrow y'_j$, $i \leq j \leq i'$, are isomorphisms which make the obvious nerve diagrams commute, the q_j respect the symmetric forms ψ_\bullet and μ_\bullet and the compatibility condition

$$q_i \circ \alpha = \beta \circ p_i \quad (14)$$

holds.

We need to prove that the functor

$$\begin{aligned} \Psi_n : \mathcal{U}_n(\mathcal{C}) &\rightarrow \mathcal{N}_{\{0, \dots, i\}}(\mathcal{C}) \times_{\mathcal{N}_{\{i\}}(\mathcal{C})}^{(2)} \mathcal{U}_{\{i, \dots, n\}}(\mathcal{C}) \\ (x_\bullet, \psi_\bullet) &\mapsto (x_0 \rightarrow \dots \rightarrow x_i, (x_i \rightarrow \dots \rightarrow x_n \rightarrow x_{n'} \rightarrow \dots \rightarrow x_{i'}, \psi_\bullet); \mathbf{1}_{x_i}) \end{aligned}$$

is an equivalence. Consider a morphism $(p_\bullet, q_\bullet) : \Psi_n(x_\bullet, \psi_\bullet) \rightarrow \Psi_n(y_\bullet, \mu_\bullet)$. Since α and β are the identity maps in this case, the compatibility condition (14) reduces to $q_i = p_i$. Any morphism $\Psi_n(x_\bullet, \psi_\bullet) \rightarrow \Psi_n(y_\bullet, \mu_\bullet)$ is therefore the image of a unique isometry $(x_\bullet, \psi_\bullet) \rightarrow (y_\bullet, \mu_\bullet)$, since for an isometry the map $x_j \rightarrow y_j$ uniquely determines the map $x_{j'} \rightarrow y_{j'}$. It follows that Ψ_n is fully faithful. To show that Ψ_n is essentially surjective suppose that we are given an object of the form (13). Consider the object of $\mathcal{U}_n(\mathcal{C})$ whose underlying diagram is the top row of

$$\begin{array}{ccccccccccc} x_0 & \rightarrow & \dots & \rightarrow & x_{i-1} & \rightarrow & x'_i & \rightarrow & \dots & \rightarrow & x'_n & \rightarrow & x'_{n'} & \rightarrow & \dots & \rightarrow & x'_{i'} & \rightarrow & P(x_{i-1}) & \rightarrow & \dots & \rightarrow & P(x_0) \\ & & & & \downarrow & \nearrow & \alpha & & & & & & \downarrow & \nearrow & \psi_{i'} & & & & \downarrow & \nearrow & & & & & \\ & & & & x_i & & & & & & & & & & P(x'_i) & & & & & & & & & & \end{array}$$

and whose symmetric form is $(\{\Theta_\bullet\}_{[0, i]}, \{\psi_\bullet\}_{[i, i']}, \{\mathbf{1}_\bullet\}_{[i', 0']})$, the subscripts indicating the index intervals to which each map applies. In the above diagram the triangles define the corresponding horizontal morphisms and the unlabelled maps are the canonical ones. The image of this object under Ψ_n is

$$(x_0 \rightarrow \dots \rightarrow x_{i-1} \rightarrow x'_i, (x'_i \rightarrow \dots \rightarrow x'_n \rightarrow x'_{n'} \rightarrow \dots \rightarrow x'_{i'}, \psi_\bullet); \mathbf{1}_{x'_i})$$

which is isomorphic via $((\mathbf{1}_{x_0}, \dots, \mathbf{1}_{x_{i-1}}, \alpha^{-1}), (\mathbf{1}_{x'_i}, \dots, \mathbf{1}_{x'_{i'}}))$ to the object (13). \square

The constructions of this section also define a simplicial set $U_\bullet(\mathcal{C}) = N_\bullet^e(\mathcal{C})_h$ and a morphism $F_\bullet : U_\bullet(\mathcal{C}) \rightarrow N_\bullet(\mathcal{C})$. While this morphism is neither left nor right relative 1-Segal, it is relevant to an interpretation of Proposition 3.2 in terms of an $(\infty, 1)$ -presheaf on \mathcal{C} , as suggested by Section 2.1. We explain this in the remainder of the section.

First, recall the Joyal-Tierney Quillen equivalence (left adjoint on the left)

$$p_1^* : (\mathbb{S}, \mathcal{J}) \rightleftarrows (\mathbb{S}_\Delta, \mathcal{S}\text{eg}_1) : i_1^*, \quad (15)$$

where \mathcal{J} is the quasi-category model structure [19, §4]. The functor i_1^* is such that the n -simplices of $i_1^* X_\bullet$ are the 0-simplices of X_n . Moreover, for each complete 1-Segal space $X_\bullet \in \mathbb{S}_\Delta$ the equivalence (15) lifts to a Quillen equivalence

$$p_1^* : \left(\mathbb{S}_{/i_1^* X_\bullet}, \mathcal{J}_{/i_1^* X_\bullet}^{\text{rfib}} \right) \rightleftarrows \left((\mathbb{S}_\Delta)_{/X_\bullet}, (\mathcal{S}\text{eg}_1)_{/X_\bullet}^{\text{rfib}} \right) : i_1^*,$$

where rfib denotes the induced right fibration model structure [3, Theorem 1.22].

Returning to unoriented nerves, let $\mathbb{N}_\bullet(\mathcal{C}) \in \mathbb{S}_\Delta$ be the complete 1-Segal space having $N_\bullet(\mathcal{N}_n(\mathcal{C}))$ as its n -simplices [32, §3.5]. In a similar way we define $\mathbb{U}_\bullet(\mathcal{C}) \in \mathbb{S}_\Delta$. Proposition 3.2 implies that the induced morphism $\mathbb{F}_\bullet : \mathbb{U}_\bullet(\mathcal{C}) \rightarrow \mathbb{N}_\bullet(\mathcal{C})$ is right relative 1-Segal. Noting that $i_1^* \mathbb{N}_\bullet(\mathcal{C}) = N_\bullet(\mathcal{C})$ and $i_1^* \mathbb{U}_\bullet(\mathcal{C}) = U_\bullet(\mathcal{C})$, we obtain a right (Kan) fibration $i_1^* \mathbb{F}_\bullet : U_\bullet(\mathcal{C}) \rightarrow N_\bullet(\mathcal{C})$. We can therefore apply [14, Proposition A] to construct from $i_1^* \mathbb{F}_\bullet$ the desired $(\infty, 1)$ -presheaf on \mathcal{C} . However, in our setting there is a simpler approach. As can be verified directly, the simplicial

set $U_\bullet(\mathcal{C})$ is 1-Segal and hence isomorphic to the nerve of the category \mathcal{S} whose objects are presented symmetric forms in \mathcal{C} and whose morphisms are

$$\mathrm{Hom}_{\mathcal{S}}((x; f_x, \psi_x), (y; f_y, \psi_y)) = \{g \in \mathrm{Hom}_{\mathcal{C}}(x, y) \mid \tilde{g} \circ f_y \circ g = f_x\},$$

where \tilde{g} is defined by $P(\tilde{g}) = \psi_y \circ g \circ \psi_x^{-1}$. As $i_1^* \mathbb{F}_\bullet$ is a right fibration, the induced functor $F : \mathcal{S} \rightarrow \mathcal{C}$ is fibred in groupoids and thus determined by the presheaf

$$h^* F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Grpd}, \quad x \mapsto \mathrm{Hom}_{\mathrm{Cat}/\mathcal{C}}(\mathcal{C}/x, \mathcal{S}).$$

More explicitly, the 2-Yoneda lemma defines an equivalence between $h^* F(x)$ and the fibre groupoid \mathcal{S}_x , the category with presented symmetric forms on x as objects and with a unique map $(x; f_x, \psi_x) \rightarrow (x; \bar{f}_x, \bar{\psi}_x)$ whenever

$$\bar{f}_x = \Theta_{\mathrm{target}(\bar{\psi}_x)}^{-1} \circ P(\psi_x \circ \bar{\psi}_x^{-1}) \circ \Theta_{\mathrm{target}(\psi_x)} \circ f_x.$$

In words, equivalences in \mathcal{S}_x correspond to isomorphic changes in the target of ψ_x . After embedding Grpd into ∞ -groupoids via the nerve, $h^* F$ agrees with the $(\infty, 1)$ -presheaf produced by [14].

3.3. Unoriented categorified twisted cyclic nerves. Let X be a topological space with a self-homeomorphism $T : X \rightarrow X$. The T -twisted loop space of X is then defined to be

$$L^T X = \{\gamma \in C^0(\mathbb{R}, X) \mid \gamma(t+1) = T(\gamma(t))\}.$$

The case of the identity map $T = \mathbf{1}_X$ recovers the ordinary loop space LX . Suppose that we are also given a continuous involution $p : X \rightarrow X$ which satisfies

$$p = T \circ p \circ T.$$

This, together with the orientation reversing involution $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, $t \mapsto 1 - t$, induces an involution

$$p_L : L^T X \rightarrow L^T X, \quad \gamma \mapsto p \circ \gamma \circ \sigma^{-1},$$

the fixed point set of which is naturally interpreted as the unoriented loop space of the stack $[X/\langle T \rangle]$. The goal of this section is to construct a relative 2-Segal simplicial space which is a categorical analogue of the unoriented loop space.

Let \mathcal{C} be a small category with an endofunctor $T : \mathcal{C} \rightarrow \mathcal{C}$. The categorified T -twisted cyclic nerve of \mathcal{C} (see [7, §3.3], [5]) is the simplicial groupoid $\mathcal{NC}_\bullet^T(\mathcal{C})$ which assigns to $[n] \in \Delta$ the groupoid of all diagrams in \mathcal{C} of the form

$$x_\bullet = \{x_0 \xrightarrow{f_0} x_1 \xrightarrow{f_1} \dots \xrightarrow{f_{n-2}} x_{n-1} \xrightarrow{f_{n-1}} x_n \xrightarrow{f_n} T(x_0)\}. \quad (16)$$

A morphism $x_\bullet \rightarrow y_\bullet$ in $\mathcal{NC}_\bullet^T(\mathcal{C})$ is a collection of isomorphisms $x_i \rightarrow y_i$, $0 \leq i \leq n$, which make the obvious diagrams commute. For $i = 1, \dots, n$ the face map $\partial_i : \mathcal{NC}_n^T(\mathcal{C}) \rightarrow \mathcal{NC}_{n-1}^T(\mathcal{C})$ omits x_i and composes f_i and f_{i-1} . The face map ∂_0 sends (16) to

$$x_1 \xrightarrow{f_1} \dots \xrightarrow{f_{n-2}} x_{n-1} \xrightarrow{f_{n-1}} x_n \xrightarrow{T(f_0) \circ f_n} T(x_1).$$

The degeneracy maps insert identity morphisms at appropriate spots.

It is proved in [7, Theorem 3.2.3] that the (non-categorified) T -twisted cyclic nerve $\mathcal{NC}_\bullet^T(\mathcal{C})$ is a unital 2-Segal simplicial set. The analogous result holds also in the categorified setting.

Theorem 3.3. *For any small category \mathcal{C} and endofunctor $T : \mathcal{C} \rightarrow \mathcal{C}$, the simplicial groupoid $\mathcal{NC}_\bullet^T(\mathcal{C})$ is unital 2-Segal.*

Proof. Omitted. A similar result will be proved in Theorem 3.5 below. \square

In addition to the pair (\mathcal{C}, T) , suppose now that we are given a duality structure (P, Θ) on \mathcal{C} and a natural transformation

$$\lambda : P \Rightarrow T \circ P \circ T^{\text{op}}$$

which satisfies the compatibility condition

$$TP(\lambda_x) \circ \lambda_{PT(x)} \circ \Theta_{T(x)} = T(\Theta_x), \quad x \in \mathcal{C}. \quad (17)$$

For example, the pair $(T, \lambda) = (\mathbf{1}_{\mathcal{C}}, \mathbf{1}_P)$ satisfies this condition.

For each $n \geq 0$ define a functor $P_n : \mathcal{N}C_n^T(\mathcal{C})^{\text{op}} \rightarrow \mathcal{N}C_n^T(\mathcal{C})$ by sending the diagram (16) to

$$P(x_n) \xrightarrow{P(f_{n-1})} P(x_{n-1}) \xrightarrow{P(f_{n-2})} \dots \xrightarrow{P(f_1)} P(x_1) \xrightarrow{P(f_0)} P(x_0) \xrightarrow{P_*(f_n)} TP(x_n)$$

where $P_*(f_n) : P(x_0) \xrightarrow{\lambda_{x_0}} TPT(x_0) \xrightarrow{TP(f_n)} TP(x_n)$. The action of P_n on morphisms is the obvious one.

Lemma 3.4. *The data $\Theta_{n, x_\bullet} = (\Theta_{x_0}, \dots, \Theta_{x_n}, T(\Theta_{x_0}))$, $x_\bullet \in \mathcal{N}C_n^T(\mathcal{C})$, define a natural isomorphism $\Theta_n : \mathbf{1}_{\mathcal{N}C_n^T(\mathcal{C})} \Rightarrow P_n \circ P_n^{\text{op}}$, giving a triple $(\mathcal{N}C_\bullet^T(\mathcal{C}), P_\bullet, \Theta_\bullet)$ which satisfies the hypotheses of Lemma 3.1.*

Proof. We will prove that Θ_{n, x_\bullet} defines a morphism $x_\bullet \rightarrow P_n^2(x_\bullet)$; the remaining statements of the lemma can be verified directly. Keeping the notation (16), we need to show that the diagram

$$\begin{array}{ccc} x_n & \xrightarrow{f_n} & T(x_0) \\ \Theta_{x_n} \downarrow & & \downarrow T(\Theta_{x_0}) \\ P^2(x_n) & \xrightarrow{P_*^2(f_n)} & TP^2(x_0) \end{array}$$

commutes. The definition of P_* implies that

$$P_*^2(f_n) = TP(\lambda_{x_0}) \circ TPTP(f_n) \circ \lambda_{P(x_n)}.$$

The natural transformation λ associates to $x_n \xrightarrow{f_n} T(x_0)$ the commutative diagram

$$\begin{array}{ccc} TPTP(x_n) & \xrightarrow{TPTP(f_n)} & TPTPT(x_0) \\ \lambda_{P(x_n)} \uparrow & & \uparrow \lambda_{PT(x_0)} \\ P^2(x_n) & \xrightarrow{P^2(f_n)} & P^2T(x_0). \end{array}$$

Combining this with the equality $P^2(f_n) \circ \Theta_{x_n} = \Theta_{T(x_0)} \circ f_n$, we compute

$$\begin{aligned} P_*^2(f_0) \circ \Theta_{x_n} &= TP(\lambda_{x_0}) \circ TPTP(f_n) \circ \lambda_{P(x_n)} \circ \Theta_{x_n} \\ &= TP(\lambda_{x_0}) \circ \lambda_{PT(x_0)} \circ \Theta_{T(x_0)} \circ f_n \\ &= T(\Theta_{x_0}) \circ f_n, \end{aligned}$$

where in the final step the compatibility condition (17) was used. \square

Lemmas 3.1 and 3.4 imply that $\mathcal{N}C_\bullet^{T,e}(\mathcal{C})$ is a simplicial groupoid with duality.

Definition. *The Hermitian groupoid $\mathcal{N}C_\bullet^{T,e}(\mathcal{C})_h$, denoted by $\mathcal{N}U_\bullet^T(\mathcal{C})$, is called the unoriented categorified T -twisted cyclic nerve of $(\mathcal{C}, P, \Theta; T, \lambda)$.*

Write $(x_\bullet, \psi_\bullet) \in \mathcal{N}U_n^T(\mathcal{C})$ for the diagram

$$x_0 \xrightarrow{f_0} x_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-1}} x_n \xrightarrow{f_n} x_{n'} \xrightarrow{f_{n'}} x_{(n-1)'} \xrightarrow{f_{(n-1)'}} \cdots \xrightarrow{f_{1'}} x_{0'} \xrightarrow{f_{0'}} T(x_0)$$

together with symmetric isomorphisms ψ_\bullet which satisfy $\psi_{i+1} \circ f_i = P(f_{(i+1)'}) \circ \psi_i$, $i = 0, \dots, n-1$, and

$$\psi_{n'} \circ f_n = P(f_n) \circ \psi_n, \quad T(\psi_0) \circ f_{0'} = P_*(f_{0'}) \circ \psi_{0'}.$$

The following statement gives a relative 2-Segal space which plays the role of the unoriented loop space.

Theorem 3.5. *For any small category with duality (\mathcal{C}, P, Θ) and endofunctor $T : \mathcal{C} \rightarrow \mathcal{C}$ with compatibility data λ as above, the morphism $F_\bullet : \mathcal{N}U_\bullet^T(\mathcal{C}) \rightarrow \mathcal{N}C_\bullet^T(\mathcal{C})$ is a unital relative 2-Segal simplicial groupoid.*

Proof. We need to prove that the 1-Segal morphisms

$$\Xi_n : \mathcal{N}U_n^T(\mathcal{C}) \rightarrow \mathcal{N}U_i^T(\mathcal{C}) \times_{\mathcal{N}U_{\{i\}}^T(\mathcal{C})}^{(2)} \mathcal{N}U_{n-i}^T(\mathcal{C})$$

and the relative 2-Segal morphisms

$$\Psi_n : \mathcal{N}U_n^T(\mathcal{C}) \rightarrow \mathcal{N}C_n^T(\mathcal{C}) \times_{\mathcal{N}C_{\{0,n\}}^T(\mathcal{C})}^{(2)} \mathcal{N}U_{\{0,n\}}^T(\mathcal{C})$$

are equivalences. Since the proofs are similar we will only prove the latter. Explicitly, the functor Ψ_n sends $(x_\bullet, \psi_\bullet) \in \mathcal{N}U_n^T(\mathcal{C})$ to

$$(x_0 \rightarrow \cdots \rightarrow x_n \rightarrow T(x_0), (x_0 \rightarrow x_n \rightarrow x_{n'} \rightarrow x_{0'} \rightarrow T(x_0), \psi_\bullet); (\mathbf{1}_{x_0}, \mathbf{1}_{x_n})).$$

To see that Ψ_n is fully faithful, let $\Psi_n(x_\bullet, \psi_\bullet) \rightarrow \Psi_n(y_\bullet, \mu_\bullet)$ be a morphism determined by maps $((p_0, \dots, p_n), (q_0, q_n))$, the notation an obvious modification of that from the proof of Proposition 3.2. Since the isomorphisms α_0, α_n and β_0, β_n used to define arbitrary morphisms in $\mathcal{N}C_n^T(\mathcal{C}) \times_{\mathcal{N}C_{\{0,n\}}^T(\mathcal{C})}^{(2)} \mathcal{N}U_{\{0,n\}}^T(\mathcal{C})$ are the identities in this case, the compatibility conditions (analogous to equation (14)) imply $p_0 = q_0$ and $p_n = q_n$. Using the symmetry conditions imposed by the isometry condition, any morphism $\Psi_n(x_\bullet, \psi_\bullet) \rightarrow \Psi_n(y_\bullet, \mu_\bullet)$ is therefore the image of a unique morphism $(x_\bullet, \psi_\bullet) \rightarrow (y_\bullet, \mu_\bullet)$.

To prove that Ψ_n is essentially surjective, let

$$(x_0 \rightarrow \cdots \rightarrow x_n \rightarrow T(x_0), (x'_0 \rightarrow x'_{n'} \rightarrow x'_{n'} \rightarrow x'_{0'} \rightarrow T(x'_0), \psi_\bullet); (\alpha_0, \alpha_n)) \quad (18)$$

be an arbitrary object of $\mathcal{N}C_n^T(\mathcal{C}) \times_{\mathcal{N}C_{\{0,n\}}^T(\mathcal{C})}^{(2)} \mathcal{N}U_{\{0,n\}}^T(\mathcal{C})$. Define an object of $\mathcal{N}U_n^T(\mathcal{C})$ by the top row of the diagram

$$\begin{array}{ccccccc} x'_0 & \rightarrow & x_1 & \rightarrow & \cdots & \rightarrow & x_{n-1} & \rightarrow & x'_n & \rightarrow & x'_{n'} & \rightarrow & P(x_{n-1}) & \rightarrow & \cdots & \rightarrow & P(x_1) & \rightarrow & x'_{0'} & \rightarrow & T(x'_0) \\ \alpha_0^{-1} \downarrow & \nearrow & & & & & & & \nearrow & \alpha_n & \psi_{n'} \downarrow & \nearrow & & & & & & & \downarrow & \nearrow & \psi_{0'}^{-1} \\ & & x_0 & & & & x_n & & P(x'_n) & & & & & & & & & & & P(x'_0) & & \end{array}$$

with symmetric form $(\{\Theta_\bullet\}_{[1,n-1]}, \{\psi_\bullet\}_{\{0,n,n',0'\}}, \{\mathbf{1}_\bullet\}_{[(n-1)',1']})$. The image of this object under Ψ_n is

$$(x'_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_{n-1} \rightarrow x'_n \rightarrow T(x'_0), (x'_0 \rightarrow \cdots \rightarrow x'_{0'} \rightarrow T(x'_0), \psi_\bullet); (\mathbf{1}_{x'_0}, \mathbf{1}_{x'_n})).$$

Then the tuple $(\alpha_0^{-1}, \mathbf{1}_{x_1}, \dots, \mathbf{1}_{x_{n-1}}, \alpha_n^{-1}, (\mathbf{1}_{x'_0}, \dots, \mathbf{1}_{x'_{0'}}))$ defines an isomorphism from the previous object to the object (18). Hence Ψ_n is an equivalence.

The proof that relative unitality holds is similar. \square

Remark. The homotopical counterpart of the $\mathrm{SO}(2)$ -action on LX is the cyclic structure of $NC_\bullet(\mathcal{C})$ in the sense of Connes. This action extends to $\mathrm{O}(2)$ by allowing orientation reversal of loops and corresponds to the dihedral structure of $NC_\bullet(\mathcal{C})$ induced by a (strict) duality P on \mathcal{C} . More generally, $NC_\bullet^T(\mathcal{C})$ has a paradihedral or, if $T^r = 1$ for some $r \geq 2$, an r -dihedral structure. Similar comments apply to categorified nerves. See [9, §1.6] for further discussion.

3.4. The \mathcal{R}_\bullet -construction. The goal of this section is to prove that the \mathcal{R}_\bullet -construction (also called the Hermitian \mathcal{S}_\bullet -construction) from Grothendieck-Witt theory is relative 2-Segal over the Waldhausen \mathcal{S}_\bullet -construction. We work in the proto-exact setting of Section 1.4.

Definition. A proto-exact category with duality is a category with duality (\mathcal{C}, P, Θ) such that $\mathcal{C} = (\mathcal{C}, \mathfrak{I}, \mathfrak{D})$ is proto-exact and P is an exact functor. Explicitly, the latter condition comprises of the following:

- (i) $P(0) \simeq 0$,
- (ii) a morphism $U \xrightarrow{\phi} V$ is in \mathfrak{I} if and only if $P(V) \xrightarrow{P(\phi)} P(U)$ is in \mathfrak{D} , and
- (iii) P sends biCartesian squares to biCartesian squares.

Let N be a symmetric form in a proto-exact category with duality \mathcal{C} and let $i : U \rightarrow N$ be an inflation. The orthogonal U^\perp is defined to be the pullback

$$\begin{array}{ccc} U^\perp & \xrightarrow{\quad} & N \\ \downarrow & & \downarrow \\ 0 & \xrightarrow{\quad} & P(U). \end{array}$$

The inflation i is called isotropic if the composition $P(i)\psi_N i$ is zero and the canonical monomorphism $U \rightarrow U^\perp$ is an inflation.

Example.

- (1) Let $\mathrm{Rep}_k(Q)$ be the abelian category of representations of quiver Q . Given a contravariant involution of Q together with some combinatorial data, there is an associated exact duality structure on $\mathrm{Rep}_k(Q)$ whose symmetric forms are (generalizations of) the orthogonal or symplectic quiver representations of Derksen and Weyman [4]. See also [48, §1.4]. A similar construction applies when $\mathrm{Rep}_k(Q)$ is replaced with $\mathrm{Rep}_{\mathbb{F}_1}(Q)$.
- (2) Let $\mathrm{Vect}(X)$ be the exact category of vector bundles over a scheme X . Fix a line bundle $L \rightarrow X$ and a sign $s \in \{\pm 1\}$. Then $(P, \Theta) = (\mathrm{Hom}_{\mathcal{O}_X}(-, L), s \cdot \mathrm{can})$ defines an exact duality structure on $\mathrm{Vect}(X)$ whose symmetric forms are vector bundles over X with L -valued orthogonal ($s = 1$) or symplectic ($s = -1$) forms. \triangleleft

In this section we will make the following assumption on (\mathcal{C}, P, Θ) .

Reduction Assumption. Let N be a symmetric form in \mathcal{C} and let $U \rightarrow N$ be isotropic with orthogonal $k : U^\perp \rightarrow N$. Define $M \in \mathcal{C}$ to be the pushout

$$\begin{array}{ccc} U & \xrightarrow{\quad} & U^\perp \\ \downarrow & & \downarrow \pi \\ 0 & \xrightarrow{\quad} & M. \end{array}$$

Then there is a unique symmetric form ψ_M on M such that $P(k)\psi_N k = P(\pi)\psi_M \pi$.

The symmetric form (M, ψ_M) is called the isotropic reduction of N by U and is denoted by $N//U$. When \mathcal{C} is exact the Reduction Assumption is known to hold;

see [30, Lemma 5.2], [35, Lemma 2.6]. In a number of (non-exact) proto-exact examples, such as $\text{Rep}_{\mathbb{F}_1}(Q)$, the Reduction Assumption also holds.

The category $[n]$ has a strict duality structure given at the level of objects by $i \mapsto n - i$. Then $[\text{Ar}_n, \mathcal{C}]$ and its full subcategory $\mathcal{W}_n(\mathcal{C})$ inherit duality structures. Moreover, the duality structures on $\mathcal{W}_\bullet(\mathcal{C})$ satisfy the assumptions of Lemma 3.1 so that $\mathcal{W}_\bullet^e(\mathcal{C})$ is a simplicial object of CatD . Explicitly, the dual of a diagram $\{A_{\{p,q\}}\}_{0 \leq p \leq q \leq 0'} \in \mathcal{W}_n^e(\mathcal{C})$ is the diagram $\{P(A_{\{q',p'\}})\}_{0 \leq p \leq q \leq 0'}$ and the double dual identification at (p, q) is $\Theta_{A_{\{p,q\}}}$.

The \mathcal{R}_\bullet -construction of (\mathcal{C}, P, Θ) , denoted by $\mathcal{R}_\bullet(\mathcal{C})$, is the Hermitian groupoid $\mathcal{W}_\bullet^e(\mathcal{C})_h$; see [39], [15].¹ Objects of $\mathcal{R}_n(\mathcal{C})$ are diagrams $\{A_{\{p,q\}}\}_{0 \leq p \leq q \leq 0'} \in \mathcal{W}_n^e(\mathcal{C})$ together with symmetric isomorphisms $\psi_{p,q} : A_{\{p,q\}} \rightarrow P(A_{\{q',p'\}})$ which make all appropriate diagrams commute. In particular,

- (i) for every $1 \leq i \leq n$ the pair $(A_{\{i,i'\}}, \psi_{i,i'})$ is a symmetric form,
- (ii) for every $0 \leq i \leq j \leq n$ the inflation $A_{\{i,j\}} \rightsquigarrow A_{\{i,i'\}}$ is isotropic with orthogonal $A_{\{i,j'\}} \rightsquigarrow A_{\{i,i'\}}$, and
- (iii) for every $0 \leq i \leq j \leq n$ the symmetric form $(A_{\{j,j'\}}, \psi_{j,j'})$ is isometric to the reduction $(A_{\{i,i'\}}, \psi_{i,i'}) // A_{\{i,j\}}$.

For example, an object of $\mathcal{R}_0(\mathcal{C})$ is a diagram

$$\begin{array}{c} 0 \rightsquigarrow A_{\{0,0'\}} \\ \downarrow \\ 0 \end{array}$$

together with a symmetric form on $A_{\{0,0'\}}$. Hence $\mathcal{R}_0(\mathcal{C})$ is equivalent to the Hermitian groupoid \mathcal{C}_h . Similarly, an object of $\mathcal{R}_1(\mathcal{C})$ is a diagram

$$\begin{array}{ccccc} 0 \rightsquigarrow & A_{\{0,1\}} & \rightsquigarrow & A_{\{0,1'\}} & \rightsquigarrow & A_{\{0,0'\}} \\ & \downarrow & & \downarrow & & \downarrow \\ & 0 \rightsquigarrow & & A_{\{1,1'\}} & \rightsquigarrow & A_{\{1,0'\}} \\ & & & \downarrow & & \downarrow \\ & & & 0 \rightsquigarrow & & A_{\{1',0'\}} \\ & & & & & \downarrow \\ & & & & & 0 \end{array}$$

all of whose squares are biCartesian, together with the data of

- (i) a symmetric form on $A_{\{0,0'\}}$ such that $A_{\{0,1\}} \rightsquigarrow A_{\{0,0'\}}$ is isotropic with orthogonal $A_{\{0,1'\}} \rightsquigarrow A_{\{0,0'\}}$,
- (ii) a symmetric form on $A_{\{1,1'\}}$ presenting $A_{\{1,1'\}}$ as $A_{\{0,0'\}} // A_{\{0,1\}}$, and
- (iii) isomorphisms $A_{\{0,1\}} \simeq P(A_{\{1',0'\}})$ and $A_{\{0,1'\}} \simeq P(A_{\{1,0'\}})$ such that each morphism above the diagonal agrees with the corresponding morphism below the diagonal.

In short, objects of $\mathcal{R}_n(\mathcal{C})$ are isotropic n -flags together with presentations of all subquotients and subreductions. The forgetful map

$$F_\bullet : \mathcal{R}_\bullet(\mathcal{C}) \rightarrow \mathcal{S}_\bullet(\mathcal{C}), \quad \mathcal{R}_n(\mathcal{C}) \ni \{A_{\{p,q\}}, \psi_{p,q}\}_{0 \leq p \leq q \leq 0'} \mapsto \{A_{\{p,q\}}\}_{0 \leq p \leq q \leq n}$$

is a morphism of simplicial groupoids.

¹In [15] the notation $\mathcal{R}_\bullet(\mathcal{C})$ is used for $\mathcal{W}_\bullet^e(\mathcal{C})$, whereas what we call $\mathcal{R}_\bullet(\mathcal{C})$ is denoted by $\mathcal{R}_\bullet^h(\mathcal{C})$.

Theorem 3.6. *Let \mathcal{C} be a proto-exact category with duality which satisfies the Reduction Assumption. Then the morphism $F_\bullet : \mathcal{R}_\bullet(\mathcal{C}) \rightarrow \mathcal{S}_\bullet(\mathcal{C})$ is a unital relative 2-Segal simplicial groupoid.*

Proof. For each $n \geq 0$ let $\mathcal{I}_n(\mathcal{C})$ be the groupoid of isotropic n -flags in \mathcal{C} . An object of $\mathcal{I}_n(\mathcal{C})$ is a diagram

$$0 \succ A_1 \succ \cdots \succ A_n \succ A_{n'} \succ \cdots \succ A_{1'} \succ (A_{0'}, \psi_{0'}) \quad (19)$$

with $(A_{0'}, \psi_{0'})$ a symmetric form such that, for each $0 \leq i \leq n$, the inflation $A_i \succ A_{0'}$ is isotropic with orthogonal $A_{i'} \succ A_{0'}$. We claim that the forgetful functor

$$\nu_n : \mathcal{R}_n(\mathcal{C}) \rightarrow \mathcal{I}_n(\mathcal{C}), \quad \{A_{\{i,j\}}, \psi_{i,j}\}_{0 \leq i \leq j \leq 0'} \mapsto \{A_{\{0,j\}}, \psi_{0,0'}\}_{0 \leq j \leq 0'}$$

is an equivalence. Construct a quasi-inverse η_n of ν_n as follows. Given an isotropic flag (19), put $A_{\{0,k\}} = A_k$, $1 \leq k \leq 1'$, and $(A_{\{0,0'\}}, \psi_{0,0'}) = (A_{0'}, \psi_{0'})$ and let $A_{\{1,k\}}$ be the pushout

$$\begin{array}{ccc} A_{\{0,1\}} & \xrightarrow{\quad} & A_{\{0,k\}} \\ \downarrow & & \downarrow \\ 0 & \xrightarrow{\quad\quad\quad} & A_{\{1,k\}}. \end{array}$$

By the Reduction Assumption the symmetric form $(A_{\{0,0'\}}, \psi_{0,0'})$ induces a unique compatible symmetric form on $A_{\{1,1'\}}$. Define also $A_{\{1,0'\}} = P(A_{\{0,1'\}})$ and let $A_{\{0,0'\}} \twoheadrightarrow A_{\{1,0'\}}$ be the composition

$$A_{\{0,0'\}} \xrightarrow{\psi_{0,0'}} P(A_{\{0,0'\}}) \twoheadrightarrow P(A_{\{0,1'\}}).$$

This construction defines the top two rows of $\eta_n(A_{\{0,\bullet\}})$ and can be iterated to define the remaining $n - 2$ rows. It is clear that ν_n is a quasi-inverse to η_n .

We can now prove the theorem. The 1-Segal morphism for $\mathcal{R}_\bullet(\mathcal{C})$ is

$$\Xi_n : \mathcal{R}_n(\mathcal{C}) \rightarrow \mathcal{R}_{\{0,\dots,i\}}(\mathcal{C}) \times_{\mathcal{R}_{\{i\}}(\mathcal{C})}^{(2)} \mathcal{R}_{\{i,\dots,n\}}(\mathcal{C}).$$

Arguing as in the proof of Theorem 2.7 and using that ν_n is an equivalence, to prove that Ξ_n is an equivalence it suffices to prove that the functor

$$\tilde{\Xi}_n : \mathcal{I}_n(\mathcal{C}) \rightarrow \mathcal{I}_i(\mathcal{C}) \times_{\mathcal{I}_{\{i\}}(\mathcal{C})}^{(2)} \mathcal{I}_{n-i}(\mathcal{C})$$

is an equivalence. A quasi-inverse of $\tilde{\Xi}_n$ is defined by assigning to a pair

$$(0 \succ A'_1 \succ \cdots \succ A'_i \succ A'_{i'} \succ \cdots \succ A'_{1'} \succ (A'_{0'}, \psi'_{0'})) \in \mathcal{I}_i(\mathcal{C})$$

and

$$(0 \succ A''_{i+1} \succ \cdots \succ A''_n \succ A''_{n'} \succ \cdots \succ A''_{(i+1)'}) \in \mathcal{I}_{n-i}(\mathcal{C})$$

together with an isometry $(A'_{0'}, \psi'_{0'}) // A'_i \simeq (A''_{i'}, \psi''_{i'})$ the object

$$(0 \succ A'_1 \succ \cdots \succ A'_n \succ A'_{n'} \succ \cdots \succ A'_{1'} \succ (A'_{0'}, \psi'_{0'})) \in \mathcal{I}_n(\mathcal{C})$$

where A'_j , $i + 1 \leq j \leq (i + 1)'$, is defined to be the pullback

$$\begin{array}{ccc} A'_j & \xrightarrow{\quad\quad\quad} & A'_{i'} \\ \downarrow & & \downarrow \\ A''_j & \xrightarrow{\quad\quad\quad} & A''_{i'}. \end{array}$$

That this is indeed a quasi-inverse follows from the Hermitian variant of the Second Isomorphism Theorem [30, Proposition 6.5], which generalizes to the proto-exact setting under the Reduction Assumption.

Arguing in the same way, to prove that the functor

$$\Psi_n : \mathcal{R}_n(\mathcal{C}) \rightarrow \mathcal{S}_n(\mathcal{C}) \times_{\mathcal{S}_{\{0,n\}}(\mathcal{C})}^{(2)} \mathcal{R}_{\{0,n\}}(\mathcal{C})$$

is an equivalence it suffices to prove that the functor

$$\tilde{\Psi}_n : \mathcal{I}_n(\mathcal{C}) \rightarrow \mathcal{F}_n(\mathcal{C}) \times_{\mathcal{F}_{\{0,n\}}(\mathcal{C})}^{(2)} \mathcal{I}_{\{0,n\}}(\mathcal{C})$$

is an equivalence, which is obvious.

Finally, relative unitality is the condition that, for each $0 \leq i \leq n-1$, the functor

$$\Upsilon_n : \mathcal{R}_{n-1}(\mathcal{C}) \rightarrow \mathcal{S}_{\{i\}}(\mathcal{C}) \times_{\mathcal{S}_{\{i,i+1\}}(\mathcal{C})}^{(2)} \mathcal{R}_n(\mathcal{C})$$

is an equivalence. Note that $\mathcal{S}_0(\mathcal{C})$ is a point, $\mathcal{S}_1(\mathcal{C})$ is the maximal groupoid of \mathcal{C} and the map $\mathcal{S}_0(\mathcal{C}) \xrightarrow{s_0} \mathcal{S}_1(\mathcal{C})$ sends the point to the zero object. An object of $\mathcal{S}_{\{i\}}(\mathcal{C}) \times_{\mathcal{S}_{\{i,i+1\}}(\mathcal{C})}^{(2)} \mathcal{R}_n(\mathcal{C})$ is therefore an object of $\mathcal{R}_n(\mathcal{C})$ whose maps between the i th and $(i+1)$ st rows/columns are the identities. It follows from this that Υ_n is an equivalence. \square

When \mathcal{C} is exact, the morphism $F_\bullet : \mathcal{R}_\bullet(\mathcal{C}) \rightarrow \mathcal{S}_\bullet(\mathcal{C})$ is closely related to the higher Grothendieck-Witt theory of \mathcal{C} . Indeed, $GW_i(\mathcal{C})$ is the i th homotopy group of the homotopy fibre over $0 \in \mathcal{C}$ of the map $|BF_\bullet| : |B\mathcal{R}_\bullet(\mathcal{C})| \rightarrow |B\mathcal{S}_\bullet(\mathcal{C})|$ [36].

4. APPLICATIONS TO HALL ALGEBRAS

4.1. Categorical Hall algebra representations. After reviewing the relationship between 2-Segal spaces and categorical Hall algebras [7], in this section we construct from a relative 2-Segal space a categorical representation of the Hall algebra of the base.

Fix a combinatorial model category \mathcal{C} , such as \mathbb{S} with its Kan model structure or \mathbf{Grpd} with its Bousfield model structure; the latter will be the main source of examples. While the Quillen model structure on \mathbf{Top} is not combinatorial, it is Quillen equivalent to \mathbb{S} so that the results of this section, for all intents and purposes, also apply to simplicial spaces. We write $\text{pt} \in \mathcal{C}$ for the final object.

Let $\mathbf{Span}(\mathcal{C})$ be the bicategory of spans in \mathcal{C} . Objects of $\mathbf{Span}(\mathcal{C})$ are simply objects of \mathcal{C} while 1-morphisms $A \rightarrow B$ in $\mathbf{Span}(\mathcal{C})$ are spans from A to B , that is, diagrams in \mathcal{C} of the form

$$A \leftarrow X \rightarrow B.$$

Composition of spans is given by homotopy pullback. A 2-morphism in $\mathbf{Span}(\mathcal{C})$ is a homotopy commutative diagram

$$\begin{array}{ccc} & X & \\ & \swarrow & \searrow \\ A & & B \\ & \nwarrow & \nearrow \\ & X' & \end{array}$$

in \mathcal{C} . Give $\mathbf{Span}(\mathcal{C})$ the Cartesian monoidal structure.

Associated to the bicategory $\mathbf{Span}(\mathcal{C})$ is the category $\mathbf{Span}(\mathcal{C})^\sim$, having the same objects as $\mathbf{Span}(\mathcal{C})$ but with morphisms being the 2-isomorphism classes spans. The constructions of this section are simplified if one uses $\mathbf{Span}(\mathcal{C})^\sim$ in place of $\mathbf{Span}(\mathcal{C})$, the downside being that less of the (relative) 2-Segal structure is used.

Definition ([7, §8.1]). *A transfer structure on \mathcal{C} is a pair $(\mathcal{S}, \mathcal{P})$ consisting of collections of morphisms \mathcal{S} and \mathcal{P} in \mathcal{C} called smooth and proper, respectively, which satisfy the following properties:*

- (1) *Both collections \mathcal{S} and \mathcal{P} are closed under composition.*

(2) Given a homotopy Cartesian diagram

$$\begin{array}{ccc} X & \xrightarrow{s} & Y \\ p \downarrow & & \downarrow p' \\ X' & \xrightarrow{s'} & Y' \end{array} \quad (20)$$

in \mathcal{C} with $s' \in \mathcal{S}$ and $p' \in \mathcal{P}$, then $s \in \mathcal{S}$ and $p \in \mathcal{P}$.

Example. The pair $(\mathcal{S}, \mathcal{P}) = (\mathcal{C}, \mathcal{C})$ is called the trivial transfer structure on \mathcal{C} . \triangleleft

Fix a transfer structure $(\mathcal{S}, \mathcal{P})$. Spans of the form $A \xleftarrow{s} X \xrightarrow{p} B$ with $s \in \mathcal{S}$ and $p \in \mathcal{P}$ are called $(\mathcal{S}, \mathcal{P})$ -admissible and form a subcategory $\text{Span}(\mathcal{S}, \mathcal{P}) \subset \text{Span}(\mathcal{C})$.

Theorem 4.1 ([7, Proposition 8.1.7]). *Let X_\bullet be a 2-Segal object of \mathcal{C} . Assume that the span*

$$m_X = \left\{ X_{\{0,1\}} \times X_{\{1,2\}} \xleftarrow{(\partial_2, \partial_0)} X_{\{0,1,2\}} \xrightarrow{\partial_1} X_{\{0,2\}} \right\} \in \text{Hom}_{\text{Span}(\mathcal{C})}(X_1 \otimes X_1, X_1) \quad (21)$$

is $(\mathcal{S}, \mathcal{P})$ -admissible. Then (X_1, m_X) is a semigroup in $\text{Span}(\mathcal{S}, \mathcal{P})$. Moreover, if X_\bullet is unital and the span

$$I_X = \left\{ \text{pt} \xleftarrow{\text{can}} X_0 \xrightarrow{s_0} X_1 \right\} \in \text{Hom}_{\text{Span}(\mathcal{C})}(\text{pt}, X_1)$$

is $(\mathcal{S}, \mathcal{P})$ -admissible, then (X_1, m_X, I_X) is a monoid in $\text{Span}(\mathcal{S}, \mathcal{P})$.

The pair $\mathcal{H}(X_\bullet) = (X_1, m_X)$ is called the $(\mathcal{S}, \mathcal{P})$ -universal Hall algebra of X_\bullet . Slightly abusively, the associativity isomorphism a_X for m_X is omitted from the notation. Note that at the universal level the transfer structure $(\mathcal{S}, \mathcal{P})$ simply determines the subcategory $\text{Span}(\mathcal{S}, \mathcal{P}) \subset \text{Span}(\mathcal{C})$ to which $\mathcal{H}(X_\bullet)$ belongs. In particular, we can always take the trivial transfer structure. A more important role is played by the transfer structure when passing from the universal Hall algebra to its concrete realizations. To explain this procedure, fix a monoidal category $(\mathcal{V}, \otimes, \mathbf{1}_{\mathcal{V}})$.

Definition ([7, §8.1]). *A \mathcal{V} -valued theory with transfer on \mathcal{C} is the data of*

- (1) a transfer structure $(\mathcal{S}, \mathcal{P})$ on \mathcal{C} ,
- (2) a contravariant functor $(-)^* : \mathcal{S} \rightarrow \mathcal{V}$ and a covariant functor $(-)_* : \mathcal{P} \rightarrow \mathcal{V}$ with common values on objects, denoted by \mathfrak{h} , and which map weak equivalences to isomorphisms, and
- (3) an isomorphism $\mathfrak{h}(\text{pt}) \simeq \mathbf{1}_{\mathcal{V}}$ and multiplicativity data for \mathfrak{h} , that is, natural maps

$$\mathfrak{h}(X) \otimes \mathfrak{h}(Y) \rightarrow \mathfrak{h}(X \otimes Y)$$

which satisfy associativity and unitality conditions

such that for a homotopy Cartesian diagram of the form (20) with $s, s' \in \mathcal{S}$ and $p, p' \in \mathcal{P}$, we have $p'_* \circ s^* = s'^* \circ p_*$.

Applying a \mathcal{V} -valued theory with transfer \mathfrak{h} to the $(\mathcal{S}, \mathcal{P})$ -universal Hall algebra $\mathcal{H}(X_\bullet)$ gives a semigroup in \mathcal{V} , denoted by

$$\mathcal{H}(X_\bullet; \mathfrak{h}) = (\mathfrak{h}(X_1), \partial_{1*} \circ (\partial_2 \times \partial_0)^*).$$

Dually, if the opposite span $m_X^{\text{op}} \in \text{Hom}_{\text{Span}(\mathcal{C})}(X_1, X_1 \otimes X_1)$ is $(\mathcal{S}, \mathcal{P})$ -admissible, then (X_1, m_X^{op}) is a cosemigroup in $\text{Span}(\mathcal{S}, \mathcal{P})$, called the $(\mathcal{S}, \mathcal{P})$ -universal Hall coalgebra, and passing to theories with transfer gives cosemigroups in monoidal categories. These statements can be proved in the same way as Theorem 4.1.

Example. Let $\mathcal{S}_\bullet(\mathcal{C})$ be the Waldhausen groupoid of an essentially small exact category \mathcal{C} . Suppose that \mathcal{C} is finitary in the sense that $\text{Ext}_{\text{env}(\mathcal{C})}^n(U, V)$, $n = 0, 1$, is finite for all $U, V \in \mathcal{C}$. Here $\text{env}(\mathcal{C})$ denotes the abelian envelope of \mathcal{C} [31]. Then $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}))$ categorifies the Hall algebra of \mathcal{C} , as defined in various contexts in [33], [16], [34]. To see this, let k be a field of characteristic zero. Consider the transfer structure on Grpd in which \mathcal{S} and \mathcal{P} are the collections of weakly proper and locally proper morphisms, respectively (see [7, §8.2]). A Vect_k -valued theory with transfer \mathfrak{F}_0 on Grpd is then defined by taking finitely supported k -valued functions which are constant on isomorphism classes. The resulting k -algebra $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ is the standard Hall algebra of \mathcal{C} . Explicitly, $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ is the k -vector space with basis $\{\mathbf{1}_U\}_{U \in \pi_0(\mathcal{S}_1(\mathcal{C}))}$ and multiplication

$$\mathbf{1}_U \cdot \mathbf{1}_V = \sum_{W \in \pi_0(\mathcal{S}_1(\mathcal{C}))} F_{U,V}^W \mathbf{1}_W,$$

where $F_{U,V}^W$ is the number of admissible subobjects of W which are isomorphic to U and have quotient isomorphic to V .

For certain categories \mathcal{C} , such as $\text{Coh}(X)$ for a smooth projective variety X or $\text{mod}(A)$ for a finitely generated algebra A , the \mathcal{S}_\bullet -construction defines a 2-Segal simplicial Artin stack. This allows to recover the perverse sheaf theoretic [26], motivic [20], [23] and cohomological [24] Hall algebras of \mathcal{C} . See [7, §8.5]. \triangleleft

By using relative 2-Segal spaces we can easily modify the above results to construct module objects.

Theorem 4.2. *Let X_\bullet be as in Theorem 4.1 and let $F_\bullet : Y_\bullet \rightarrow X_\bullet$ be a relative 2-Segal object of \mathcal{C} . Assume that the left action span*

$$\mu_Y^l = \left\{ X_{\{0,1\}} \times Y_{\{1\}} \xleftarrow{(F_1, \partial_0)} Y_{\{0,1\}} \xrightarrow{\partial_1} Y_{\{1\}} \right\} \in \text{Hom}_{\text{Span}(\mathcal{C})}(X_1 \otimes Y_0, Y_0) \quad (22)$$

is $(\mathcal{S}, \mathcal{P})$ -admissible. Then (Y_0, μ_Y^l) is a left (X_1, m_X) -module in $\text{Span}(\mathcal{S}, \mathcal{P})$. Moreover, if F_\bullet is unital, then (Y_0, μ_Y^l) is a left (X_1, m_X, I_X) -module. Analogous statements hold for the right action span

$$\mu_Y^r = \left\{ Y_{\{0\}} \times X_{\{0,1\}} \xleftarrow{\partial_1 \times F_1} Y_{\{0,1\}} \xrightarrow{\partial_0} Y_{\{1\}} \right\} \in \text{Hom}_{\text{Span}(\mathcal{C})}(Y_0 \otimes X_1, Y_0).$$

Proof. We will prove the theorem for the left action span. It follows directly from the definitions that we have

$$\mu_Y^l \circ (m_X \otimes \mathbf{1}_{Y_0}) = \left\{ \begin{array}{c} (X_{\{0,1,2\}} \times Y_{\{2\}}) \times_{X_{\{0,2\}} \times Y_{\{2\}}}^R Y_{\{0,2\}} \longrightarrow Y_{\{0,2\}} \longrightarrow Y_{\{0\}} \\ \downarrow \\ X_{\{0,1,2\}} \times Y_{\{2\}} \longrightarrow X_{\{0,2\}} \times Y_{\{2\}} \\ \downarrow \\ X_{\{0,1\}} \times X_{\{1,2\}} \times Y_{\{2\}} \end{array} \right\}$$

and

$$\mu_Y^l \circ (\mathbf{1}_{X_1} \otimes \mu_Y^l) = \left\{ \begin{array}{c} (X_{\{0,1\}} \times Y_{\{1,2\}}) \times_{X_{\{0,1\}} \times Y_{\{1\}}}^R Y_{\{0,1\}} \longrightarrow Y_{\{0,1\}} \longrightarrow Y_{\{0\}} \\ \downarrow \\ X_{\{0,1\}} \times Y_{\{1,2\}} \longrightarrow X_{\{0,1\}} \times Y_{\{1\}} \\ \downarrow \\ X_{\{0,1\}} \times X_{\{1,2\}} \times Y_{\{2\}} \end{array} \right\}$$

as spans $X_1 \otimes X_1 \otimes Y_0 \rightarrow Y_0$. We claim that both $\mu_Y^l \circ (m_X \otimes \mathbf{1}_{Y_0})$ and $\mu_Y^l \circ (\mathbf{1}_{X_1} \otimes \mu_Y^l)$ are 2-isomorphic to the span

$$\sigma^l = \left\{ X_{\{0,1\}} \times X_{\{1,2\}} \times Y_{\{2\}} \xleftarrow{(F_{\{0,1\}} \circ \partial_2, F_{\{1,2\}} \circ \partial_0, \partial_0 \circ \partial_1)} Y_{\{0,1,2\}} \xrightarrow{\partial_1 \circ \partial_2} Y_{\{0\}} \right\}.$$

Indeed, the composition

$$\begin{aligned} Y_{\{0,1,2\}} &\xrightarrow{\alpha_1} (X_{\{0,1,2\}} \times Y_{\{2\}}) \times_{X_{\{0,2\}} \times Y_{\{2\}}}^R Y_{\{0,2\}} \\ &\rightarrow X_{\{0,1,2\}} \times_{X_{\{0,2\}}}^R Y_{\{0,2\}} \end{aligned}$$

is a weak equivalence by the relative 2-Segal condition on F_\bullet . Since the second functor is a weak equivalence for trivial reasons, it follows that α_1 is a weak equivalence and so defines a 2-isomorphism $\sigma^l \rightarrow \mu_Y^l \circ (m_X \otimes \mathbf{1}_{Y_0})$. Similarly, the composition

$$\begin{aligned} Y_{\{0,1,2\}} &\xrightarrow{\alpha_2} (X_{\{0,1\}} \times Y_{\{1,2\}}) \times_{X_{\{0,1\}} \times Y_{\{1\}}}^R Y_{\{0,1\}} \\ &\rightarrow Y_{\{1,2\}} \times_{Y_{\{1\}}}^R Y_{\{0,1\}} \end{aligned}$$

is a weak equivalence by the 1-Segal condition on Y_\bullet and α_2 defines a 2-isomorphism $\sigma^l \rightarrow \mu_Y^l \circ (\mathbf{1}_{X_1} \otimes \mu_Y^l)$. We claim that the composition

$$\alpha_Y^l : \mu_Y^l \circ (m_X \otimes \mathbf{1}_{Y_0}) \xrightarrow{\alpha_1^{-1}} \sigma^l \xrightarrow{\alpha_2} \mu_Y^l \circ (\mathbf{1}_{X_1} \otimes \mu_Y^l)$$

is a module associator for the left action of (X_1, m_X) on Y_0 determined by μ_Y^l . Without the unital assumption, this amounts to verifying that α_Y^l satisfies module theoretic Mac Lane coherence; see [28, §2.3] or diagram (23) below. We will verify this using the setting of Section 2.3. The poset of the 11 polyhedral subdivisions of the relative pentagon P_3^∞ , ordered by refinement, is illustrated in Figure 3. At the level of the map $F_\bullet : Y_\bullet \rightarrow X_\bullet$, each node \mathcal{P} of Figure 3 defines a span

$$X_1 \times X_1 \times X_1 \times Y_0 \leftarrow (\Delta^{\mathcal{P}}, F_\bullet)_R \rightarrow Y_0.$$

Similarly, each arrow of Figure 3 defines a 2-isomorphism of spans. The spans associated to the five vertices of the pentagon are precisely the spans appearing in the diagram expressing module theoretic Mac Lane coherence while the composed 2-isomorphisms along the edges of the pentagon are precisely the arrows in the coherence diagram. It follows that module theoretic Mac Lane coherence holds. Hence (Y_0, μ_Y^l) is a left (X_1, m_X) -module.


Finally, in the unital setting the action of I_X on Y_0 is given by the span

$$\begin{array}{ccccc} (X_{\{0\}} \times Y_{\{1\}}) \times_{X_{\{0,1\}} \times Y_{\{1\}}}^R Y_{\{0,1\}} & \longrightarrow & Y_{\{0,1\}} & \longrightarrow & Y_{\{0\}} \\ \downarrow & & \downarrow & & \\ X_{\{0\}} \times Y_{\{1\}} & \longrightarrow & X_{\{0,1\}} \times Y_{\{1\}} & & \\ \downarrow & & & & \\ \text{pt} \times Y_{\{1\}} & & & & \end{array}$$

which is 2-isomorphic to the identity span $Y_0 \xleftarrow{\mathbf{1}_{Y_0}} Y_0 \xrightarrow{\mathbf{1}_{Y_0}} Y_0$. Indeed, the composition of functors

$$\begin{aligned} Y_{\{0\}} &\xrightarrow{I_Y} (X_{\{0\}} \times Y_{\{1\}}) \times_{X_{\{0,1\}} \times Y_{\{1\}}}^R Y_{\{0,1\}} \\ &\rightarrow X_{\{0\}} \times_{X_{\{0,1\}}}^R Y_{\{0,1\}} \end{aligned}$$

is an equivalence by the unital relative 2-Segal condition while the second functor is trivially an equivalence. To complete the proof we need to verify that I_Y is compatible with the module associator in the sense that diagram (24) below commutes.


 FIGURE 3. The poset of polyhedral subdivisions of P_3^∞ .

This is a straightforward exercise which can be completed in much the same way as the above verification of Mac Lane coherence. \square

The left $\mathcal{H}(X_\bullet)$ -module $\mathcal{M}(Y_\bullet) = (Y_0, \mu_Y^l)$ is called the $(\mathcal{S}, \mathcal{P})$ -universal left Hall module of F_\bullet . From a \mathcal{V} -valued theory with transfer \mathfrak{h} we obtain a left $\mathcal{H}(X_\bullet; \mathfrak{h})$ -module $\mathcal{M}(Y_\bullet; \mathfrak{h})$ in \mathcal{V} . In the same way we get right modules over $\mathcal{H}(X_\bullet)$ and $\mathcal{H}(X_\bullet; \mathfrak{h})$. Note that the left and right module structures do not define a bimodule.

Example. Let X_\bullet be 2-Segal object of \mathcal{C} . By Proposition 2.5 the right path space $F_n^\triangleright : P^\triangleright X_\bullet \rightarrow X_\bullet$ is relative 2-Segal. Since $F_n^\triangleright = \partial_{n'}$, the spans $\mu_{P^\triangleright X}^l$ and m_X are equal. Hence $\mathcal{M}(X_n^\triangleright) = \mathcal{H}(X_\bullet)$ as left $\mathcal{H}(X_\bullet)$ -modules. On the other hand, the right $\mathcal{H}(X_\bullet)$ -module structure on $\mathcal{M}(X_n^\triangleright)$ is closely related to the coproduct Δ on $\mathcal{H}(X_\bullet)$. Consider for example $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ for a finitary exact category \mathcal{C} . Recall that the Green bilinear form on $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$, defined by

$$(\mathbf{1}_U, \mathbf{1}_V) = \frac{\delta_{U,V}}{|\text{Aut}(U)|},$$

satisfies the Hopf property $(\mathbf{1}_U \otimes \mathbf{1}_V, \Delta \mathbf{1}_W) = (\mathbf{1}_U \cdot \mathbf{1}_V, \mathbf{1}_W)$. Then the right and left $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ -actions are adjoint with respect to $(-, -)$. \triangleleft

Remarks.

- (1) Generalizing the previous observation, it is proved in [45, Proposition 5.6.10] that for a relative 2-Segal simplicial groupoid $Y_\bullet \rightarrow X_\bullet$, the right and left $\mathcal{H}(X_\bullet)$ -module structures of $\mathcal{M}(Y_\bullet)$ are related via a categorical Green bilinear form.
- (2) The universal Hall algebra $\mathcal{H}(X_\bullet)$ is itself a $\mathcal{H}(X_\bullet)$ -bimodule which, however, does not arise from a relative 2-Segal space over X_\bullet .

Example. Let $\mathcal{S}_\bullet^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}) \rightarrow \mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}})$ be the relative 2-Segal groupoid associated to a stability function Z and a framing Φ on an abelian category \mathcal{C} . The universal left $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}_\phi^{Z\text{-ss}}))$ -module $\mathcal{M}(\mathcal{S}_\bullet^{\text{st-fr}}(\mathcal{C}_\phi^{Z\text{-ss}}))$ categorifies the stable framed Hall algebra representations of [41], [11] which appear in framed Donaldson-Thomas theory. \triangleleft

Example. Let $\mathcal{R}_\bullet(\mathcal{C}) \rightarrow \mathcal{S}_\bullet(\mathcal{C})$ be the relative 2-Segal groupoid associated to a proto-exact category with duality via the \mathcal{R}_\bullet -construction. For finitary exact \mathcal{C} , the universal left $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}))$ -module $\mathcal{M}(\mathcal{R}_\bullet(\mathcal{C}))$ categorifies the Hall algebra representations of [47]. Explicitly, $\mathcal{M}(\mathcal{R}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ is the k -vector space with basis $\{\mathbf{1}_{(M, \psi_M)}\}_{(M, \psi_M) \in \pi_0(\mathcal{R}_0(\mathcal{C}))}$ and left $\mathcal{H}(\mathcal{S}_\bullet(\mathcal{C}); \mathfrak{F}_0)$ -module structure

$$\mathbf{1}_U \star \mathbf{1}_{(M, \psi_M)} = \sum_{(N, \psi_N) \in \pi_0(\mathcal{R}_0(\mathcal{C}))} G_{U, M}^N \mathbf{1}_{(N, \psi_N)}.$$

The structure constant $G_{U, M}^N$ is the number of isotropic subobjects of N which are isomorphic to U and have reduction isometric to M .

Similarly, the simplicial stack version of $\mathcal{R}_\bullet(\mathcal{C})$ is relative 2-Segal over $\mathcal{S}_\bullet(\mathcal{C})$ and recovers the perverse sheaf theoretic [10], motivic and cohomological [48] Hall algebra representations which appear in the representation theory of quantum enveloping algebras and orientifold Donaldson-Thomas theory. \triangleleft

4.2. Hall monoidal module categories. We briefly describe a variant of the constructions of Section 4.1. Let k be a field of characteristic zero. Let X_\bullet be a unital 2-Segal groupoid² which is admissible with respect to weakly proper and locally proper maps. Write $\text{Fun}_0(X_1)$ for the (abelian) category of finitely supported functors $X_1 \rightarrow \text{Vect}_k$. The span m_X from Theorem 4.1 induces a bifunctor

$$\otimes^X = \partial_{1*} \circ (\partial_2 \times \partial_0)^* : \text{Fun}_0(X_1) \otimes \text{Fun}_0(X_1) \rightarrow \text{Fun}_0(X_1).$$

It is proved in [6, Theorem 2.49] that the triple $\mathcal{H}^\otimes(X_\bullet) = (\text{Fun}_0(X_1), \otimes^X, I_X)$ is a monoidal category. More generally, each componentwise cocontinuous monoidal left derivator of groupoids \mathbb{D} determines a monoidal category structure on $\mathbb{D}(X_1)$; see [45, Theorem 5.0.1.(1)]. The example $\mathcal{H}^\otimes(X_\bullet)$ arises for $\mathbb{D} = \text{Fun}_0(-, \text{Vect}_k)$.

In a similar way, we have the following relative construction. The same statement is proved in [45, Proposition 5.2.6.(2)], as a special case of [45, Theorem 5.0.1.(2)].

Theorem 4.3. *Assume that $F_\bullet : Y_\bullet \rightarrow X_\bullet$ is an admissible unital relative 2-Segal groupoid. Then the left action span $\mu_Y^!$ defines a bifunctor*

$$\otimes^Y = \partial_{1*} \circ (F_1 \times \partial_0)^* : \text{Fun}_0(X_1) \otimes \text{Fun}_0(Y_0) \rightarrow \text{Fun}_0(Y_0)$$

which gives $\mathcal{M}^\otimes(Y_\bullet) = (\text{Fun}_0(Y_0), \otimes^Y)$ the structure of a left $\mathcal{H}^\otimes(X_\bullet)$ -module category in the sense of [28].

Proof. The proof is very similar to that Theorem 4.2 and is therefore omitted. \square

We give two related instances of Theorem 4.3.

Example. Let $\text{Vect}_{\mathbb{F}_1}$ be the proto-exact category of finite dimensional vector spaces over \mathbb{F}_1 . Objects of $\text{Vect}_{\mathbb{F}_1}$ are finite pointed sets and morphisms are partial bijections. Let $\text{Vect}_{\mathbb{F}_1}^{\text{sk}} \subset \text{Vect}_{\mathbb{F}_1}$ be the skeleton of standard ordinals. Writing \mathfrak{S}_n for the symmetric group on n letters, we have an equivalence of groupoids

$$\mathcal{S}_1(\text{Vect}_{\mathbb{F}_1}^{\text{sk}}) \simeq \bigsqcup_{n \geq 0} B\mathfrak{S}_n.$$

Objects of $\mathcal{H}^\otimes(\mathcal{S}_\bullet(\text{Vect}_{\mathbb{F}_1}^{\text{sk}}))$ (henceforth denoted by $\mathcal{H}^\otimes(\text{Vect}_{\mathbb{F}_1}^{\text{sk}})$) are thus sequences of finite dimensional representations of symmetric groups over k , only finitely many of which are non-trivial. The monoidal product is induction of representations. Using the results of [27] it follows that $\mathcal{H}^\otimes(\text{Vect}_{\mathbb{F}_1}^{\text{sk}})$ is equivalent to the category \mathbf{P} of polynomial functors $\text{Vect}_k \rightarrow \text{Vect}_k$ [6, §2.5.1].

The functor $P = \text{Hom}_{\text{Vect}_{\mathbb{F}_1}}(-, \{*, 1\})$ defines a strict exact duality structure on $\text{Vect}_{\mathbb{F}_1}$. Note that each object of $\text{Vect}_{\mathbb{F}_1}$ is canonically isomorphic to its dual. In

²With minor modifications X_\bullet could be a simplicial object of a combinatorial model category.

particular, P preserves $\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}$. A symmetric form on \mathbb{F}_1^n is an element $\pi \in \mathfrak{S}_n$ which squares to the identity; conjugate such elements determine isometric symmetric forms. It follows that symmetric forms on \mathbb{F}_1^n are determined uniquely by their Witt index $0 \leq w \leq \lfloor \frac{n}{2} \rfloor$, the number of 2-cycles of any representative π . The isometry group of π , which is its centralizer in \mathfrak{S}_n , is isomorphic to $(\mathbb{Z}_2 \wr \mathfrak{S}_w) \times \mathfrak{S}_{n-2w}$. We therefore obtain an equivalence of groupoids

$$\mathcal{R}_0(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}) \simeq \bigsqcup_{w \geq 0} \bigsqcup_{d \geq 0} B((\mathbb{Z}_2 \wr \mathfrak{S}_w) \times \mathfrak{S}_d)$$

and we see that objects of $\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}})$ are finite sequences of representations of groups of the form $(\mathbb{Z}_2 \wr \mathfrak{S}_w) \times \mathfrak{S}_d$. The left $\mathcal{H}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}})$ -action on $\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}})$ is induction of representations along subgroups of the form

$$\mathfrak{S}_n \times ((\mathbb{Z}_2 \wr \mathfrak{S}_w) \times \mathfrak{S}_d) \leq (\mathbb{Z}_2 \wr \mathfrak{S}_{n+w}) \times \mathfrak{S}_d.$$

From this description we see that

$$\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}) = \bigoplus_{d=0}^{\infty} \mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; d)$$

as $\mathcal{H}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}})$ -modules, the index d labelling the fixed difference between the dimension and twice the Witt index. Moreover, we have

$$\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; d) \simeq \mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; 0) \times \mathbf{P}^d$$

where $\mathbf{P}^d \subset \mathbf{P}$ is the full subcategory of degree d homogeneous polynomial functors. Using again results of [27] we find that $\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; 0)$ is equivalent to $\mathbf{P}_{\mathbb{Z}_2}$, the category of polynomial functors $\mathbf{Vect}_k^{\mathbb{Z}_2\text{-gr}} \rightarrow \mathbf{Vect}_k$, where $\mathbf{Vect}_k^{\mathbb{Z}_2\text{-gr}}$ denotes the category of \mathbb{Z}_2 -graded finite dimensional vector spaces over k and we view $\mathbf{P}_{\mathbb{Z}_2}$ as a \mathbf{P} -module category via the forgetful functor $\mathbf{Vect}_k^{\mathbb{Z}_2\text{-gr}} \rightarrow \mathbf{Vect}_k$.

Upon passing to Grothendieck groups we obtain an isomorphism

$$K_0(\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; d)) \simeq \left(\bigoplus_{w=0}^{\infty} R_k(\mathbb{Z}_2 \wr \mathfrak{S}_w) \right) \otimes_{\mathbb{Z}} R_k(\mathfrak{S}_d)$$

as modules over the algebra

$$K_0(\mathcal{H}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}})) \simeq \bigoplus_{n=0}^{\infty} R_k(\mathfrak{S}_n),$$

where $R_k(-)$ denotes the representation ring over k . In the case $k = \mathbb{C}$ these modules have been studied in [40] and are closely related to the work of Zelevinsky [49], who studied the algebra structure on $K_0(\mathcal{M}^\otimes(\mathbf{Vect}_{\mathbb{F}_1}^{\text{sk}}; 0))$ arising from induction of representations of wreath symmetric products. \triangleleft

We now give a sort of quantization of the previous example.

Example. Let q be a prime power and consider the exact category $\mathbf{Vect}_{\mathbb{F}_q}$. Let $\mathbf{Vect}_{\mathbb{F}_q}^{\text{sk}} \subset \mathbf{Vect}_{\mathbb{F}_q}$ be the skeleton consisting of the vector spaces \mathbb{F}_q^n , $n \geq 0$. We have an equivalence of groupoids

$$S_1(\mathbf{Vect}_{\mathbb{F}_q}^{\text{sk}}) \simeq \bigsqcup_{n \geq 0} B\text{GL}_n(\mathbb{F}_q).$$

The category $\mathcal{H}^\otimes(\mathbf{Vect}_{\mathbb{F}_q}^{\text{sk}})$, whose monoidal product is parabolic induction of finite dimensional representations of GL , has appeared in the work of Joyal and Street [18]. When $k = \mathbb{C}$ the associated algebra $K_0(\mathcal{H}^\otimes(\mathbf{Vect}_{\mathbb{F}_q}^{\text{sk}}))$, which is the complex representation ring of the tower of general linear groups over \mathbb{F}_q , has been studied by Green [13] and Zelevinsky [49].

Assume now that q is odd and fix a sign $s \in \{\pm 1\}$. Take the exact duality structure with $P = \text{Hom}_{\text{Vect}_{\mathbb{F}_q}}(-, \mathbb{F}_q)$ with $\Theta = s \cdot \text{can}$. Symmetric forms in $\text{Vect}_{\mathbb{F}_q}$ are orthogonal or symplectic vector spaces. Identify the dual of \mathbb{F}_q^n with itself via the dual basis. It follows that we have an equivalence of groupoids

$$R_0(\text{Vect}_{\mathbb{F}_q}^{\text{sk}}) \simeq \bigsqcup_{n \geq 0} \bigsqcup_{\varepsilon \in W_n} BG_n^\varepsilon$$

where W_n is the Witt group of \mathbb{F}_q^n (\mathbb{Z}_2 if $s = 1$ and trivial if $s = -1$) and $G_n^\varepsilon = O_n^\varepsilon(\mathbb{F}_q)$ if $s = +1$ and $G_n = \text{Sp}_{2n}(\mathbb{F}_q)$ if $s = -1$. The left $\mathcal{H}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}})$ -action on $\mathcal{M}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}})$ is given by parabolic induction between GL and G representations. When $k = \mathbb{C}$ the $K_0(\mathcal{H}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}}))$ -modules $K_0(\mathcal{M}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}}))$ were studied by van Leeuwen [44], who showed that they are generated by cuspidal elements with respect to the natural Hall comodule structure.

It is natural to regard the module $\mathcal{M}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}})$ as a sort of q -analogue of the modules $\mathcal{M}^\otimes(\text{Vect}_{\mathbb{F}_1}^{\text{sk}}; d)$, $d = 0, 1$, consistent with the philosophy that the $q \rightarrow 1$ limit of $\text{G}(\mathbb{F}_q)$ is its Weyl group, namely $\mathbb{Z}_2 \wr \mathfrak{S}_n$ for the symplectic Sp_{2n} and even orthogonal O_{2n} groups and $(\mathbb{Z}_2 \wr \mathfrak{S}_n) \times \mathbb{Z}_2$ for the odd orthogonal group O_{2n+1} . The extra factor of \mathbb{Z}_2 for O_{2n+1} reflects the fact that the subcategories of $\mathcal{M}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}})$ with fixed Witt type ε are isomorphic as left $\mathcal{H}^\otimes(\text{Vect}_{\mathbb{F}_q}^{\text{sk}})$ -modules, the same statement holding also at $q = 1$. Working instead with Ringel-style Hall algebras and modules allows for the following more precise statement, which can be verified directly: the $q = 1$ specialization of the $\mathcal{H}(\text{Vect}_{\mathbb{F}_q}^{\text{sk}}; \mathfrak{F}_0)$ -module $\mathcal{M}(\text{Vect}_{\mathbb{F}_q}^{\text{sk}}; \mathfrak{F}_0)$ is isomorphic to the direct sum of the $\mathcal{H}(\text{Vect}_{\mathbb{F}_1}^{\text{sk}}; \mathfrak{F}_0)$ -modules $\mathcal{M}(\text{Vect}_{\mathbb{F}_1}^{\text{sk}}; \mathfrak{F}_0)$, $d = 0, 1$. \triangleleft

4.3. Modules over multivalued categories. In this section and the next we describe two higher categorical interpretations of relative 2-Segal simplicial sets; the first is in terms of modules over multivalued categories while the second is of a Hall algebraic nature.

Let $2\text{-SegS} \subset \mathbb{S}$ be the full subcategory of unital 2-Segal simplicial sets. Let also μCat denote the category of small multivalued categories. Objects of μCat are tuples $\mathfrak{X} = (\mathfrak{X}_0, \mathfrak{X}_1, m_{\mathfrak{X}}, a_{\mathfrak{X}}, e_{\mathfrak{X}}, i_{\mathfrak{X}}^l, i_{\mathfrak{X}}^r)$ consisting of sets \mathfrak{X}_0 and \mathfrak{X}_1 with source and target maps $\partial_1, \partial_0 : \mathfrak{X}_1 \rightarrow \mathfrak{X}_0$, a composition law $m_{\mathfrak{X}} \in \text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1, \mathfrak{X}_1)$, an associator isomorphism

$$(m_{\mathfrak{X}} \circ (m_{\mathfrak{X}} \times \mathbf{1}_{\mathfrak{X}_1}) \xrightarrow{a_{\mathfrak{X}}} m_{\mathfrak{X}} \circ (\mathbf{1}_{\mathfrak{X}_1} \times m_{\mathfrak{X}})) \in \text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1, \mathfrak{X}_1)$$

which satisfies Mac Lane coherence, a unit map $e_{\mathfrak{X}} : \mathfrak{X}_0 \rightarrow \mathfrak{X}_1$ and compatible left and right unit isomorphisms $i_{\mathfrak{X}}^l, i_{\mathfrak{X}}^r \in \text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1, \mathfrak{X}_1)$. A morphism $\varphi : \mathfrak{X} \rightarrow \mathfrak{X}'$ is the data of maps $\varphi_i : \mathfrak{X}_i \rightarrow \mathfrak{X}'_i$, $i = 0, 1$, compatible with the source, target and unit maps and a morphism of spans $\tilde{\varphi}_2 : \varphi_1 \circ m_{\mathfrak{X}} \rightarrow m_{\mathfrak{X}'} \circ (\varphi_1 \times_{\varphi_0} \varphi_1)$.

Theorem 4.4 ([7, Theorem 3.3.6]). *The generalized nerve construction defines an equivalence of categories $\mathbf{N}_\bullet : \mu\text{Cat} \xrightarrow{\sim} 2\text{-SegS}$.*

In the relative setting we will require the notion of a module over a multivalued category.

Definition. *Let \mathfrak{X} be a multivalued category. A unital left \mathfrak{X} -module is a tuple $\mathfrak{Y} = (\mathfrak{Y}_0, \mathfrak{F}_0, \mu_{\mathfrak{Y}}, \alpha_{\mathfrak{Y}}, \iota_{\mathfrak{Y}})$ consisting of*

- (i) *a set \mathfrak{Y}_0 together with a map $\mathfrak{F}_0 : \mathfrak{Y}_0 \rightarrow \mathfrak{X}_0$,*
- (ii) *a left action span $\mu_{\mathfrak{Y}} \in \text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0, \mathfrak{Y}_0)$,*
- (iii) *a module associator isomorphism*

$$(\mu_{\mathfrak{Y}} \circ (m_{\mathfrak{X}} \times \mathbf{1}_{\mathfrak{Y}_0}) \xrightarrow{\alpha_{\mathfrak{Y}}} \mu_{\mathfrak{Y}} \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mu_{\mathfrak{Y}})) \in \text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0, \mathfrak{Y}_0)$$

which satisfies module theoretic Mac Lane coherence: the diagram

$$\begin{array}{ccc}
& \mu \circ (m \circ (m \times \mathbf{1}_{\mathfrak{X}_1}) \times \mathbf{1}_{\mathfrak{Y}_0}) & \\
\alpha \circ (\mu \times \mathbf{1}_{\mathfrak{X}_1 \times \mathbf{1}_{\mathfrak{Y}_0}}) \swarrow & & \searrow \mu \circ (a \times \mathbf{1}_{\mathfrak{Y}_0}) \\
\mu \circ (m \times \mu) & & \mu \circ ((m \circ (\mathbf{1}_{\mathfrak{X}_1} \times m)) \times \mathbf{1}_{\mathfrak{Y}_0}) \\
\alpha \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mathbf{1}_{\mathfrak{X}_1} \times \mu) \downarrow & & \downarrow \alpha \circ (\mathbf{1}_{\mathfrak{X}_1} \times m \times \mathbf{1}_{\mathfrak{Y}_0}) \\
\mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mu)) & \xleftarrow{\mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times \alpha)} & \mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times (\mu \circ (m \circ \mathbf{1}_{\mathfrak{Y}_0})))
\end{array} \tag{23}$$

commutes in $\text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0, \mathfrak{Y}_0)$, and
(iv) a unit isomorphism $\iota_{\mathfrak{Y}} : \mu_{\mathfrak{Y}} \circ (e_{\mathfrak{X}} \times \mathbf{1}_{\mathfrak{Y}_0}) \rightarrow \mathbf{1}_{\mathfrak{Y}_0}$ in $\text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{Y}_0, \mathfrak{Y}_0)$
for which the diagram

$$\begin{array}{ccc}
\mu \circ (m \circ (\mathbf{1}_{\mathfrak{X}_1} \times e) \times \mathbf{1}_{\mathfrak{Y}_0}) & \xrightarrow{\alpha \circ (\mathbf{1}_{\mathfrak{X}_1} \times e \times \mathbf{1}_{\mathfrak{Y}_0})} & \mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mu \circ (e \times \mathbf{1}_{\mathfrak{Y}_0})) \\
\searrow \mu \circ (i_{\mathfrak{X}}^1 \times \mathbf{1}_{\mathfrak{Y}_0}) & & \swarrow \mu \circ (\mathbf{1}_{\mathfrak{X}_1} \times \iota_{\mathfrak{Y}}) \\
& \mu &
\end{array} \tag{24}$$

commutes in $\text{Hom}_{\text{Span}(\text{Set})}(\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0, \mathfrak{Y}_0)$.

Define a category $\mu\text{Cat-mod}$ as follows. Objects are unital left modules over multivalued categories. A morphism $(\varphi, \vartheta) : (\mathfrak{Y} \rightarrow \mathfrak{X}) \rightarrow (\mathfrak{Y}' \rightarrow \mathfrak{X}')$ consists of a morphism $\varphi : \mathfrak{X} \rightarrow \mathfrak{X}'$, a map $\vartheta_0 : \mathfrak{Y}_0 \rightarrow \mathfrak{Y}'_0$ which satisfies $\mathfrak{F}'_0 \circ \vartheta_0 = \varphi_0 \circ \mathfrak{F}_0$ and a morphism of spans

$$\tilde{\vartheta}_1 : \vartheta_0 \circ \mu_{\mathfrak{Y}} \rightarrow \mu_{\mathfrak{Y}'} \circ (\varphi_1 \times_{\varphi_0} \vartheta_0).$$

Informally, ϑ defines a morphism of \mathfrak{X} -modules $\mathfrak{Y} \rightarrow \varphi^* \mathfrak{Y}'$.

Let $2\text{-SegRelS} \subset \mathbb{S}^{[1]}$ be the full subcategory of unital relative 2-Segal simplicial sets.

Theorem 4.5. *The equivalence $\mathbf{N}_{\bullet} : \mu\text{Cat} \xrightarrow{\sim} 2\text{-SegS}$ lifts to an equivalence $\mathbf{N}_{\bullet}^{\text{rel}} : \mu\text{Cat-mod} \xrightarrow{\sim} 2\text{-SegRelS}$.*

Proof. We begin by modifying the proof of [7, Theorem 3.3.6] so as to define a functor $2\text{-SegRelS} \rightarrow \mu\text{Cat-mod}$. Let $F_{\bullet} : Y_{\bullet} \rightarrow X_{\bullet}$ be a unital relative 2-Segal simplicial set. The multivalued category \mathfrak{X} associated to X_{\bullet} via Theorem 4.4 has $\mathfrak{X}_0 = X_0$ and $\mathfrak{X}_1 = X_1$ with the canonical maps $\partial_1, \partial_0 : \mathfrak{X}_1 \rightarrow \mathfrak{X}_0$ and composition span

$$m_{\mathfrak{X}} = \{\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \xleftarrow{(\partial_2, \partial_0)} X_2 \xrightarrow{\partial_1} \mathfrak{X}_1\}.$$

Turning to the relative data, let $\mathfrak{Y}_0 = Y_0$ and $\mathfrak{F}_0 = F_0$. Define an action span by

$$\mu_{\mathfrak{Y}} = \{\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0 \xleftarrow{(F_1, \partial_0)} Y_1 \xrightarrow{\partial_1} \mathfrak{Y}_0\}.$$

The morphisms of spans

$$\mu_{\mathfrak{Y}} \circ (m_{\mathfrak{X}} \times \mathbf{1}_{\mathfrak{Y}_0}) \leftarrow \{\mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{X}_1 \times_{\mathfrak{X}_0} \mathfrak{Y}_0 \leftarrow Y_2 \rightarrow \mathfrak{Y}_0\} \rightarrow \mu_{\mathfrak{Y}} \circ (\mathbf{1}_{\mathfrak{X}_1} \times \mu_{\mathfrak{Y}}),$$

each of which is constructed as in the proof of Theorem 4.2 and is an isomorphism by the relative 2-Segal conditions, combine to define the associator $\alpha_{\mathfrak{Y}}$. The unit isomorphism $\iota_{\mathfrak{Y}}$ is defined to be the inverse of the relative unit bijection $Y_{\{0\}} \rightarrow X_{\{0\}} \times_{X_{\{0,1\}}} Y_{\{0,1\}}$. Mac Lane coherence and unit compatibility are verified as in the proof of Theorem 4.2. Hence \mathfrak{Y} is a unital left \mathfrak{X} -module. At the level of morphisms the functor $2\text{-SegRelS} \rightarrow \mu\text{Cat-mod}$ is defined in the obvious way.

To describe a quasi-inverse $\mathbf{N}_{\bullet}^{\text{rel}} : \mu\text{Cat-mod} \rightarrow 2\text{-SegRelS}$, let $\mathfrak{X} \in \mu\text{Cat}$ with associated unital 2-Segal simplicial set $X_{\bullet} = \mathbf{N}_{\bullet}(\mathfrak{X})$. Given a unital left \mathfrak{X} -module \mathfrak{Y} , let $Y_0 = \mathfrak{Y}_0$ and let Y_1 be the middle set of the span $\mu_{\mathfrak{Y}}$. We have canonical maps

$\partial_i : Y_1 \rightarrow Y_0$ and $F_i : Y_i \rightarrow X_i$, $i = 0, 1$, and, by inverting the map which defines $\iota_{\mathfrak{Y}}$, a map $s_0 : Y_0 \rightarrow Y_1$. Moreover, these maps obey the 1-truncated simplicial identities. For each $n \geq 2$ define

$$\tilde{Y}_n = \varprojlim^{\text{Set}}_{\{\sigma: \Delta^k \hookrightarrow \Delta^{P_n^\infty}\}_{k \leq 2}} F_{P_n^\infty}(\sigma),$$

where we use the notation introduced in Section 2.3. Given $0 \leq i < j < k \leq n$ and an element $\tilde{y} \in \tilde{Y}_n$, write y_{ij} and y_{ijk} for the corresponding elements of $Y_{\{i,j\}}$ and $X_{\{i,j,k\}}$, respectively. The associator $\alpha_{\mathfrak{Y}}$ defines a collection of bijections

$$\alpha_{ijk} : X_{\{i,j,k\}} \times_{X_{\{i,k\}}} Y_{\{i,k\}} \rightarrow Y_{\{i,j\}} \times_{Y_{\{j\}}} Y_{\{j,k\}}$$

which we use to define

$$Y_n = \{\tilde{y} \in \tilde{Y}_n \mid \alpha_{ijk}(y_{ijk}, y_{ik}) = (y_{ij}, y_{jk}) \text{ for all } 0 \leq i < j < k \leq n\}.$$

Module theoretic Mac Lane coherence implies that the Y_n assemble to a 1-Segal simplicial set Y_\bullet and that the canonical maps $F_n : Y_n \rightarrow X_n$ assemble to a simplicial morphism F_\bullet . To see that F_\bullet satisfies the remaining relative 2-Segal conditions, consider the commutative diagram

$$\begin{array}{ccc} Y_n & \xrightarrow{\quad\quad\quad} & X_n \times_{X_{\{0,n\}}} Y_{\{0,n\}} \\ \downarrow & & \downarrow \\ Y_1 \times_{Y_0} \cdots \times_{Y_0} Y_1 & \xrightarrow{\quad\quad\quad} & X_{\{0,1,2\}} \times_{X_{\{0,2\}}} \cdots \times_{X_{\{0,n-1\}}} X_{\{0,n-1,n\}} \times_{X_{\{0,n\}}} Y_{\{0,n\}}. \end{array}$$

The vertical morphisms are bijections by the 1- and 2-Segal conditions on Y_\bullet and X_\bullet , respectively. The bottom arrow is the iterated application of the module associator, relating the bracketings in which n elements of \mathfrak{X} act on \mathfrak{Y} from right to left and from left to right, and is therefore a bijection. Hence F_\bullet is relative 2-Segal. The relative unit bijection $Y_{\{0\}} \rightarrow X_{\{0\}} \times_{X_{\{0,1\}}} Y_{\{0,1\}}$ is the inverse of $\iota_{\mathfrak{Y}}$. The compatibility of $\iota_{\mathfrak{Y}}$, $i_{\mathfrak{X}}^l$ and $\alpha_{\mathfrak{Y}}$ implies that the higher relative unit bijections hold. The functor $\mathbf{N}_\bullet^{\text{rel}}$ assigns to a morphism (φ, ϑ) the pair $(\phi_\bullet, \theta_\bullet)$, where $\phi_\bullet = \mathbf{N}_\bullet(\varphi)$, θ_0 is equal to $\tilde{\vartheta}_0$ and θ_n , $n \geq 1$, are the canonically induced maps, which are well-defined by the definition of Y_n and Y'_n , $n \geq 2$. \square

Remark. Theorem 4.5 could also have been formulated in terms of right modules.

There is a semi-simplicial variant of Theorem 4.5 where $\mu\text{Cat-mod}$ is replaced with $\frac{1}{2}\mu\text{Cat-mod}$, the category of left modules over multivalued semicategories, the prefix ‘semi’ indicating that all data related to unit morphisms is omitted. As a simple special case, let X_\bullet be a 2-Segal semi-simplicial set with $X_0 = X_1 = \text{pt}$. Then X_\bullet defines a distributive monoidal endofunctor on Set by $F \otimes F' = X_2 \times F \times F'$. As explained in [7, §3.7], the associator reduces to a bijection $a : X_2 \times X_2 \rightarrow X_2 \times X_2$ which satisfies the pentagon equation

$$a_{23} \circ a_{13} \circ a_{12} = a_{12} \circ a_{23}.$$

Conversely, a bijective solution a of the pentagon equation on a set X_2 extends to a 2-Segal semi-simplicial set with $\mathbf{N}_0(X_2, a) = \mathbf{N}_1(X_2, a) = \text{pt}$ and $\mathbf{N}_n(X_2, a)$, $n \geq 2$, the set of tuples $\{x_{ijk} \in X_2 \mid 0 \leq i < j < k \leq n\}$ which satisfy

$$a(x_{ijk}, x_{ikl}) = (x_{ijl}, x_{jkl}), \quad 0 \leq i < j < k < l \leq n.$$

Modifying this construction, a relative 2-Segal semi-simplicial set $Y_\bullet \rightarrow X_\bullet$ with $Y_0 = \text{pt}$ defines a monoidal endofunctor $F \boxtimes G = Y_1 \times F \times G$ which is a left \otimes -module. The module associator is a bijection $\alpha : X_2 \times Y_1 \rightarrow Y_1 \times Y_1$ which satisfies the a -pentagon equation

$$\alpha_{23} \circ \alpha_{13} \circ \alpha_{12} = \alpha_{12} \circ \alpha_{23}.$$

Moreover, from a bijective solution to the a -pentagon equation we can construct a relative 2-Segal semi-simplicial set with $\mathbf{N}_0(Y_1, \alpha) = \text{pt}$, $\mathbf{N}_1(Y_1, \alpha) = Y_1$ and $\mathbf{N}_n(Y_1, \alpha)$, $n \geq 2$, the subset of $\mathbf{N}_n(X_2, a) \times \{y_{ij} \in Y_1 \mid 0 \leq i < j \leq n\}$ consisting of tuples which satisfy

$$\alpha(x_{ijk}, y_{ik}) = (y_{ij}, y_{jk}), \quad 0 \leq i < j < k \leq n.$$

The structure map $\mathbf{N}_\bullet(Y_1, \alpha) \rightarrow \mathbf{N}_\bullet(X_2, a)$ is the canonical projection.

Example. For any group \mathbf{G} the map

$$a : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G} \times \mathbf{G}, \quad (x, y) \mapsto (xy, y)$$

is a bijective solution to the pentagon equation and so defines a 2-Segal semi-simplicial set $\mathbf{N}_\bullet(\mathbf{G})$; see [21], [7, Example 3.7.7].

Let now $\rho : \mathbf{G} \rightarrow \text{Aut}(M)$ be a left \mathbf{G} -action on a set M . Then

$$\alpha : \mathbf{G} \times M \rightarrow M \times M, \quad (x, m) \mapsto (\rho(x)m, m)$$

solves the a -pentagon equation. Moreover, α is a bijection if and only if ρ gives M the structure of a \mathbf{G} -torsor. Hence, associated to each \mathbf{G} -torsor M is a relative 2-Segal simplicial set $\mathbf{N}_\bullet(\mathbf{G}, M) \rightarrow \mathbf{N}_\bullet(\mathbf{G})$. When $M = \mathbf{G}$ with \mathbf{G} acting by left multiplication the above construction recovers the left path $P^{\triangleleft} \mathbf{N}_\bullet(\mathbf{G}) \rightarrow \mathbf{N}_\bullet(\mathbf{G})$. \triangleleft

4.4. Presheaves on Hall 2-categories. We begin by constructing a module over a Hall-type algebra from a relative 2-Segal semi-simplicial set. Fix a field k . Let X_\bullet be a 2-Segal simplicial set for which the map (∂_2, ∂_0) from the span (21) has finite fibres. The Hall category $H(X_\bullet)$ [7, §3.4] is the k -linear category with $\text{Ob}(H(X_\bullet)) = X_0$ and $\text{Hom}_{H(X_\bullet)}(a, b) = \mathfrak{F}_0(X_{a \rightarrow b})$, the finitely supported k -valued functions on the set

$$X_{a \rightarrow b} = \{a\} \times_{X_0} X_1 \times_{X_0} \{b\}.$$

Composition of morphisms is defined by push-pull along the span

$$X_{b \rightarrow c} \times X_{a \rightarrow b} \leftarrow \{p \in X_2 \mid \partial_{\{0\}}p = a, \partial_{\{1\}}p = b, \partial_{\{2\}}p = c\} \rightarrow X_{a \rightarrow c}. \quad (25)$$

Writing $\mathbf{1}_x$ for the characteristic function of $x \in X_{a \rightarrow b}$, composition in $H(X_\bullet)$ becomes $\mathbf{1}_x \cdot \mathbf{1}_{x'} = \sum_{x''} f_{x, x'}^{x''} \mathbf{1}_{x''}$ where

$$f_{x, x'}^{x''} = |\{p \in X_2 \mid \partial_2 p = x, \partial_1 p = x', \partial_0 p = x''\}|.$$

Similarly, a relative 2-Segal simplicial set $F_\bullet : Y_\bullet \rightarrow X_\bullet$ for which the map (F_1, ∂_0) from the span (22) has finite fibres defines a presheaf $\mathcal{F} : H(X_\bullet)^{\text{op}} \rightarrow \text{Set}$ by setting $\mathcal{F}(a) = \mathfrak{F}_0(F_0^{-1}(a))$, $a \in X_0$, and using push-pull along the span

$$X_{a \rightarrow b} \times F_0^{-1}(b) \leftarrow \{q \in Y_1 \mid \partial_1 q \in F_0^{-1}(a), \partial_0 q \in F_0^{-1}(b)\} \rightarrow F_0^{-1}(a) \quad (26)$$

to define the action map $\text{Hom}_{H(X_\bullet)}(a, b) \times \mathcal{F}(b) \rightarrow \mathcal{F}(a)$. Writing $\mathbf{1}_\xi \in \mathcal{F}(a)$ for the characteristic function of $\xi \in F_0^{-1}(a)$, we have $\mathbf{1}_x \star \mathbf{1}_{\xi'} = \sum_{\xi''} g_{x, \xi'}^{\xi''} \mathbf{1}_{\xi''}$ where

$$g_{x, \xi'}^{\xi''} = |\{q \in Y_1 \mid \partial_1 q = \xi'', \partial_0 q = \xi', F_1(q) = x\}|.$$

Before categorifying the above construction we recall some preliminary definitions from [7, §3.5.B]. A category \mathcal{C} is called \sqcup -semisimple if it is equivalent to an overcategory $\text{Set}/_B$, in which case a simple object of \mathcal{C} is an object which is isomorphic to a point $\{b\} \in \text{Set}/_B$. Denote by $\|\mathcal{C}\|$ the set of isomorphism classes of simple objects of \mathcal{C} . A functor between \sqcup -semisimple categories is called additive (resp. simple) if it preserves coproducts (resp. simple objects). The \sqcup -semisimple categories, their additive functors and their natural transformations form a bicategory Cat^{\sqcup} . Let $\text{Cat}^{\sqcup!} \subset \text{Cat}^{\sqcup}$ be the subcategory whose morphisms are instead simple additive functors.

Define a 2-functor by $A : \text{Span}(\text{Set}) \rightarrow \text{Cat}^{\sqcup}$ by

$$Z \mapsto \text{Set}/Z, \quad (Z \xleftarrow{s} W \xrightarrow{p} Z') \mapsto (\text{Set}/Z \xrightarrow{p_* \circ s^*} \text{Set}/Z')$$

with the obvious action on 2-morphisms. Define also $B : \text{Cat}^{\sqcup} \rightarrow \text{Span}(\text{Set})$ by

$$\mathcal{C} \mapsto \|\mathcal{C}\|, \quad (\mathcal{C} \xrightarrow{\phi} \mathcal{C}') \mapsto (\|\mathcal{C}\| \leftarrow \bigsqcup_{a \in \|\mathcal{C}\|} F'(\phi(a)) \rightarrow \|\mathcal{C}'\|)$$

where $F' : \mathcal{C}' \rightarrow \text{Set}/Z'$ is some equivalence.

Proposition 4.6 ([7, Proposition 3.5.4]). *The 2-functors $A : \text{Span}(\text{Set}) \leftrightarrow \text{Cat}^{\sqcup} : B$ are mutually inverse 2-equivalences which restrict to 2-equivalences $\text{Set} \leftrightarrow \text{Cat}^{\sqcup}$.*

For later purposes, fix natural equivalences

$$\kappa : \mathbf{1}_{\text{Cat}^{\sqcup}} \Rightarrow A \circ B, \quad \lambda : B \circ A \Rightarrow \mathbf{1}_{\text{Span}(\text{Set})}$$

realizing the 2-equivalences of Proposition 4.6.

A semibicategory is the data of a bicategory with all mention of unit 1-morphisms omitted. A semibicategory is called \sqcup -semisimple if it is small, its morphism categories are \sqcup -semisimple and composition of morphisms is additive in each variable. A lax 2-functor between \sqcup -semisimple semibicategories is called admissible if the associated functors on morphism categories are simple additive.

Definition. *Two admissible lax 2-functors $\phi, \phi' : \mathbb{X} \rightarrow \mathbb{X}'$ of \sqcup -semisimple semibicategories are called equivalent if they agree on objects and there exists a collection of natural isomorphisms*

$$\begin{array}{ccc} & \xrightarrow{\phi_{a,b}} & \\ \text{Hom}_{\mathbb{X}}(a,b) & \xrightarrow{U_{a,b}} \Downarrow & \text{Hom}_{\mathbb{X}'}(\phi(a), \phi(b)), \quad a, b \in \text{Ob}(\mathbb{X}) \\ & \xrightarrow{\phi'_{a,b}} & \end{array}$$

which commute with the coherence maps for ϕ, ϕ' and the associativity isomorphisms.

If \mathbb{X}, \mathbb{X}' are bicategories, then $\phi, \phi' : \mathbb{X} \rightarrow \mathbb{X}'$ are equivalent if there exists an invertible icon $U : \phi \Rightarrow \phi'$. Slightly abusively, we will refer to $\{U_{a,b}\}_{a,b \in \text{Ob}(\mathbb{X})}$ as an icon even if \mathbb{X}, \mathbb{X}' are not bicategories. The \sqcup -semisimple semibicategories and their equivalence classes of admissible lax 2-functors form a category $\frac{1}{2}\text{biCat}^{\sqcup}$.

Let X_{\bullet} be a 2-Segal semi-simplicial set. Its Hall 2-category $\mathbb{H}(X_{\bullet})$ is the \sqcup -semisimple semibicategory with

$$\text{Ob}(\mathbb{H}(X_{\bullet})) = X_0, \quad \text{Hom}_{\mathbb{H}(X_{\bullet})}(a,b) = \text{Set}/X_{a \rightarrow b}.$$

Push-pull along the span (25) defines composition of 1-morphisms. Associated to a morphism $\phi_{\bullet} : X_{\bullet} \rightarrow X'_{\bullet}$ is the admissible lax 2-functor $\phi : \mathbb{H}(X_{\bullet}) \rightarrow \mathbb{H}(X'_{\bullet})$ given by ϕ_0 on objects and by pushforward along ϕ_1 on morphisms.

Theorem 4.7 ([7, Theorem 3.5.8]). *The assignment $X_{\bullet} \mapsto \mathbb{H}(X_{\bullet})$ extends to an equivalence between the category of 2-Segal semi-simplicial sets and $\frac{1}{2}\text{biCat}^{\sqcup}$.*

Proof. A proof is outlined in [7]. We fill in some details which will be needed below. We first construct a functor $\Xi : \frac{1}{2}\text{biCat}^{\sqcup} \rightarrow \frac{1}{2}\mu\text{Cat}$. Define $\mathfrak{X} = \Xi(\mathbb{X}) \in \frac{1}{2}\mu\text{Cat}$ by

$$\mathfrak{X}_0 = \text{Ob}(\mathbb{X}), \quad \mathfrak{X}_1 = \bigsqcup_{a,b \in \text{Ob}(\mathbb{X})} \|\text{Hom}_{\mathbb{X}}(a,b)\|$$

with span $m_{\mathfrak{X}}$ obtained by applying Proposition 4.6 to the functors which define composition of 1-morphisms in \mathbb{X} . Given $\phi : \mathbb{X} \rightarrow \mathbb{X}'$, define $\varphi = \Xi(\phi)$ so that φ_0 is equal to ϕ on objects and φ_1 is given by the functors $\phi_{a,b}$. If $\phi, \phi' : \mathbb{X} \rightarrow \mathbb{X}'$

are equivalent, say via an icon $U : \phi \Rightarrow \phi'$, then $\varphi_0 = \varphi'_0$ and $\varphi_1 = \varphi'_1$ due to the isomorphisms $U_{a,b}(f) : \phi_{a,b}(f) \xrightarrow{\sim} \phi'_{a,b}(f)$, $f \in \text{Hom}_{\mathbb{X}}(a, b)$.

Define also a functor $\Omega : \frac{1}{2}\mu\text{Cat} \rightarrow \frac{1}{2}\text{biCat}^{\sqcup}$ by setting $\mathbb{X} = \Omega(\mathfrak{X})$, where $\text{Ob}(\mathbb{X}) = \mathfrak{X}_0$ and $\text{Hom}_{\mathbb{X}}(a, b) = \text{Set}/_{\mathfrak{X}_{a \rightarrow b}}$. Given $\varphi : \mathfrak{X} \rightarrow \mathfrak{X}'$, let $\phi = \Omega(\varphi)$ be the morphism which is equal to φ_0 on objects and whose component functor $\phi_{a,b}$ is pushforward along $\varphi_1 : \mathfrak{X}_{a \rightarrow b} \rightarrow \mathfrak{X}'_{\varphi_0(a) \rightarrow \varphi_0(b)}$.

We claim that Ξ and Ω are mutually inverse equivalences. Let $\epsilon : \Xi \circ \Omega \Rightarrow \mathbf{1}_{\frac{1}{2}\mu\text{Cat}}$ be the natural isomorphism whose component $\epsilon_{\mathfrak{X}} : \Xi(\Omega(\mathfrak{X})) \rightarrow \mathfrak{X}$ has $(\epsilon_{\mathfrak{X}})_0$ equal to the identity and has $(\epsilon_{\mathfrak{X}})_1$ equal to the map

$$\bigsqcup_{a,b \in \mathfrak{X}_0} \lambda_{\mathfrak{X}_{a \rightarrow b}} : \bigsqcup_{a,b \in \mathfrak{X}_0} \|\text{Set}/_{\mathfrak{X}_{a \rightarrow b}}\| \rightarrow \mathfrak{X}_1.$$

Similarly, let $\eta : \mathbf{1}_{\frac{1}{2}\text{biCat}^{\sqcup}} \Rightarrow \Omega \circ \Xi$ be the natural transformation whose component $\eta_{\mathbb{X}} : \mathbb{X} \rightarrow \Omega(\Xi(\mathbb{X}))$ is the identity on objects and is equal to the functor $\kappa_{\text{Hom}_{\mathbb{X}}(a,b)}$ on morphism categories. Given $\phi : \mathbb{X} \rightarrow \mathbb{X}'$, we need to check that the diagram

$$\begin{array}{ccc} \mathbb{X} & \xrightarrow{\eta_{\mathbb{X}}} & \Omega(\Xi(\mathbb{X})) \\ \phi \downarrow & & \downarrow \Omega(\Xi(\phi)) \\ \mathbb{X}' & \xrightarrow{\eta_{\mathbb{X}'}} & \Omega(\Xi(\mathbb{X}')) \end{array}$$

commutes in $\frac{1}{2}\text{biCat}^{\sqcup}$, which amounts to giving an invertible icon $U : \eta_{\mathbb{X}'} \circ \phi \Rightarrow \Omega(\Xi(\phi)) \circ \eta_{\mathbb{X}}$. The required natural transformation $U_{a,b}$ is defined to be

$$\begin{array}{ccc} \text{Hom}_{\mathbb{X}}(a, b) & \xrightarrow{\kappa_{\text{Hom}_{\mathbb{X}}(a,b)}} & \text{Set}/_{\|\text{Hom}_{\mathbb{X}}(a,b)\|} \\ \phi_{a,b} \downarrow & \searrow \kappa_{\phi_{a,b}} & \downarrow \|\phi_{a,b}\|_* \\ \text{Hom}_{\mathbb{X}'}(\phi(a), \phi(b)) & \xrightarrow{\kappa_{\text{Hom}_{\mathbb{X}'}(\phi(a), \phi(b))}} & \text{Set}/_{\|\text{Hom}_{\mathbb{X}'}(\phi(a), \phi(b))\|} \end{array}$$

This establishes an equivalence $\frac{1}{2}\text{biCat}^{\sqcup} \simeq \frac{1}{2}\mu\text{Cat}$. Using the semi-simplicial variant of Theorem 4.4 we get an equivalence from $\frac{1}{2}\text{biCat}^{\sqcup}$ to the category of 2-Segal semi-simplicial sets which is a quasi-inverse of the functor $X_{\bullet} \mapsto \mathbb{H}(X_{\bullet})$. \square

Motivated by Proposition 2.2, the goal of the remainder of this section is to interpret relative 2-Segal semi-simplicial sets in terms of a certain class of presheaves. To this end, define a Cat^{\sqcup} -valued presheaf on a \sqcup -semisimple semibicategory to be a 2-functor $\mathbb{F} : \mathbb{X}^{\text{op}} \rightarrow \text{Cat}^{\sqcup}$ with $\mathbb{X} \in \frac{1}{2}\text{biCat}^{\sqcup}$. Here \mathbb{X}^{op} is the semibicategory obtained from \mathbb{X} by reversing only its 1-cells. A morphism

$$(\phi, \theta) : (\mathbb{F} : \mathbb{X}^{\text{op}} \rightarrow \text{Cat}^{\sqcup}) \rightarrow (\mathbb{F}' : \mathbb{X}'^{\text{op}} \rightarrow \text{Cat}^{\sqcup})$$

consists of a lax 2-functor $\phi : \mathbb{X} \rightarrow \mathbb{X}'$ and an oplax natural transformation $\theta : \mathbb{F} \Rightarrow \mathbb{F}' \circ \phi^{\text{op}}$. The pair (ϕ, θ) is called admissible if ϕ is admissible and the functors $\theta_a : \mathbb{F}(a) \rightarrow \mathbb{F}'(\phi(a))$ are simple additive. The composition $\mathbb{F} \xrightarrow{(\phi, \theta)} \mathbb{F}' \xrightarrow{(\xi, \zeta)} \mathbb{F}''$ is defined by the right whiskering of ζ and ϕ^{op} : $(\xi, \zeta) \circ (\phi, \theta) = (\xi \circ \phi, (\zeta \phi^{\text{op}}) \circ \theta)$.

Definition. *Two admissible morphisms*

$$(\phi, \theta), (\phi', \theta') : (\mathbb{F} : \mathbb{X}^{\text{op}} \rightarrow \text{Cat}^{\sqcup}) \rightarrow (\mathbb{F}' : \mathbb{X}'^{\text{op}} \rightarrow \text{Cat}^{\sqcup})$$

are called *equivalent* if there exists an invertible icon $U : \phi \Rightarrow \phi'$ and an invertible modification

$$\begin{array}{ccc} & \mathbb{F} & \\ & \curvearrowright & \\ \mathbb{X}^{\text{op}} & \xrightarrow{(\mathbb{F}'U^{\text{op}})\circ\theta} & \text{Cat}^{\sqcup} \\ & \curvearrowleft & \\ & \mathbb{F}'\circ\phi'^{\text{op}} & \end{array}$$

(A diagram showing a span from \mathbb{X}^{op} to Cat^{\sqcup} . The top arrow is $(\mathbb{F}'U^{\text{op}})\circ\theta$ and the bottom arrow is $\mathbb{F}'\circ\phi'^{\text{op}}$. A curved arrow labeled \mathbb{F} connects the two arrows. A central square diagram shows Γ as a modification between θ and θ' .

Denote by Psh^{\sqcup} the category of Cat^{\sqcup} -valued presheaves on \sqcup -semisimple semibi-categories and their equivalence classes of admissible morphisms.

We now construct a functor from the category of relative 2-Segal semi-simplicial sets to Psh^{\sqcup} . Given $F_{\bullet} : Y_{\bullet} \rightarrow X_{\bullet}$, define $\mathbb{F} : \mathbb{H}(X_{\bullet})^{\text{op}} \rightarrow \text{Cat}^{\sqcup}$ as follows. Put $\mathbb{F}(a) = \text{Set}_{/F_0^{-1}(a)}$, $a \in X_0$, and use the span (26) to define a functor

$$\mathbb{F}_{a,b} : \text{Set}_{/X_{a \rightarrow b}} \rightarrow [\text{Set}_{/F_0^{-1}(b)}, \text{Set}_{/F_0^{-1}(a)}],$$

the image of which consists of additive functors by Proposition 4.6. Here and in what follows we use the equivalence $[\mathcal{C} \times \mathcal{D}, \mathcal{E}] \simeq [\mathcal{C}, [\mathcal{D}, \mathcal{E}]]$ for categories \mathcal{C}, \mathcal{D} and \mathcal{E} , the first two of which are small. The relative 2-Segal bijections induce the coherence isomorphisms for \mathbb{F} . Given a morphism

$$\begin{array}{ccc} Y_{\bullet} & \xrightarrow{\theta_{\bullet}} & Y'_{\bullet} \\ F_{\bullet} \downarrow & & \downarrow F'_{\bullet} \\ X_{\bullet} & \xrightarrow{\phi_{\bullet}} & X'_{\bullet} \end{array}$$

of relative 2-Segal semi-simplicial sets, we need to define a morphism

$$(\phi, \theta) : (\mathbb{F} : \mathbb{H}(X_{\bullet})^{\text{op}} \rightarrow \text{Cat}^{\sqcup}) \rightarrow (\mathbb{F}' : \mathbb{H}(X'_{\bullet})^{\text{op}} \rightarrow \text{Cat}^{\sqcup}).$$

Let ϕ be the lax 2-functor associated to ϕ_{\bullet} . Let $\theta_a : \mathbb{F}(a) \rightarrow \mathbb{F}'(\phi(a))$ be the pushforward along $\theta_0|_{F_0^{-1}(a)} : F_0^{-1}(a) \rightarrow F_0'^{-1}(\phi(a))$. The natural transformation

$$\begin{array}{ccc} \text{Set}_{/X_{a \rightarrow b}} & \xrightarrow{(\mathbb{F}'\circ\phi^{\text{op}})_{a,b}} & [\text{Set}_{/F_0'^{-1}(\phi(b))}, \text{Set}_{/F_0'^{-1}(\phi(a))}] \\ \mathbb{F}_{a,b} \downarrow & \nearrow \theta_{a,b} & \downarrow -\circ\theta_b \\ [\text{Set}_{/F_0^{-1}(b)}, \text{Set}_{/F_0^{-1}(a)}] & \xrightarrow{\theta_a \circ -} & [\text{Set}_{/F_0^{-1}(b)}, \text{Set}_{/F_0^{-1}(\phi(a))}] \end{array}$$

is induced by the morphism of spans determined by the restriction of the map $\theta_1 : Y_1 \rightarrow Y'_1$ to the middle set of the span (26).

Theorem 4.8. *The assignment $F_{\bullet} \mapsto \mathbb{F}$ extends to an equivalence between the category of relative 2-Segal semi-simplicial sets and Psh^{\sqcup} .*

Proof. Lift $\Xi : \frac{1}{2}\text{biCat}^{\sqcup} \rightarrow \frac{1}{2}\mu\text{Cat}$ to a functor $\Xi^{\text{rel}} : \text{Psh}^{\sqcup} \rightarrow \frac{1}{2}\mu\text{Cat-mod}$ as follows. Define $\Xi^{\text{rel}}(\mathbb{F} : \mathbb{X}^{\text{op}} \rightarrow \text{Cat}^{\sqcup}) = (\mathfrak{F} : \mathfrak{Y} \rightarrow \mathfrak{X})$ by $\mathfrak{X} = \Xi(\mathbb{X})$ and $\mathfrak{Y}_0 = \bigsqcup_{a \in \text{Ob}(\mathbb{X})} \|\mathbb{F}(a)\|$ with the canonical map $\mathfrak{F}_0 : \mathfrak{Y}_0 \rightarrow \mathfrak{X}_0$. The span $\mu_{\mathfrak{Y}}$ is obtained by applying Proposition 4.6 to the functors $\mathbb{F}_{a,b}$ while the associator $\alpha_{\mathfrak{Y}}$ is defined using the coherence isomorphisms for \mathbb{F} . Given a morphism $(\phi, \theta) : \mathbb{F} \rightarrow \mathbb{F}'$, set $\varphi = \Xi(\phi)$, let $\vartheta_0 : \mathfrak{Y}_0 \rightarrow \mathfrak{Y}'_0$ be the map determined by the functors θ_a and let $\vartheta_1 : \vartheta_0 \circ \mu_{\mathfrak{Y}} \rightarrow \mu_{\mathfrak{Y}'} \circ (\varphi_1 \times_{\varphi_0} \vartheta_0)$ be the morphism obtained by applying Proposition

4.6 to the coherence natural transformations

$$\begin{array}{ccc}
\mathrm{Hom}_{\mathbb{X}}(a, b) \times \mathbb{F}(b) & \xrightarrow{\phi_{a,b} \times \theta_b} & \mathrm{Hom}_{\mathbb{X}'}(\phi(a), \phi(b)) \times \mathbb{F}'(\phi(b)) \\
\mathbb{F}_{a,b} \downarrow & \nearrow \theta_{a,b} & \downarrow \mathbb{F}'_{a,b} \\
\mathbb{F}(a) & \xrightarrow{\theta_a} & \mathbb{F}'(\phi(a)).
\end{array}$$

Suppose that (ϕ, θ) and (ϕ', θ') are equivalent morphisms, say via an icon U and a modification Γ . Then $\varphi = \varphi'$. Since $U_a : \phi(a) \rightarrow \phi'(a)$ is the identity we have $((\mathbb{F}'U^{\mathrm{op}}) \circ \theta)_a = \theta_a$ and hence a natural isomorphism $\Gamma_a : \theta_a \Rightarrow \theta'_a$, showing that $\vartheta_0 = \vartheta'_0$. The equality $\tilde{\vartheta}_1 = \tilde{\vartheta}'_1$ follows from the fact that the components $\{\Gamma_a\}_{a \in \mathrm{Ob}(\mathbb{X})}$ commute with morphisms in \mathbb{X} .

Similarly, construct a lift $\Omega^{\mathrm{rel}} : \frac{1}{2}\mu\mathrm{Cat}\text{-mod} \rightarrow \mathrm{Psh}^{\sqcup}$ of $\Omega : \frac{1}{2}\mu\mathrm{Cat} \rightarrow \frac{1}{2}\mathrm{biCat}^{\sqcup}$. Define $\Omega^{\mathrm{rel}}(\mathfrak{F} : \mathfrak{Y} \rightarrow \mathfrak{X}) = (\mathbb{F} : \mathbb{X}^{\mathrm{op}} \rightarrow \mathrm{Cat}^{\sqcup})$ by $\mathbb{X} = \Omega(\mathfrak{X})$ with $\mathbb{F}(a) = \mathrm{Set}_{/\mathfrak{F}_0^{-1}(a)}$ and functors $\mathbb{F}_{a,b}$ determined by $\mu\mathfrak{Y}$. Given a morphism $(\varphi, \vartheta) : \mathfrak{F} \rightarrow \mathfrak{F}'$, put $\phi = \Omega(\varphi)$ and let θ be the oplax natural transformation whose component functor $\theta_a : \mathbb{F}(a) \rightarrow \mathbb{F}'(\phi(a))$ is pushforward along the map $\vartheta_{0|\mathfrak{F}_0^{-1}(a)} : \mathfrak{F}_0^{-1}(a) \rightarrow \mathfrak{F}'_0^{-1}(\phi(a))$ and whose coherence natural transformations $\theta_{a,b}$ are induced by the morphism $\tilde{\vartheta}_1$.

To prove that Ξ^{rel} and Ω^{rel} are inverse equivalences, we first lift $\epsilon : \Xi \circ \Omega \Rightarrow \mathbf{1}_{\frac{1}{2}\mu\mathrm{Cat}}$ to $\epsilon^{\mathrm{rel}} : \Xi^{\mathrm{rel}} \circ \Omega^{\mathrm{rel}} \Rightarrow \mathbf{1}_{\frac{1}{2}\mu\mathrm{Cat}\text{-mod}}$. On the total space $\epsilon_{\mathfrak{F}}^{\mathrm{rel}} : \Xi^{\mathrm{rel}}(\Omega^{\mathrm{rel}}(\mathfrak{F})) \rightarrow \mathfrak{F}$ is given by $\bigsqcup_{a \in \mathfrak{X}_0} \lambda_{\mathfrak{F}_0^{-1}(a)}$ on objects while the morphism $\vartheta_0 \circ \mu_{\Xi^{\mathrm{rel}}(\Omega^{\mathrm{rel}}(\mathfrak{Y}))} \rightarrow \mu_{\mathfrak{Y}} \circ (\varphi_1 \times \varphi_0 \vartheta_0)$ is $\lambda_{\mu_{\mathfrak{Y}}}$. Similarly, lift $\eta : \mathbf{1}_{\frac{1}{2}\mathrm{biCat}^{\sqcup}} \Rightarrow \Omega \circ \Xi$ to $\eta^{\mathrm{rel}} : \mathbf{1}_{\mathrm{Psh}^{\sqcup}} \Rightarrow \Omega^{\mathrm{rel}} \circ \Xi^{\mathrm{rel}}$ as follows. Writing $\eta_{\mathbb{F}}^{\mathrm{rel}} = (\phi_{\mathbb{F}}, \theta_{\mathbb{F}})$, put $\phi_{\mathbb{F}} = \eta_{\mathbb{X}}$. Since $\eta_{\mathbb{X}}$ is the identity on objects, we can define $\theta_{\mathbb{F}} : \mathbb{F} \rightarrow \Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\mathbb{F})) \circ \eta_{\mathbb{X}}^{\mathrm{op}}$ so that its component functor $(\theta_{\mathbb{F}})_a$ is $\lambda_{\mathbb{F}(a)}$. Associated to the functor $\mathbb{F}_{a,b}$ is the natural transformation

$$\begin{array}{ccc}
\mathrm{Hom}_{\mathbb{X}}(a, b) \times \mathbb{F}(b) & \xrightarrow{\lambda_{\mathrm{Hom}_{\mathbb{X}}(a,b) \times \mathbb{F}(b)}} & \mathrm{Set}_{/\|\mathrm{Hom}_{\mathbb{X}}(a,b) \times \mathbb{F}(b)\|} \\
\mathbb{F}_{a,b} \downarrow & \nearrow \lambda_{\mathbb{F}_{a,b}} & \downarrow \|\mathbb{F}_{a,b}\|_* \\
\mathbb{F}(a) & \xrightarrow{\lambda_{\mathbb{F}(a)}} & \mathrm{Set}_{/\|\mathbb{F}(a)\|}
\end{array}$$

which we take to be the coherence data $(\theta_{\mathbb{F}})_{a,b}$. It remains to check that, for an admissible morphism $(\phi, \theta) : \mathbb{F} \rightarrow \mathbb{F}'$, the diagram

$$\begin{array}{ccc}
\mathbb{F} & \xrightarrow{\eta_{\mathbb{F}}^{\mathrm{rel}}} & \Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\mathbb{F})) \\
(\phi, \theta) \downarrow & & \downarrow \Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\phi, \theta)) \\
\mathbb{F}' & \xrightarrow{\eta_{\mathbb{F}'}^{\mathrm{rel}}} & \Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\mathbb{F}'))
\end{array}$$

commutes in Psh^{\sqcup} . That is, we need an invertible icon $U : \phi_{\mathbb{F}'} \circ \phi \Rightarrow \eta_{\mathbb{X}}(\phi) \circ \phi_{\mathbb{F}}$, which exists by Theorem 4.7, and an invertible modification

$$\begin{array}{ccc}
& & \mathbb{F} \\
& \nearrow & \circlearrowleft \\
\mathbb{X}^{\mathrm{op}} & \xrightarrow{(\Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\mathbb{F}'))U^{\mathrm{op}}) \circ (\theta_{\mathbb{F}'} \circ \theta)} & \Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\theta)) \circ \theta_{\mathbb{F}} \\
& \searrow & \circlearrowright \\
& & \mathbb{F}'
\end{array}
\quad \text{Cat}^{\sqcup}.$$

$\Omega^{\mathrm{rel}}(\Xi^{\mathrm{rel}}(\mathbb{F}')) \circ (\eta_{\mathbb{X}'} \circ \phi)^{\mathrm{op}}$

The required natural transformations Γ_a are defined to be

$$\begin{array}{ccc} \mathbb{F}(a) & \xrightarrow{\lambda_{\mathbb{F}(a)}} & \mathbf{Set}_{/\|\mathbb{F}(a)\|} \\ \theta_a \downarrow & \nearrow \lambda_{\theta_a} & \downarrow \|\theta_a\|_* \\ \mathbb{F}'(\phi(a)) & \xrightarrow{\lambda_{\mathbb{F}'(\phi(a))}} & \mathbf{Set}_{/\|\mathbb{F}'(\phi(a))\|}. \end{array}$$

This establishes the equivalence $\mathbf{Psh}^{\sqcup} \simeq \frac{1}{2}\mu\mathbf{Cat}\text{-mod}$. Now apply Theorem 4.5. \square

The simplicial variant of Theorem 4.7 is an equivalence between 2-SegS and the category of \sqcup -semisimple bicategories with simple units. The modification of Theorem 4.8 is a compatible equivalence between 2-SegRelS and the category of \mathbf{Cat}^{\sqcup} -valued presheaves over \sqcup -semisimple bicategories with simple units.

Example. In the above language, the semi-simplicial set $\tilde{\mathbb{T}}_{\bullet}(M)$ of Section 2.5 defines a \sqcup -semisimple semibicategory $\mathbb{X}(M)$ with $\text{Ob}(\mathbb{X}(M)) = M$ and $\text{Hom}_{\mathbb{X}(M)}(a, b) = \mathbf{Set}_{/M_{a \rightarrow b}}$, where $M_{a \rightarrow b} \subset C^0([-\infty, \infty], M)$ is the subset of paths from a to b , with composition given by counting (suitably interpreted) pseudoholomorphic polygons in M with prescribed boundary conditions. The relative 2-Segal semi-simplicial morphism $\tilde{\mathbb{T}}_{\bullet}^{\tau}(M) \rightarrow \tilde{\mathbb{T}}_{\bullet}(M)$ of Theorem 2.8 becomes the presheaf $\mathbb{F}^{\tau} : \mathbb{X}(M)^{\text{op}} \rightarrow \mathbf{Cat}^{\sqcup}$ given by $\mathbb{F}^{\tau}(a) = \mathbf{Set}_{/a^{\tau} \sqcup M_{a \rightarrow \tau(a)}^{\tau}}$, where $a^{\tau} = a$ if $a \in M^{\tau}$ and $a^{\tau} = \emptyset$ otherwise and $M_{a \rightarrow \tau(a)}^{\tau} \subset M_{a \rightarrow \tau(a)}$ is the subset of real paths. Counts of real pseudoholomorphic n -gons, with $n \leq 4$, obeying one ordinary and two real boundary conditions determine the required functors $\mathbb{F}^{\tau}(b) \rightarrow \mathbb{F}^{\tau}(a)$. \triangleleft

REFERENCES

- [1] J. Bergner, A. Osorno, V. Ozornova, M. Rovelli, and C. Scheimbauer. 2-Segal sets and the Waldhausen construction. arXiv:1609.02853, 2016.
- [2] T. Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.
- [3] P. de Brito. Segal objects and the Grothendieck construction. arXiv:1605.00706, 2016.
- [4] H. Derksen and J. Weyman. Generalized quivers associated to reductive groups. *Colloq. Math.*, 94(2):151–173, 2002.
- [5] V. Drinfeld. On the notion of geometric realization. *Mosc. Math. J.*, 4(3):619–626, 782, 2004.
- [6] T. Dyckerhoff. Higher categorical aspects of Hall algebras. arXiv:1505.06940, 2015.
- [7] T. Dyckerhoff and M. Kapranov. Higher Segal spaces I. arXiv:1212.3563, 2012.
- [8] T. Dyckerhoff and M. Kapranov. Triangulated surfaces in triangulated categories. arXiv:1306.2545, 2013.
- [9] T. Dyckerhoff and M. Kapranov. Crossed simplicial groups and structured surfaces. In *Stacks and categories in geometry, topology, and algebra*, volume 643 of *Contemp. Math.*, pages 37–110. Amer. Math. Soc., Providence, RI, 2015.
- [10] N. Enomoto. A quiver construction of symmetric crystals. *Int. Math. Res. Not.*, 12:2200–2247, 2009.
- [11] H. Franzen. On cohomology rings of non-commutative Hilbert schemes and CoHa-modules. *Math. Res. Lett.*, 23(3):804–840, 2016.
- [12] I. Gálvez-Carrillo, J. Kock, and A. Tonks. Decomposition spaces, incidence algebras and Möbius inversion I: basic theory. arXiv:1512.07573, 2015.
- [13] J. Green. The characters of the finite general linear groups. *Trans. Amer. Math. Soc.*, 80:402–447, 1955.
- [14] G. Heuts and I. Moerdijk. Left fibrations and homotopy colimits. *Math. Z.*, 279(3-4):723–744, 2015.
- [15] J. Hornbostel and M. Schlichting. Localization in Hermitian K -theory of rings. *J. London Math. Soc. (2)*, 70(1):77–124, 2004.
- [16] A. Hubery. From triangulated categories to Lie algebras: a theorem of Peng and Xiao. In *Trends in representation theory of algebras and related topics*, volume 406 of *Contemp. Math.*, pages 51–66. Amer. Math. Soc., Providence, RI, 2006.
- [17] A. Joyal. The theory of quasi-categories I. In preparation.
- [18] A. Joyal and R. Street. The category of representations of the general linear groups over a finite field. *J. Algebra*, 176(3):908–946, 1995.

- [19] A. Joyal and M. Tierney. Quasi-categories vs Segal spaces. In *Categories in algebra, geometry and mathematical physics*, volume 431 of *Contemp. Math.*, pages 277–326. Amer. Math. Soc., Providence, RI, 2007.
- [20] D. Joyce. Configurations in abelian categories. II. Ringel-Hall algebras. *Adv. Math.*, 210(2):635–706, 2007.
- [21] R. Kashaev. Quantization of Teichmüller spaces and the quantum dilogarithm. *Lett. Math. Phys.*, 43(2):105–115, 1998.
- [22] D. Kazhdan and Y. Varshavskii. The Yoneda lemma for complete Segal spaces. *Funktsional. Anal. i Prilozhen.*, 48(2):3–38, 2014.
- [23] M. Kontsevich and Y. Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations. arXiv:0811.2435, 2008.
- [24] M. Kontsevich and Y. Soibelman. Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants. *Commun. Number Theory Phys.*, 5(2):231–352, 2011.
- [25] J. Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [26] G. Lusztig. Quivers, perverse sheaves, and quantized enveloping algebras. *J. Amer. Math. Soc.*, 4(2):365–421, 1991.
- [27] I. Macdonald. Polynomial functors and wreath products. *J. Pure Appl. Algebra*, 18(2):173–204, 1980.
- [28] V. Ostrik. Module categories, weak Hopf algebras and modular invariants. *Transform. Groups*, 8(2):177–206, 2003.
- [29] R. Penner. *Decorated Teichmüller theory*. QGM Master Class Series. European Mathematical Society (EMS), Zürich, 2012.
- [30] H.-G. Quebbemann, W. Scharlau, and M. Schulte. Quadratic and Hermitian forms in additive and abelian categories. *J. Algebra*, 59(2):264–289, 1979.
- [31] D. Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [32] C. Rezk. A model for the homotopy theory of homotopy theory. *Trans. Amer. Math. Soc.*, 353(3):973–1007 (electronic), 2001.
- [33] C. Ringel. Hall algebras. In *Topics in algebra, Part 1 (Warsaw, 1988)*, volume 26 of *Banach Center Publ.*, pages 433–447. PWN, Warsaw, 1990.
- [34] O. Schiffmann. Lectures on Hall algebras. In *Geometric methods in representation theory. II*, volume 24 of *Sémin. Congr.*, pages 1–141. Soc. Math. France, Paris, 2012.
- [35] M. Schlichting. Hermitian K -theory of exact categories. *J. K-Theory*, 5(1):105–165, 2010.
- [36] M. Schlichting. The Mayer-Vietoris principle for Grothendieck-Witt groups of schemes. *Invent. Math.*, 179(2):349–433, 2010.
- [37] G. Segal. Classifying spaces and spectral sequences. *Inst. Hautes Études Sci. Publ. Math.*, (34):105–112, 1968.
- [38] G. Segal. Categories and cohomology theories. *Topology*, 13:293–312, 1974.
- [39] J. Shapiro and D. Yao. Hermitian \mathcal{U} -theory of exact categories with duality functors. *J. Pure Appl. Algebra*, 109(3):323–330, 1996.
- [40] S. Shelley-Abrahamson. Hopf modules and representations of finite wreath products. *Algebr. Represent. Theory*, 20(1):123–145, 2017.
- [41] Y. Soibelman. Remarks on cohomological Hall algebras and their representations. In *Arbeitstagung Bonn 2013: In Memory of Friedrich Hirzebruch*, volume 319 of *Progress in Mathematics*, pages 355–385. Birkhäuser Basel, 2016.
- [42] M. Szczesny. Representations of quivers over \mathbb{F}_1 and Hall algebras. *Int. Math. Res. Not. IMRN*, (10):2377–2404, 2012.
- [43] R. Thomason and T. Trobaugh. Higher algebraic K -theory of schemes and of derived categories. In *The Grothendieck Festschrift, Vol. III*, volume 88 of *Progr. Math.*, pages 247–435. Birkhäuser Boston, Boston, MA, 1990.
- [44] M. van Leeuwen. An application of Hopf-algebra techniques to representations of finite classical groups. *J. Algebra*, 140(1):210–246, 1991.
- [45] T. Walde. Hall monoidal categories and categorical modules. arXiv:1611.08241, 2016.
- [46] F. Waldhausen. Algebraic K -theory of spaces. In *Algebraic and geometric topology (New Brunswick, N.J., 1983)*, volume 1126 of *Lecture Notes in Math.*, pages 318–419. Springer, Berlin, 1985.
- [47] M. Young. The Hall module of an exact category with duality. *J. Algebra*, 446:291–322, 2016.
- [48] M. Young. Representations of cohomological Hall algebras and Donaldson-Thomas theory with classical structure groups. arXiv:1603.05401, 2016.

- [49] A. Zelevinsky. *Representations of finite classical groups: A Hopf algebra approach*, volume 869 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1981.

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG
E-mail address: myoung@ims.cuhk.edu.hk